arXiv:1506.08011v2 [math.RT] 23 Jul 2015

Irreducible quantum group modules with finite dimensional weight spaces. II

Dennis Hasselstrøm Pedersen

Abstract

We classify the simple quantum group modules with finite dimensional weight spaces when the quantum parameter q is transcendental and the Lie algebra is not of type G_2. This is part 2 of the story. The first part being [Ped15a]. In [Ped15a] the classification is reduced to the classification of torsion free simple modules. In this paper we follow the procedures of [Mat00] to reduce the classification to the classification of infinite dimensional admissible simple highest weight modules. We then classify the infinite dimensional admissible simple highest weight modules and show among other things that they only exist for types A and C. Finally we complete the classification of simple torsion free modules for types A and C completing the classification of the simple torsion free modules.

Contents

1 Introduction **2**
1.1 Main results . . . 2
1.2 Acknowledgements . . . 3
1.3 Notation . . . 3
1.4 Basic definitions . . . 5

2 Reductions **6**

3 U_A calculations **7**

4 Ore localization and twists of localized modules **8**

5 Coherent families **19**

6 Classification of simple torsion free $U_q(\mathfrak{sl}_2)$-modules **30**

7 An example for $U_q(\mathfrak{sl}_3)$ **31**

8 Classification of admissible simple highest weight modules **34**
8.1 Preliminaries . . . 34
8.2 Rank 2 calculations . . . 36
8.3 Type A, D, E . . . 37
8.4 Quantum Shale-Weil representation . . . 41
8.5 Type B, C, F . . . 45

9 Classification of simple torsion free modules. Type A. 47

10 Classification of simple torsion free modules. Type C. 54

1 Introduction

This is part 2 of the classification of simple quantum group modules with finite dimensional weight spaces. In this paper we focus on the non root of unity case. Let $\mathfrak{g}$ be a simple Lie algebra. Let $q \in \mathbb{C}$ be a non root of unity and let U_q be the quantized enveloping algebra over $\mathbb{C}$ with q as the quantum parameter (defined below). We want to classify all simple weight modules for U_q with finite dimensional weight spaces. In the papers [Fer90] and [Mat00] this is done for $\mathfrak{g}$-modules. Fernando proves in [Fer90] that the classification of simple $\mathfrak{g}$ weight modules with finite dimensional weight spaces essentially boils down to classifying two classes of simple modules: Finite dimensional simple modules over a reductive Lie algebra and so called 'torsion free' simple modules over a simple Lie algebra. The classification of finite dimensional modules is well known in the classical case (as well as the quantum group case) so the remaining problem is to classify the torsion free simple modules. O. Mathieu classifies all torsion free $\mathfrak{g}$-modules in [Mat00]. The classification uses the concept of a $\mathfrak{g}$ coherent family which are huge $\mathfrak{g}$ modules with weight vectors for every possible weight, see [Mat00, Section 4]. Mathieu shows that every torsion free simple module is a submodule of a unique irreducible semisimple coherent family [Mat00, Proposition 4.8] and each of these irreducible semisimple coherent families contains an admissible simple highest weight module as well [Mat00, Proposition 6.2 ii)]. This reduces the classification to the classification of admissible simple highest weight modules. In this paper we will follow closely the methods described in [Mat00]. We will focus only on the case when q is not a root of unity. The root of unity case is studied in [Ped15a]. Some of the results of [Mat00] translate directly to the quantum group case but in several cases there are obstructions that need to be handled differently. In particular the case by case classification in types A and C is done differently. This is because our analog of $\mathcal{EXT}(L)$ given an admissible simple infinite dimensional module L is slightly different from the classical case see e.g. Section 7. The proof when reducing to types A and C in [Fer90] and [Mat00] uses some algebraic geometry to show that torsion free modules can only exist in types A and C. In this paper we show that infinite dimensional admissible simple highest weight modules only exist in types A and C and use this fact to show that torsion free modules can not exist for modules other than types A and C. For this we have to restrict to transcendental q. Specifically we use Theorem 8.1. If this theorem is true for a general non-root-of-unity q we can remove this restriction. The author is not aware of such a result in the litterature.

1.1 Main results

To classify simple weight modules with finite dimensional modules we follow the procedures of S. Fernando and O. Mathieu in [Fer90] and [Mat00]. The analog of the reduction done in [Fer90] is taken care of in the quantum group case in [Ped15a] so what remains is to classify the torsion free modules. We will first recall some results from [Ped15a] and [Ped15b] concerning the reduction and some formulas for commutating root vectors. This is recalled in Section 2 and Section 3. In Section 4 we do some prelimary calculations concerning Ore localization and certain 'twists' of modules necessary to define the 'Coherent families' of Section 5. Here we don't define the concept of a general coherent family but instead directly define the analog of coherent irreducible semisimple extensions $\mathcal{EXT}(L)$ of an admissible simple infinite dimensional module L. In analog with the classical case we show that for any admissible simple infinite dimensional module L, $\mathcal{EXT}(L)$ contains a submodule isomorphic to a simple highest weight module, see Theorem 5.12. We also prove a result in the other direction: If $\mathfrak{g}$ is such that there exists a simple infinite dimensional admissible module L then there exists a torsion free $U_q(\mathfrak{g})$-module, see Theorem 5.8. So the existence of torsion free modules over the quantized enveloping algebra of a specific $\mathfrak{g}$ is equivalent to the existence of an admissible infinite dimensional highest weight simple module over $U_q(\mathfrak{g})$. Using this we show that torsion free modules exist only for types A and C in the Sections 8.1, 8.2, 8.3, 8.4 and 8.5 where we also classify the admissible simple highest weight modules which are infinite dimensional. Finally in Section 9 and Section 10 we complete the classification in types A and C, respectively, by showing exactly which submodules of $\mathcal{EXT}(L(\lambda))$ are torsion free for a λ of a specific form see Theorem 9.8 and Theorem 10.7.

1.2 Acknowledgements

I would like to thank my advisor Henning H. Andersen for great supervision and many helpful comments and discussions and Jacob Greenstein for introducing me to this problem when I was visiting him at UC Riverside in the fall of 2013. The authors research was supported by the center of excellence grant 'Center for Quantum Geometry of Moduli Spaces' from the Danish National Research Foundation (DNRF95).

1.3 Notation

We will fix some notation: We denote by $\mathfrak{g}$ a fixed simple Lie algebra over the complex numbers $\mathbb{C}$. We assume $\mathfrak{g}$ is not of type G_2 to avoid unpleasant computations.

Fix a triangular decomposition of $\mathfrak{g}$: $\mathfrak{g} = \mathfrak{g}^- \oplus \mathfrak{h} \oplus \mathfrak{g}^+$: Let $\mathfrak{h}$ be a maximal toral subalgebra and let $\Phi \subset \mathfrak{h}^*$ be the roots of $\mathfrak{g}$ relative to $\mathfrak{h}$. Choose a simple system of roots $\Pi = \{\alpha_1, \dots, \alpha_n\} \subset \Phi$. Let Φ^+ (resp. Φ^-) be the positive (resp. negative) roots. Let $\mathfrak{g}^\pm$ be the positive and negative part of $\mathfrak{g}$ corresponding to the simple system Π. Let W be the Weyl group generated by the simple reflections $s_i := s_{\alpha_i}$. For a $w \in W$ let $l(w)$ be the length of W i.e. the smallest amount of simple reflections such that $w = s_{i_1} \cdots s_{i_{l(w)}}$. Let $(\cdot|\cdot)$ be a standard W-invariant bilinear form on $\mathfrak{h}^*$ and $\langle \alpha, \beta^\vee \rangle = \frac{2(\alpha|\beta)}{(\beta|\beta)}$. Since $(\cdot|\cdot)$ is standard we have $(\alpha|\alpha) = 2$ for any short root $\alpha \in \Phi$ and since $\mathfrak{g}$ is not of type G_2 we have $(\beta|\beta) = 4$ for any long root $\beta \in \Phi$. Let $Q = \operatorname{span}_{\mathbb{Z}} \{\alpha_1, \dots, \alpha_n\}$ denote the root lattice and $\Lambda = \operatorname{span}_{\mathbb{Z}} \{\omega_1, \dots, \omega_n\} \subset \mathfrak{h}^*$ the integral lattice where $\omega_i \in \mathfrak{h}^*$ is the fundamental weights defined by $(\omega_i|\alpha_j) = \delta_{ij}$.

Let $U_v = U_v(\mathfrak{g})$ be the corresponding quantized enveloping algebra defined over $\mathbb{Q}(v)$, see e.g. [Jan96] with generators $E_\alpha, F_\alpha, K_\alpha^{\pm 1}$, $\alpha \in \Pi$ and certain relations which can be found in chapter 4 of [Jan96]. We define $v_\alpha = v^{(\alpha|\alpha)/2}$ (i.e. $v_\alpha = v$ if α is a short root and $v_\alpha = v^2$ if α is a long root) and for $n \in \mathbb{Z}$, $[n]_v = \frac{v^n - v^{-n}}{v - v^{-1}}$. Let $[n]_\alpha := [n]_{v_\alpha} = \frac{v_\alpha^n - v_\alpha^{-n}}{v_\alpha - v_\alpha^{-1}}$. We omit the subscripts when it is clear from the context. For later use we also define the quantum binomial coefficients: For $r \in \mathbb{N}$ and $a \in \mathbb{Z}$:

$$\begin{bmatrix} a \\ r \end{bmatrix}_v = \frac{[a][a-1]\cdots[a-r+1]}{[r]!}$$

where $[r]! := [r][r-1]\cdots[2][1]$. Let $A = \mathbb{Z}[v, v^{-1}]$ and let U_A be Lusztigs A-form, i.e. the A subalgebra generated by the divided powers $E_\alpha^{(n)} := \frac{1}{[n]_\alpha !} E_\alpha^n$, $F_\alpha^{(n)} := \frac{1}{[n]_\alpha !} F_\alpha^n$ and $K_\alpha^{\pm 1}$, $\alpha \in \Pi$.

Let $q \in \mathbb{C}^* = \mathbb{C} \backslash \{0\}$ be a nonzero complex number that is not a root of unity and set $U_q = U_A \otimes_A \mathbb{C}_q$ where $\mathbb{C}_q$ is the A-algebra $\mathbb{C}$ where v is sent to q.

We have a triangular decomposition of Lusztigs A-form $U_A = U_A^- \otimes U_A^0 \otimes U_A^+$ with $U_A^- = \left\langle F_\alpha^{(n)} | \alpha \in \Pi, n \in \mathbb{N} \right\rangle \in U_A$, $U_A^+ = \left\langle E_\alpha^{(n)} | \alpha \in \Pi, n \in \mathbb{N} \right\rangle \in U_A$ and $U_A^0 = \left\langle K_\alpha^{\pm 1}, \begin{bmatrix} K_\alpha ; c \\ r \end{bmatrix} | \alpha \in \Pi, c \in \mathbb{Z}, r \in \mathbb{N} \right\rangle$ where

$$\begin{bmatrix} K_\alpha ; c \\ r \end{bmatrix} := \prod_{j=1}^{r} \frac{K_\alpha v_\alpha^{c-j+1} - K_\alpha^{-1} v_\alpha^{-c+j-1}}{v_\alpha^j - v_\alpha^{-j}}.$$

We have the corresponding triangular decomposition of U_q: $U_q = U_q^- \otimes U_q^0 \otimes U_q^+$ with $U_q^\pm = U_A^\pm \otimes_A \mathbb{C}_q$ and $U_q^0 = U_A^0 \otimes_A \mathbb{C}_q$.

For a $q \in \mathbb{C}^*$ define $\begin{bmatrix} a \\ r \end{bmatrix}_q$ as the image of $\begin{bmatrix} a \\ r \end{bmatrix}_v$ in $\mathbb{C}_q$. We will omit the subscript from the notation when it is clear from the context. $q_\beta \in \mathbb{C}$ and $[n]_\beta \in \mathbb{C}$ are defined as the image of $v_\beta \in A$ and $[n]_\beta \in A$, respectively abusing notation. Similarly, we will abuse notation and write $\begin{bmatrix} K_\alpha ; c \\ r \end{bmatrix}$ also for the image of $\begin{bmatrix} K_\alpha ; c \\ r \end{bmatrix} \in U_A$ in U_q. Define for $\mu \in Q$, $K_\mu = \prod_{i=1}^n K_{\alpha_i}^{a_i}$ if $\mu = \sum_{i=1}^n a_i \alpha_i$ with $a_i \in \mathbb{Z}$.

There is a braid group action on U_v which we will describe now. We use the definition from [Jan96, Chapter 8]. The definition is slightly different from the original in [Lus90, Theorem 3.1] (see [Jan96, Warning 8.14]). For each simple reflection s_i there is a braid operator that we will denote by T_{s_i} satisfying the following: $T_{s_i} : U_v \to U_v$ is a $\mathbb{Q}(v)$ automorphism. For $i \neq j \in \{1, \ldots, n\}$

$$\begin{aligned}
T_{s_i}(K_\mu) =& K_{s_i(\mu)} \\
T_{s_i}(E_{\alpha_i}) =& -F_{\alpha_i} K_{\alpha_i} \\
T_{s_i}(F_{\alpha_i}) =& -K_{\alpha_i}^{-1} E_{\alpha_i} \\
T_{s_i}(E_{\alpha_j}) =& \sum_{i=0}^{-\langle \alpha_j, \alpha_i^\vee \rangle} (-1)^i v_{\alpha_i}^{-i} E_{\alpha_i}^{(r-i)} E_{\alpha_j} E_{\alpha_i}^{(i)} \\
T_{s_i}(F_{\alpha_j}) =& \sum_{i=0}^{-\langle \alpha_j, \alpha_i^\vee \rangle} (-1)^i v_{\alpha_i}^{i} E_{\alpha_i}^{(i)} E_{\alpha_j} E_{\alpha_i}^{(r-i)}.
\end{aligned}$$

The inverse $T_{s_i}^{-1}$ is given by conjugating with the $\mathbb{Q}$-algebra anti-automorphism

Ψ from [Lus90, section 1.1] defined as follows:

$$\Psi(E_{\alpha_i}) = E_{\alpha_i}, \quad \Psi(F_{\alpha_i}) = F_{\alpha_i}, \quad \Psi(K_{\alpha_i}) = K_{\alpha_i}^{-1}, \quad \Psi(q) = q.$$

The braid operators T_{s_i} satisfy braid relations so we can define T_w for any $w \in W$: Choose a reduced expression of w: $w = s_{i_1} \cdots s_{i_n}$. Then $T_w = T_{s_{i_1}} \cdots T_{s_{i_n}}$ and T_w is independent of the chosen reduced expression, see e.g. [Lus90, Theorem 3.2]. We have $T_w(K_\mu) = K_{w(\mu)}$. The braid group operators restrict to automorphisms $U_A \to U_A$ and extend to automorphisms $U_q \to U_q$.

Let M be a U_q-module and $\lambda : U_q^0 \to \mathbb{C}$ a character (i.e. an algebra homomorphism into $\mathbb{C}$). Then

$$M_\lambda = \{m \in M | \forall u \in U_q^0, um = \lambda(u)m\}.$$

Let X denote the set of characters $U_q^0 \to \mathbb{C}$. Since $U_q^0 \cong \mathbb{C}[X_1^{\pm 1}, \dots, X_n^{\pm 1}]$ we can identify X with $(\mathbb{C}^*)^n$ by $U_q^0 \ni \lambda \mapsto (\lambda(K_{\alpha_1}), \dots, \lambda(K_{\alpha_n})) \in (\mathbb{C}^*)^n$.

1.4 Basic definitions

Definition 1.1 *Let* $\operatorname{wt} M$ *denote all the weights of* M*, i.e.* $\operatorname{wt} M = \{\lambda \in X | M_\lambda \neq 0\}$.

For $\mu \in \Lambda$ *and* $b \in \mathbb{C}^*$ *define the character* b^μ *by* $b^\mu(K_\alpha) = b^{(\mu|\alpha)}$, $\alpha \in \Pi$. *In particular for* $b = q$ *we get* $q^\mu(K_\alpha) = q^{(\mu|\alpha)}$. *We say that* M *only has integral weights if* $\lambda(K_\alpha) \in \pm q_\alpha^{\mathbb{Z}}$ *for all* $\lambda \in \operatorname{wt} M$, $\alpha \in \Pi$.

There is an action of W on X. For $\lambda \in X$ define $w\lambda$ by

$$(w\lambda)(u) = \lambda(T_{w^{-1}}(u))$$

Note that $wq^\mu = q^{w(\mu)}$.

Definition 1.2 *Let* M *be a* U_q*-module and* $w \in W$*. Define the twisted module* ${}^wM$ *by the following:*

As a vector space ${}^wM = M$ *but the action is given by twisting with* w^{-1}*: For* $m \in {}^wM$ *and* $u \in U_q$*:*

$$u \cdot m = T_{w^{-1}}(u)m.$$

We also define ${}^{\overline{w}}M$ *to be the inverse twist, i.e. for* $m \in {}^{\overline{w}}M$, $u \in U_q$*:*

$$u \cdot m = T_{w^{-1}}^{-1}(u)m.$$

Hence for any U_q*-module* ${}^{\overline{w}}({}^wM) = M = {}^w({}^{\overline{w}}M)$.

Note that $\operatorname{wt} {}^wM = w(\operatorname{wt} M)$ and that ${}^w({}^{w'}M) \cong {}^{ww'}M$ for $w, w' \in W$ with $l(ww') = l(w) + l(w')$ because the braid operators T_w satisfy braid relations. Also ${}^{\overline{w}}({}^{\overline{w'}}M) \cong {}^{\overline{w'w}}M$

Definition 1.3 *We define the category* $\mathcal{F} = \mathcal{F}(\mathfrak{g})$ *as the full subcategory of* $U_q - \mathrm{Mod}$ *such that for every* $M \in \mathcal{F}$ *we have*

1. M *is finitely generated as a* U_q*-module.*

2. $M = \bigoplus_{\lambda \in X} M_\lambda$ *and* $\dim M_\lambda < \infty$.

Note that the assignment $M \mapsto {}^w M$ is an endofunctor on $\mathcal{F}$ (in fact an auto-equivalence).

Let w_0 be the longest element in W and let $s_{i_1} \cdots s_{i_N}$ be a reduced expression of w_0. We define root vectors E_β and F_β for any $\beta \in \Phi^+$ by the following:

First of all set

$$\beta_j = s_{i_1} \cdots s_{i_{j-1}}(\alpha_{i_j}), \text{ for } i = 1, \ldots, N$$

Then $\Phi^+ = \{\beta_1, \ldots, \beta_N\}$. Set

$$E_{\beta_j} = T_{s_{i_1}} \cdots T_{s_{i_{j-1}}}(E_{\alpha_{i_j}})$$

and

$$F_{\beta_j} = T_{s_{i_1}} \cdots T_{s_{i_{j-1}}}(F_{\alpha_{i_j}})$$

In this way we have defined root vectors for each $\beta \in \Phi^+$. These root vectors depend on the reduced expression chosen for w_0 above. For a different reduced expression we might get different root vectors. It is a fact that if $\beta \in \Pi$ then the root vectors E_β and F_β defined above are the same as the generators with the same notation (cf. e.g. [Jan96, Proposition 8.20]) so the notation is not ambigious in this case. By "Let E_β be a root vector" we will just mean a root vector constructed as above for some reduced expression of w_0.

2 Reductions

We recall the following results from [Ped15a].

Proposition 2.1 *Let* β *be a positive root and* E_β, F_β *root vectors corresponding to* β. *Let* $M \in \mathcal{F}$. *The sets* $M^{[\beta]} = \{m \in M | \dim \langle E_\beta \rangle m < \infty\}$ *and* $M^{[-\beta]} = \{m \in M | \dim \langle F_\beta \rangle m < \infty\}$ *are submodules of* M *and independent of the chosen root vectors* E_β, F_β.

Proof. This is shown for E_β in Proposition 2.3 and Lemma 2.4 in [Ped15a] and the proofs are the same for F_β. □

Definition 2.2 *Let* $M \in \mathcal{F}$. *Let* $\beta \in \Phi$. M *is called* β*-free if* $M^{[\beta]} = 0$ *and* β*-finite if* $M^{[\beta]} = M$.

Suppose $L \in \mathcal{F}$ is a simple module and β a root. Then by Proposition 2.1 L is either β-finite or β-free.

Definition 2.3 *Let* $M \in \mathcal{F}$. *Define* $F_M = \{\beta \in \Phi | M \text{ is } \beta\text{-finite}\}$ *and* $T_M = \{\beta \in \Phi | M \text{ is } \beta\text{-free}\}$. *For later use we also define* $F_M^s := F_M \cap (-F_M)$ *and* $T_M^s := T_M \cap (-T_M)$ *to be the symmetrical parts of* F_M *and* T_M.

Note that $\Phi = F_L \cup T_L$ for a simple module L and this is a disjoint union.

Definition 2.4 *A module* M *is called torsion free if* $T_M = \Phi$.

Proposition 2.5 *Let* $L \in \mathcal{F}$ *be a simple module and* β *a root.* L *is* β*-free if and only if* $q^{\mathbb{N}\beta} \operatorname{wt} L \subset \operatorname{wt} L$.

Proof. Proposition 2.9 in [Ped15a]. $\square$

Proposition 2.6 *Let $L \in \mathcal{F}$ be a simple module. T_L and F_L are closed subsets of Q. That is, if $\beta, \gamma \in F_L$ (resp. $\beta, \gamma \in T_L$) and $\beta + \gamma \in \Phi$ then $\beta + \gamma \in F_L$ (resp. $\beta + \gamma \in T_L$).*

Proof. Proposition 2.10 and Proposition 2.11 in [Ped15a]. $\square$

Theorem 2.7 *Let $\lambda \in X$. There is a $1-1$ correspondence between simple U_q-modules with weight λ and simple $(U_q)_0$-modules with weight λ given by: For V a U_q-module, V_λ is the corresponding simple $(U_q)_0$-module.*

Proof. Theorem 2.21 in [Ped15a]. $\square$

Theorem 2.8 *Let $L \in \mathcal{F}$ be a simple $U_q(\mathfrak{g})$-module. Then there exists a $w \in W$, subalgebras $U_q(\mathfrak{p}), U_q(\mathfrak{l}), U_q(\mathfrak{u}), U_q(\mathfrak{u}^-)$ of U_q with $U_q = U_q(\mathfrak{u}^-)U_q(\mathfrak{p})$, $U_q(\mathfrak{p}) = U_q(\mathfrak{l})U_q(\mathfrak{u})$ and a simple $U_q(\mathfrak{l})$-module N such that ${}^wL$ is the unique simple quotient of $U_q \otimes_{U_q(\mathfrak{p})} N$ where N is considered a $U_q(\mathfrak{p})$-module with $U_q(\mathfrak{u})$ acting trivially.*

Furthermore there exists subalgebras U_{fr}, U_{fin} of $U_q(\mathfrak{l})$ such that $U_q(\mathfrak{l}) \cong U_{fr} \otimes U_{fin}$ and simple U_{fr} and U_{fin} modules X_{fr} and X_{fin} where X_{fin} is finite dimensional and X_{fr} is torsion free such that $N \cong X_{fin} \otimes X_{fr}$ as a $U_{fr} \otimes U_{fin}$-module.

U_{fr} is the quantized enveloping algebra of a semisimple Lie algebra $\mathfrak{t} = \mathfrak{t}_1 \oplus \cdots \oplus \mathfrak{t}_r$ where $\mathfrak{t}_1, \ldots, \mathfrak{t}_r$ are some simple Lie algebras. There exists simple torsion free $U_q(\mathfrak{t}_i)$-modules X_i, $i = 1, \ldots, r$ such that $X_{fr} \cong X_1 \otimes \cdots \otimes X_r$ as $U_q(\mathfrak{t}_1) \otimes \cdots \otimes U_q(\mathfrak{t}_r)$-modules.

Proof. Theorem 2.23 in [Ped15a]. $\square$

So the problem of classifying simple modules in $\mathcal{F}$ is reduced to the problem of classifying finite dimensional simple modules and classifying simple torsion free modules of $U_q(\mathfrak{t})$ where $\mathfrak{t}$ is a simple Lie algebra.

3 U_A calculations

In this section we recall from [Ped15b] some formulas for commuting root vectors with each other that will be used later on. Recall that $A = \mathbb{Z}[v, v^{-1}]$ where v is an indeterminate and U_A is the A-subspace of U_v generated by the divided powers $E_\alpha^{(n)}$ and $F_\alpha^{(n)}$, $n \in \mathbb{N}$.

Definition 3.1 *Let $x \in (U_q)_\mu$ and $y \in (U_q)_\gamma$ then we define*

$$[x, y]_q := xy - q^{-(\mu|\gamma)}yx$$

Theorem 3.2 *Suppose we have a reduced expression of $w_0 = s_{i_1} \cdots s_{i_N}$ and define root vectors $F_{\beta_1}, \ldots, F_{\beta_N}$. Let $i < j$. Let $A = \mathbb{Z}[q, q^{-1}]$ and let A' be the localization of A in $[2]$ if the Lie algebra contains any B_n, C_n or F_4 part. Then*

$$[F_{\beta_j}, F_{\beta_i}]_q = F_{\beta_j}F_{\beta_i} - q^{-(\beta_i|\beta_j)}F_{\beta_i}F_{\beta_j} \in \operatorname{span}_{A'}\left\{F_{\beta_{j-1}}^{a_{j-1}} \cdots F_{\beta_{i+1}}^{a_{i+1}}\right\}$$

Proof. [LS91, Proposition 5.5.2]. A proof following [DP93, Theorem 9.3] can also be found in [Ped15b, Theorem 2.9]. □

Definition 3.3 *Let $u \in U_A$ and $\beta \in \Phi^+$. Define* $\operatorname{ad}(F_\beta^i)(u) := [[\dots[u, F_\beta]_q \dots]_q, F_\beta]_q$ *and* $\widetilde{\operatorname{ad}}(F_\beta^i)(u) := [F_\beta, [\dots, [F_\beta, u]_q \dots]]_q$ *where the commutator is taken i times from the left and right respectively.*

Proposition 3.4 *Let $a \in \mathbb{N}$, $u \in (U_A)_\mu$ and $r = \langle \mu, \beta^\vee \rangle$. In U_A we have the identities*

$$\begin{aligned} uF_\beta^a &= \sum_{i=0}^{a} v_\beta^{(i-a)(r+i)} \begin{bmatrix} a \\ i \end{bmatrix}_\beta F_\beta^{a-i} \operatorname{ad}(F_\beta^i)(u) \\ &= \sum_{i=0}^{a} (-1)^i v_\beta^{a(r+i)-i} \begin{bmatrix} a \\ i \end{bmatrix}_\beta F_\beta^{a-i} \widetilde{\operatorname{ad}}(F_\beta^i)(u) \end{aligned}$$

Proof. Proposition 2.13 in [Ped15b]. □

Let $s_{i_1} \dots s_{i_N}$ be a reduced expression of w_0 and construct root vectors F_{β_i}, $i = 1, \dots, N$. In the next lemma F_{β_i} refers to the root vectors constructed as such. In particular we have an ordering of the root vectors.

Lemma 3.5 *Let $n \in \mathbb{N}$. Let $1 \leq j < k \leq N$.*
$\operatorname{ad}(F_{\beta_j}^i)(F_{\beta_k}^n) = 0$ *and* $\widetilde{\operatorname{ad}}(F_{\beta_k}^i)(F_{\beta_j}^n) = 0$ *for $i \gg 0$.*

Proof. Lemma 2.16 in [Ped15b]. □

4 Ore localization and twists of localized modules

In this section we present some results towards classifying simple torsion free modules following [Mat00].

We need the equivalent of Lemma 3.3 in [Mat00]. The proofs are essentially the same but for completeness we include most of the proofs here.

Definition 4.1 *A cone C is a finitely generated submonoid of the root lattice Q containing 0. If L is a simple module define the cone of L, $C(L)$, to be the submonoid of Q generated by T_L.*

Lemma 4.2 *Let $L \in \mathcal{F}$ be an infinite dimensional simple module. Then the group generated by the submonoid $C(L)$ is Q.*

Compare [Mat00] Lemma 3.1

Proof. First consider the case where $T_L \cap (-F_L) = \emptyset$. Then in this case we have $\Phi = T_L^s \cup F_L^s$. We claim that T_L^s and F_L^s correspond to different connected components of the Dynkin diagram: Suppose $\alpha \in F_L^s$ is a simple root and suppose $\alpha' \in \Pi$ is a simple root that is connected to α in the Dynkin diagram. So $\alpha + \alpha'$ is a root. There are two possibilities. Either $\alpha + \alpha' \in F_L$ or $\alpha + \alpha' \in T_L$. If $\alpha + \alpha' \in F_L$: Since F_L^s is symmetric we have $-\alpha \in F_L^s$ and since F_L is closed (Proposition 2.6) $\alpha' = \alpha + \alpha' + (-\alpha) \in F_L$. If $\alpha + \alpha' \in T_L$ and $\alpha' \in T_L$ then we get similarly $\alpha \in T_L$ which is a contradiction. So $\alpha' \in F_L$. We have shown that if $\alpha \in F_L$ then any simple root connected to α is in F_L also. So F_L and

T_L contains different connected components of the Dynkin diagram. Since L is simple and infinite we must have $\Phi = T_L^s$ and therefore $C(L) = Q$.

Next assume $T_L \cap (-F_L) \neq \emptyset$. By Lemma 4.16 in [Fer90] $P_L = T_L^s \cup F_L$ and $P_L^- = T_L \cup F_L^s$ are two opposite parabolic subsystems of Φ. So we have that $T_L \cap (-F_L)$ and $(-T_L) \cap F_L$ must be the roots corresponding to the nilradicals $\mathfrak{v}^{\pm}$ of two opposite parabolic subalgebras $\mathfrak{p}^{\pm}$ of $\mathfrak{g}$. Since we have $\mathfrak{g} = \mathfrak{v}^+ + \mathfrak{v}^- + [\mathfrak{v}^+, \mathfrak{v}^-]$ we get that $T_L \cap (-F_L)$ generates Q. Since $C(L)$ contains $T_L \cap (-F_L)$ it generates Q. $\square$

We define ρ and δ like in [Mat00, Section 3]:

Definition 4.3 *Let $x \geq 0$ be a real number. Define $\rho(x) = \operatorname{Card} B(x)$ where $B(x) = \{\mu \in Q | \sqrt{(\mu|\mu)} \leq x\}$*

Let M be a weight module with support lying in a single Q-coset, say $q^Q \lambda := \{q^\mu \lambda | \mu \in Q\}$. The density of M is $\delta(M) = \liminf_{x \to \infty} \rho(x)^{-1} \sum_{\mu \in B(x)} \dim M_{q^\mu \lambda}$

For a cone C we define $\delta(C) = \liminf_{x \to \infty} \rho(x)^{-1} \operatorname{Card}(C \cap B(x))$

Lemma 4.4 *There exists a real number $\varepsilon > 0$ such that $\delta(L) > \varepsilon$ for all infinite dimensional simple modules L.*

Proof. Note that since $q^{C(L)}\lambda \subset \operatorname{wt} L$ for all $\lambda \in \operatorname{wt} L$ we have $\delta(L) \geq \delta(C(L))$.

Since $C(L)$ is the cone generated by T_L and $T_L \subset \Phi$ (a finite set) there can only be finitely many different cones.

Since there are only finitely many different cones attached to infinite simple dimensional modules and since any cone C that generates Q has $\delta(C) > 0$ we conclude via Lemma 4.2 that there exists an $\varepsilon > 0$ such that $\delta(L) > \varepsilon$ for all infinite dimensional simple modules. $\square$

Definition 4.5 *A module $M \in \mathcal{F}$ is called admissible if its weights are contained in a single coset of X/q^Q and if the dimensions of the weight spaces are uniformly bounded. M is called admissible of degree d if d is the maximal dimension of the weight spaces in M.*

Of course all finite dimensional simple modules are admissible but the interesting admissible modules are the infinite dimensional simple ones. In particular simple torsion free modules are admissible. We show later that each infinite dimensional admissible simple module L gives rise to a 'coherent family' $\mathcal{EXT}(L)$ containing at least one torsion free module and at least one simple highest weight module that is admissible of the same degree.

Lemma 4.6 *Let $M \in \mathcal{F}$ be an admissible module. Then M has finite Jordan-Hölder length.*

Proof. The length of M is bounded by $A + \delta(M)/\varepsilon$ where $A = \sum_{\lambda \in Y} \dim M_\lambda$ and $Y = \{\nu \in X | \nu = \sigma q^\mu, |\langle \mu, \alpha^\vee \rangle| \leq 1, \sigma(K_\alpha) \in \{\pm 1\} \text{ for all } \alpha \in \Pi\}$. Check [Mat00, Lemma 3.3] for details. Here we use the fact that finite dimensional simple quantum group modules have the same character as their corresponding Lie algebra simple modules. This is proved for transcendental q in [Jan96, Theorem 5.15] and for general non-roots-of-unity in [APW91, Corollary 7.7]. $\square$

Lemma 4.7 *Let β be a positive root and let F_β be a corresponding root vector. The set $\{F_\beta^n | n \in \mathbb{N}\}$ is an Ore subset of U_q.*

Proof. A proof can be found in [And03] for β a simple root. If β is not simple then F_β is defined as $T_w(F_\alpha)$ for some $w \in W$ and some $\alpha \in \Pi$. Since $S := \{F_\alpha^n | n \in \mathbb{N}\}$ is an Ore subset of U_q we get for any $n \in \mathbb{N}$ and $u \in U_q$ that

$$F_\alpha^n U_q \cap uS \neq \emptyset.$$

Let $u' \in U_q$ and set $u = T_w^{-1}(u')$, then from the above

$$\emptyset \neq T_w(F_\alpha^n)T_w(U_q) \cap T_w(u)T_w(S) = F_\beta^n U_q \cap u' T_w(S).$$

Since $T_w(S) = \{F_\beta^n | n \in \mathbb{N}\}$ we have proved the lemma. $\square$

We denote the Ore localization of U_q in the above set by $U_{q(F_\beta)}$.

Lemma 4.8 *Let p be Laurent polynomial. If*

$$p(q^{r_1}, \ldots, q^{r_n}) = 0$$

for all $r_1, \ldots, r_n \in \mathbb{N}$ then $p = 0$.

Proof. If $n = 1$ we have a Laurent polynomial of one variable with infinitely many zero-points so $p = 0$. Let $n > 1$, then for constant $r_1 \in \mathbb{N}$, $p(q^{r_1}, -, \ldots, -)$ is a Laurent polynomial in $n - 1$ variables equal to zero in $(q^{r_2}, \ldots, q^{r_n})$ for all $r_2, \ldots, r_n \in \mathbb{N}$ so by induction $p(q^{r_1}, c_2, \ldots, c_n) = 0$ for all $c_2, \ldots, c_n$. Now for arbitrary $c_2, \ldots, c_n \in \mathbb{C}^*$ we get $p(-, c_2, \ldots, c_n)$ is a Laurent polynomial in one variable that is zero for all q^{r_1}, $r_1 \in \mathbb{N}$ hence $p(c_1, \ldots, c_n) = 0$ for all $c_1 \in \mathbb{C}^*$. $\square$

The next lemma is crucial for the rest of the results in this paper. We will use this result again and again.

Lemma 4.9 *Let $\beta \in \Phi^+$ and let F_β be a corresponding root vector. There exists automorphisms $\varphi_{F_\beta,b} : U_{q(F_\beta)} \to U_{q(F_\beta)}$ for each $b \in \mathbb{C}^*$ such that $\varphi_{F_\beta,q^i}(u) = F_\beta^{-i} u F_\beta^i$ for $i \in \mathbb{Z}$ and such that for $u \in U_{q(F_\beta)}$ the map $\mathbb{C}^* \to U_{q(F_\beta)}$, $b \mapsto \varphi_{F_\beta,b}(u)$ is of the form $b \mapsto p(b)$ for some Laurent polynomial $p \in U_{q(F_\beta)}[X, X^{-1}]$. Furthermore for $b, b' \in \mathbb{C}^*$, $\varphi_{F_\beta,b} \circ \varphi_{F_\beta,b'} = \varphi_{F_\beta,bb'}$.*

Proof. We can assume β is simple since if $F_\beta = T_w(F_{\alpha'})$ for some $\alpha' \in \Pi$ then we can just define the homomorphism on $T_w(E_\alpha), T_w(K_\alpha^{\pm 1}), T_w(F_\alpha)$ for $\alpha \in \Pi$ i.e. in this case we define $\varphi_{F_\beta,b}(u) = T_w(\varphi_{\alpha',b}(T_w^{-1}(u)))$ where we extend T_w to a homomorphism $T_w : U_{q(F_{\alpha'})} \to U_{q(F_\beta)}$ by $T_w(F_{\alpha'}^{-1}) = F_\beta^{-1}$.

So β is assumed simple. For $b \in \mathbb{C}^*$ define $b_\beta = b^{(\beta|\beta)/2}$ i.e. $b_\beta = b$ if β is short and $b_\beta = b^2$ when β is long. We will define the map on the generators $E_\alpha, K_\alpha, F_\alpha$ for $\alpha \in \Pi$. If $\alpha = \beta$ the map is defined as follows:

$$\begin{aligned}
\varphi_{F_\beta,b}(F_\beta^{\pm 1}) &= F_\beta^{\pm 1} \\
\varphi_{F_\beta,b}(K_\beta^{\pm 1}) &= b_\beta^{\mp 2} K_\beta^{\pm 1} \\
\varphi_{F_\beta,b}(E_\beta) &= E_\beta + F_\beta^{-1} \frac{(b_\beta - b_\beta^{-1})(q_\beta b_\beta^{-1} K_\beta - q_\beta^{-1} b_\beta K_\beta^{-1})}{(q_\beta - q_\beta^{-1})^2}.
\end{aligned}$$

Assume $\alpha \neq \beta$. Let $r = \langle \alpha, \beta^\vee \rangle$. Note that $\mathrm{ad}(F_\beta^{-r+1})(F_\alpha) = 0$ because this is one of the defining relations of U_q. We define the map as follows:

$$\begin{aligned}
\varphi_{F_\beta,b}(F_\alpha) =& \sum_{i=0}^{-r} b_\beta^{-r-i} q_\beta^{i(i+r)} \prod_{t=1}^{i} \frac{b_\beta q_\beta^{1-t} - b_\beta^{-1} q_\beta^{t-1}}{q_\beta^t - q_\beta^{-t}} F_\beta^{-i}\, \mathrm{ad}(F_\beta^i)(F_\alpha) \\
\varphi_{F_\beta,b}(K_\alpha) =& b_\beta^{-r} K_\alpha = b^{-(\alpha|\beta)} K_\alpha \\
\varphi_{F_\beta,b}(E_\alpha) =& E_\alpha.
\end{aligned}$$

Note that if $b = q^j$ for some $j \in \mathbb{Z}$ then $\prod_{t=1}^{i} \frac{b_\beta q_\beta^{1-t} - b_\beta^{-1} q_\beta^{t-1}}{q_\beta^t - q_\beta^{-t}} = \begin{bmatrix} j \\ i \end{bmatrix}_\beta$. Since the map $b \mapsto \varphi_{F_\beta,b}(u)$ is of the form $b \mapsto \sum_{i=1}^{r} p_i(b) u_i$ with p_i Laurent polynomial in b for each generator of U_q it is of this form for all $u \in U_q$. It's easy to check that $\varphi_{F_\beta,b}(u) = F_\beta^{-i} u F_\beta^i$ when $b = q^i$, $i \in \mathbb{N}$. So $\varphi_{F_\beta,b}$ satisfies the generating relations of U_q for $b = q^i$, $i \in \mathbb{N}$. By Lemma 4.8 $\varphi_{F_\beta,b}$ must satisfy the generating relations for all $b \in \mathbb{C}$.

Consider the last claim of the lemma: Let $u \in U_q$, then by the above $b \mapsto \varphi_{F_\beta,b}(u)$ is a Laurent polynomial and so $b \mapsto \varphi_{F_\beta,bb'}(u)$ and $\varphi_{F_\beta,b}(\varphi_{F_\beta,b'}(u))$ for a constant $b' \in \mathbb{C}^*$ is a Laurent polynomial as well. Now we know from above that for $b' = q^j$ for some $j \in \mathbb{Z}$ and $i \in \mathbb{Z}$:

$$\begin{aligned}
\varphi_{F_\beta,q^i} \circ \varphi_{F_\beta,b'}(u) =& F_\beta^{-i} F_\beta^{-j} u F_\beta^j F_\beta^i \\
=& F_\beta^{-i-j} u F_\beta^{i+j} \\
=& \varphi_{F_\beta,q^i q^j}(u)
\end{aligned}$$

So $\varphi_{F_\beta,b}(\varphi_{F_\beta,q^j}(u)) = \varphi_{F_\beta,bq^j}(u)$ for all $b \in \mathbb{C}^*$ since both sides are Laurent polynomials in b and they are equal in infinitely many points. In the same way we get the result for all $b' \in \mathbb{C}$. ☐

Note that if β is long then the above automorphism is a Laurent polynomial in b^2. So if $b_1^2 = b_2^2$ for $b_1, b_2 \in \mathbb{C}^*$ then $\varphi_{F_\beta,b_1} = \varphi_{F_\beta,b_2}$. We could have defined another automorphism $\varphi'_{F_\beta,b} := \varphi_{F_\beta,b^{2/(\beta|\beta)}}$ and proved the lemma above with the modification that $\varphi'_{F_\beta,q_\beta^i}(u) = F_\beta^{-i} u F_\beta^i$. The author has chosen the first option to avoid having to write the β in q_β all the time in results like Lemma 4.12 and Corollary 4.20. On the other hand this choice means that we have to take some squareroots sometimes when doing concrete calculations involving long roots see e.g the proof of Lemma 5.11. The choice of squareroot doesn't matter by the above.

We can use the formulas in Section 3 to find the value of $\varphi_{F_\beta,b}(F_{\beta'})$ and $\varphi_{F_\beta,b}(E_{\beta'})$ for general root vectors F_β, $F_{\beta'}$ and $E_{\beta'}$, $\beta, \beta' \in \Phi^+$.

Proposition 4.10 *Let $s_{i_1} \dots s_{i_N}$ be a reduced expression of w_0 and define root vectors $F_{\beta_1}, \dots, F_{\beta_N}$ and $E_{\beta_1}, \dots, E_{\beta_N}$ using this expression (i.e. $F_{\beta_j} = T_{s_{i_1}} \dots T_{s_{i_{j-1}}}(F_{\alpha_{i_j}})$*

and $E_{\beta_j} = T_{s_{i_1}} \dots T_{s_{i_{j-1}}}(E_{\alpha_{i_j}})$*). Let* $1 \leq j < k \leq N$ *and set* $r = \langle \beta_k, \beta_j^\vee \rangle$.

$$\begin{aligned}
\varphi_{F_{\beta_j},b}(F_{\beta_k}^n) =& \sum_{i\geq 0} q_{\beta_j}^{i(nr+i)} b_{\beta_j}^{-nr-i} \prod_{t=1}^{i} \frac{q_{\beta_j}^{1-t} b_{\beta_j} - q_{\beta_j}^{t-1} b_{\beta_j}^{-1}}{q_{\beta_j}^{t} - q_{\beta_j}^{-t}} F_{\beta_j}^{-i} \operatorname{ad}(F_{\beta_j}^i)(F_{\beta_k}^n)\\
\varphi_{F_{\beta_k},b}(F_{\beta_j}^n) =& \sum_{i\geq 0} (-1)^i q_{\beta_k}^{-i} b_{\beta_k}^{nr+i} \prod_{t=1}^{i} \frac{q_{\beta_k}^{1-t} b_{\beta_k} - q_{\beta_k}^{t-1} b_{\beta_k}^{-1}}{q_{\beta_k}^{t} - q_{\beta_k}^{-t}} F_{\beta_k}^{-i} \widetilde{\operatorname{ad}}(F_{\beta_k}^i)(F_{\beta_j}^n)\\
\varphi_{F_{\beta_j},b}(E_{\beta_k}) =& \sum_{i\geq 0} b_{\beta_j}^{-i} \prod_{t=1}^{i} \frac{q_{\beta_j}^{1-t} b_{\beta_j} - q_{\beta_j}^{t-1} b_{\beta_j}^{-1}}{q_{\beta_j}^{t} - q_{\beta_j}^{-t}} F_{\beta_j}^{-i} u_i\\
\varphi_{F_{\beta_k},b}(E_{\beta_j}) =& \sum_{i\geq 0} b_{\beta_k}^{i} \prod_{t=1}^{i} \frac{q_{\beta_k}^{1-t} b_{\beta_k} - q_{\beta_k}^{t-1} b_{\beta_k}^{-1}}{q_{\beta_k}^{t} - q_{\beta_k}^{-t}} F_{\beta_k}^{-i} \widetilde{u}_i
\end{aligned}$$

for some $u_i, \widetilde{u}_i \in U_q$ *(independent of* b*) such that* $u_i = \widetilde{u}_i = 0$ *for* $i \gg 0$*. In particular for any* $j, k \in \{1, \dots, N\}$*:*

$$\begin{aligned}
\varphi_{F_{\beta_j},-1}(F_{\beta_k}) &= (-1)^{(\beta_j|\beta_k)} F_{\beta_k}\\
\varphi_{F_{\beta_j},-1}(E_{\beta_k}) &= E_{\beta_k}.
\end{aligned}$$

Note that the sums are finite because of Lemma 3.5.

Proof. By Proposition 3.4 we have for any $a \in \mathbb{N}$

$$\begin{aligned}
F_{\beta_k}^n F_{\beta_j}^a =& \sum_{i=0}^{a} q_{\beta_j}^{(i-a)(nr+i)} \begin{bmatrix} a \\ i \end{bmatrix}_{\beta_j} F_{\beta_j}^{a-i} \operatorname{ad}(F_{\beta_j}^i)(F_{\beta_k}^n)\\
=& \sum_{i=0}^{\infty} q_{\beta_j}^{i(nr+i)} q_{\beta_j}^{-a(nr+i)} \prod_{t=1}^{i} \frac{q_{\beta_j}^{1-t} q_{\beta_j}^{a} - q_{\beta_j}^{t-1} q_{\beta_j}^{-a}}{q_{\beta_j}^{t} - q_{\beta_j}^{-t}} F_{\beta_j}^{a-i} \operatorname{ad}(F_{\beta_j}^i)(F_{\beta_k}^n).
\end{aligned}$$

Here we use the fact that $\left[\begin{smallmatrix} a \\ i \end{smallmatrix}\right]_{\beta_j} = 0$ for $i > a$. So

$$F_{\beta_j}^{-a} F_{\beta_k}^n F_{\beta_j}^a = \sum_{i\geq 0} q_{\beta_j}^{i(nr+i)} q_{\beta_j}^{-a(nr+i)} \prod_{t=1}^{i} \frac{q_{\beta_j}^{1-t} q_{\beta_j}^{a} - q_{\beta_j}^{t-1} q_{\beta_j}^{-a}}{q_{\beta_j}^{t} - q_{\beta_j}^{-t}} F_{\beta_j}^{-i} \operatorname{ad}(F_{\beta_j}^i)(F_{\beta_k}^n).$$

Now using the fact that $\varphi_{F_{\beta_j},q^a}(F_{\beta_k}^n) = F_{\beta_j}^{-a} F_{\beta_k}^n F_{\beta_j}^a$, the fact that $\varphi_{F_{\beta_j},b}(F_{\beta_k}^n)$ is Laurent polynomial and Lemma 4.8 we get the first identity. The second identity is shown similarly by using the second identity in Proposition 3.4.

To prove the last two identities we need to calculate $F_{\beta_j}^{-a} E_{\beta_k}^n F_{\beta_j}^a$ (resp. $F_{\beta_k}^{-a} E_{\beta_j}^n F_{\beta_k}^a$) for any $a \in \mathbb{N}$. Let $w = s_{i_1} \cdots s_{i_{j-1}}$ and $w' = s_{i_{j+1}} \cdots s_{i_{k-1}}$. Then $E_{\beta_j} = T_w(E_{\alpha_{i_j}})$ and $F_{\beta_k} = T_w T_{s_{i_j}} T_{w'}(F_{\alpha_{i_k}})$.

$$\begin{aligned}
E_{\beta_j} F_{\beta_k}^a =& T_w \left(E_{\alpha_{i_j}} T_{s_{i_j}} T_{w'}(F_{\alpha_{i_k}}^a) \right)\\
=& T_w T_{s_{i_j}} \left(-K_{\alpha_{i_j}}^{-1} F_{\alpha_{i_j}} T_{w'}(F_{\alpha_{i_k}}^a) \right).
\end{aligned}$$

Expand $s_{i_j} \cdots s_{i_N}$ from the right to a reduced expression $s_{i_j} \cdots s_{i_N} s_{m_1} \cdots s_{m_{j-1}}$ of w_0. Do the same with $s_{i_{j+1}} \cdots s_{i_N} s_{m_1} \cdots s_{m_{j-1}}$ to get a reduced expression $s_{i_{j+1}} \cdots s_{i_N} s_{m_1} \cdots s_{m_j}$. We claim that if we use the reduced expression

$s_{i_{j+1}}\cdots s_{i_N}s_{m_1}\cdots s_{m_j}$ to construct roots $\beta'_1\dots,\beta'_N$ and root vectors $F'_{\beta'_j}$ then $F'_{\beta'_N} = T_{s_{i_{j+1}}}\cdots T_{s_{i_N}}T_{s_{m_1}}\cdots T_{s_{m_{j-1}}}(F_{\alpha_{m_j}}) = F_{\alpha_{i_j}}$. This is easy to see since β'_N is positive but $s_{i_j}\beta'_N = w_0(\alpha_{m_j}) < 0$. We have $T_{w'}(F^a_{\alpha_{i_k}}) = F^a_{\beta'_{k-j}}$. Since $k-j<N$ we can use what we just calculated above: (set $d=k-j$)

$$F'^{-a}_{\beta'_d}F'_{\beta'_N}F'^{a}_{\beta'_d} = \sum_{i\geq 0} q_{\beta'_d}^{i(r+i)} q_{\beta'_d}^{-a(r+i)} \prod_{t=1}^{i} \frac{q_{\beta'_d}^{1-t}q_{\beta'_d}^{a} - q_{\beta'_d}^{t-1}q_{\beta'_d}^{-a}}{q_{\beta'_d}^{t} - q_{\beta'_d}^{-t}} F'^{-i}_{\beta'_d}\operatorname{ad}(F'^{i}_{\beta'_d})(F'_{\beta'_N}).$$

so

$$F^{-a}_{\beta_k}E_{\beta_j}F^{a}_{\beta_k} = K_{\beta_j}T_wT_{s_{i_j}}\left(\sum_{i\geq 0} q_{\beta'_d}^{i(r+i)} q_{\beta'_d}^{-ai} \prod_{t=1}^{i} \frac{q_{\beta'_d}^{1-t}q_{\beta'_d}^{a} - q_{\beta'_d}^{t-1}q_{\beta'_d}^{-a}}{q_{\beta'_d}^{t} - q_{\beta'_d}^{-t}} F'^{-i}_{\beta'_d}\operatorname{ad}(F'^{i}_{\beta'_d})(F'_{\beta'_N})\right).$$

This shows the third identity. The fourth is shown similarly.

Setting $b=-1$ in the above formulas we get the last claim of the proposition. □

Definition 4.11 *Let M be a $U_{q(F_\beta)}$-module. We define a new module $\varphi_{F_\beta,b}.M$ (with elements $\varphi_{F_\beta,b}.m$, $m\in M$) where the module structure is given by composing with the above automorphism $\varphi_{F_\beta,b}$. – i.e. $u\varphi_{F_\beta,b}.m = \varphi_{F_\beta,b}.\varphi_{F_\beta,b}(u)m$ for all $u\in U_{q(F_\beta)}$, $m\in M$.*

Note that $\operatorname{wt}\varphi_{F_\beta,b}.M = b^{-\beta}\operatorname{wt}M$ where $b^{-\beta}$ is the character such that $b^{-\beta}(K_\alpha) = b^{-(\alpha|\beta)}$ for $\alpha\in\Pi$.

The homomorphisms from Lemma 4.9 preserve degree so we can restrict to $(U_{q(F_\beta)})_0$ which we will do in the next lemma. The twist of a $(U_{q(F_\beta)})_0$-module is defined in the same way as the definition above. It is an important fact of these twists that they do not neccesarily preserve simplicity of U_q-modules: If L is a $U_{q(F_\beta)}$-module that is simple as a U_q-module then $\varphi_{F_\beta,b}.L$ can be nonsimple as a U_q-module for some $b\in\mathbb{C}^*$, see e.g. Lemma 4.23.

Lemma 4.12 *Let M be a $U_{q(F_\beta)}$-module. Let $i\in\mathbb{Z}$. Then*

$$\varphi_{F_\beta,q^i}.M\cong M$$

as $U_{q(F_\beta)}$-modules. Furthermore for $\lambda\in\operatorname{wt}M$ we have an isomorphism of $(U_{q(F_\beta)})_0$-modules:

$$\varphi_{F_\beta,q^i}.M_\lambda\cong M_{q^{-i\beta}\lambda}.$$

Proof. The isomorphism in both cases is given by $\varphi_{F_\beta,q^i}.m\mapsto F^i_\beta m$, $\varphi_{F_\beta,q^i}.M\to M$. The inverse is given by multiplying by F^{-i}_β. By Lemma 4.9: For $u\in U_{q(F_\beta)}$, $m\in M$; $\varphi_{F_\beta,q^i}(u) = F^{-i}_\beta uF^i_\beta$ so $u\varphi_{F_\beta,q^i}.m = \varphi_{F_\beta,q^i}.F^{-i}_\beta uF^i_\beta m\mapsto F^i_\beta F^{-i}_\beta uF^i_\beta m = uF^i_\beta m$. Thus the given map is a homomorphism. □

Definition 4.13 *Let $\Sigma\subset\Phi^+$. Then Σ is called a set of commuting roots if there exists an ordering of the roots in Σ; $\Sigma=\{\beta_1,\dots,\beta_s\}$ such that for some reduced expression of w_0 and corresponding construction of the root vectors F_β we have: $[F_{\beta_j},F_{\beta_i}]_q = 0$ for $1\leq i<j\leq s$.*

For any subset $I\subset\Pi$, let Q_I be the subgroup of Q generated by I, Φ_I the root system generated by I , $\Phi_I^+ = \Phi^+\cap\Phi_I$ and $\Phi_I^- = -\Phi_I^+$.

The following three lemmas have exactly the same proofs as their counterparts ([Ped15a, Lemma 5.6], [Ped15a, Lemma 5.22] and [Ped15a, Lemma 5.23]) in the root of unity case in [Ped15a]. We include the proofs here as well for completeness.

We have the following equivalent of Lemma 4.1 in [Mat00]:

Lemma 4.14 *1. Let $I \subset \Pi$ and let $\alpha \in I$. There exists a set of commuting roots $\Sigma' \subset \Phi_I^+$ with $\alpha \in \Sigma'$ such that Σ' is a basis of Q_I.*

2. Let J, F be subsets of Π with $F \neq \Pi$. Let $\Sigma' \subset \Phi_J^+ \backslash \Phi_{J\cap F}^+$ be a set of commuting roots which is a basis of Q_J. There exists a set of commuting roots Σ which is a basis of Q such that $\Sigma' \subset \Sigma \subset \Phi^+ \backslash \Phi_F^+$

Proof. The first part of the proof is just combinatorics of the root system so it is identical to the first part of the proof of Lemma 4.1 in [Mat00]: Let us first prove assertion 2.: If J is empty we can choose $\alpha \in \Pi\backslash F$ and replace J and Σ' by $\{\alpha\}$. So assume from now on that $J \neq \emptyset$. Set $J' = J\backslash F$, $p = |J'|$, $q = |J|$. Let $J_1, \dots, J_k$ be the connected components of J and set $J'_i = J' \cap J_i$, $F_i = F \cap J_i$, and $\Sigma'_i = \Sigma \cap \Phi_{J_i}$, for any $1 \leq i \leq k$. Since $\Sigma' \subset \Phi_J$ is a basis of Q_J, each Σ'_i is a basis of Q_{J_i}. Since Σ'_i lies in $\Phi^+_{J_i}\backslash\Phi^+_{F_i}$, the set $J'_i = J_i\backslash F_i$ is not empty. Hence J' meets every connected component of J. Therefore we can write $J = \{\alpha_1, \dots, \alpha_q\}$ in such a way that $J' = \{\alpha_1, \dots, \alpha_p\}$ and, for any s with $p + 1 \leq s \leq q$, α_s is connected to α_i for some $i < s$. Since Π is connected we can write $\Pi\backslash J = \{\alpha_{q+1}, \dots, \alpha_n\}$ in such a way that for any $s \geq q + 1$, α_s is connected to α_i for some i with $1 \leq i < s$. So $\Pi = \{\alpha_1, \dots, \alpha_n\}$ such that for $s > p$ we have that α_s is connected to some α_i with $1 \leq i < s$.

Let $\Sigma' = \{\beta_1, \dots, \beta_q\}$. We will define $\beta_{q+1}, \dots, \beta_l$ inductively such that for each $s \geq q$, $\{\beta_1, \dots, \beta_s\}$ is a commuting set of roots which is a basis of $\Phi_{\{\alpha_1,\dots,\alpha_s\}}$. So assume we have defined $\beta_1, \dots, \beta_s$. Let w_s be the longest word in $s_{\alpha_1}, \dots, s_{\alpha_s}$ and let w_{s+1} be the longest word in $s_{\alpha_1}, \dots, s_{\alpha_{s+1}}$. Choose a reduced expression of w_s such that the corresponding root vectors $\{F_{\beta_k}\}_{k=1}^s$ satisfies $[F_{\beta_j}, F_{\beta_i}]_q = 0$ for $i < j$. Choose a reduced expression of $w_{s+1} = w_s w'$ starting with the above reduced expression of w_s. Let N_s be the length of w_s and N_{s+1} be the length of w_{s+1}. So we get an ordering of the roots generated by $\{\alpha_1, \dots, \alpha_{s+1}\}$: $\Phi^+_{\{\alpha_1,\dots,\alpha_{s+1}\}} = \{\gamma_1, \dots, \gamma_{N_s}, \gamma_{N_s+1}, \dots, \gamma_{N_{s+1}}\}$ with $\Phi^+_{\{\alpha_1,\dots,\alpha_s\}} = \{\gamma_1, \dots, \gamma_{N_s}\}$. Consider $\gamma_{N_s+1} = w_s(\alpha_{s+1})$. Since w_s only consists of the simple reflections corresponding to $\alpha_1, \dots, \alpha_s$ we must have that $\gamma_{N_s+1} = \alpha_{s+1} + \sum_{i=1}^s m_i\alpha_i$ for some coefficients $m_i \in \mathbb{N}$. So $\{\beta_1, \dots, \beta_s, \gamma_{N_s+1}\}$ is a basis of $\Phi_{\{\alpha_1,\dots,\alpha_{s+1}\}}$. From Theorem 3.2 we get for $1 \leq i \leq s$

$$[F_{\gamma_{N_s+1}}, F_{\beta_i}]_q \in \operatorname{span}_{\mathbb{C}} \left\{F_{\gamma_{N_s}}^{a_{N_s}} \cdots F_{\gamma_2}^{a_2} | a_i \in \mathbb{N}\right\}$$

But since $\{\gamma_1, \dots, \gamma_{N_s}\} = \Phi^+_{\{\alpha_1,\dots,\alpha_s\}}$ and since $\gamma_{N_s+1} = \alpha_{s+1} + \sum_{i=1}^s m_i\alpha_i$ we get $[F_{\gamma_{N_s+1}}, F_{\beta_i}]_q = 0$.

All that is left is to show that $\gamma_{N_s+1} \notin \Phi_F$. By the above we must have that α_{s+1} is connected to some $\alpha_i \in J'$. We will show that the coefficient of α_i in γ_{N_s+1} is nonzero. Otherwise $(\gamma_{N_s+1}|\alpha_i) < 0$ and so $\gamma_{N_s+1} + \alpha_i \in \Phi_{\{\alpha_1,\dots,\alpha_{s+1}\}}$ and by Theorem 1 in [Pap94], $\gamma_{N_s+1} + \alpha_i = \gamma_j$ for some $1 < j \leq s$. This is impossible since $\gamma_{N_s+1} + \alpha_i \notin \Phi_{\{\alpha_1,\dots,\alpha_s\}}$. So we can set $\beta_{s+1} = \gamma_{N_s+1}$ and the induction step is finished.

To prove assertion 1. it can be assumed that $I = \Pi$. Thus assertion 1. follows from assertion 2. with $J = \{\alpha\}$ and $F = \emptyset$. □

Lemma 4.15 *Let $L \in \mathcal{F}$ be a simple module. Then there exists a $w \in W$ such that $w(F_L \backslash F_L^s) \subset \Phi^+$ and $w(T_L \backslash T_L^s) \subset \Phi^-$.*

Proof. Since L is simple we have $\Phi = F_L \cup T_L$. By Proposition 2.6 F_L and T_L are closed subsets. Then Lemma 4.16 in [Fer90] tells us that there exists a basis B of the root system Φ such that the antisymmetrical part of F_L is contained in the positive roots Φ_B^+ corresponding to the basis B and the antisymmetrical part of T_L is contained in the negative roots Φ_B^- corresponding to the basis. Since all bases of a root system are W-conjugate the claim follows. $\square$

Lemma 4.16 *Let L be an infinite dimensional admissible simple module. Let $w \in W$ be such that $w(F_L \backslash F_L^s) \subset \Phi^+$. Let $\alpha \in \Pi$ be such that $-\alpha \in w(T_L)$ (such an α always exists). Then there exists a commuting set of roots Σ with $\alpha \in \Sigma$ which is a basis of Q such that $-\Sigma \subset w(T_L)$.*

Proof. Set $L' = {}^wL$. Since $w(T_L) = T_{^wL} = T_{L'}$ we will just work with L'. Then $F_{L'} \backslash F_{L'}^s \subset \Phi^+$.

Note that it is always possible to choose a simple root $\alpha \in -T_{L'}$ since L' is infinite dimensional: If this was not possible we would have $\Phi^- \subset F_{L'}$. But since $F_{L'} \backslash F_{L'}^s \subset \Phi^+$ this would imply $F_L = \Phi$.

Set $F = F_{L'}^s \cap \Pi$. Since L' is infinite dimensional $F \neq \Pi$. By Lemma 4.14 2. applied with $J = \{\alpha\} = \Sigma'$ there exists a commuting set of roots Σ that is a basis of Q such that $\Sigma \subset \Phi^+ \backslash \Phi_F^+$. Since $F_{L'} \backslash F_{L'}^s \subset \Phi^+$ we have $\Phi^- = T_{L'}^- \cup (F_{L'}^s)^-$. To show $-\Sigma \subset T_{L'}$ we show $\left(\Phi^- \backslash \Phi_F^-\right) \cap F_{L'}^s = \emptyset$ or equivalently $(F_{L'}^s)^- \subset \Phi_F^-$.

Assume $\beta \in F_{L'}^s \cap \Phi^+$, $\beta = \sum_{\alpha \in \Pi} a_\alpha \alpha$, $a_\alpha \in \mathbb{N}$. The height of β is the sum $\sum_{\alpha \in \Pi} a_\alpha$. We will show by induction on the height of β that $-\beta \in \Phi_F^-$. If the height of β is 1 then β is a simple root and so $\beta \in F$. Clearly $-\beta \in \Phi_F^-$ in this case. Assume the height of β is greater than 1. Let $\alpha' \in \Pi$ be a simple root such that $\beta - \alpha'$ is a root. There are two possibilities: $-\alpha' \in T_{L'}$ or $\pm\alpha' \in F_{L'}^s$.

In the first case where $-\alpha' \in T_{L'}$ we must have $-\beta + \alpha' \in F_{L'}^s$ since if $-\beta + \alpha' \in T_L$ then $-\beta = (-\beta + \alpha') - \alpha' \in T_{L'}$. So $\beta - \alpha' \in F_{L'}^s$ and $\beta \in F_{L'}^s$. Since $F_{L'}$ is closed (Proposition 2.6) we get $-\alpha' = (\beta - \alpha') - \beta \in F_L$ which is a contradiction. So the first case ($-\alpha' \in T_{L'}$) is impossible.

In the second case since $F_{L'}$ is closed we get $\pm(\beta - \alpha') \in F_{L'}$ i.e. $\beta - \alpha' \in F_{L'}^s$. By the induction $-(\beta - \alpha') \in \Phi_F^-$ and since $-\beta = -(\beta - \alpha') - \alpha'$ we are done.$\square$

Proposition 4.17 *Let $\Sigma = \{\beta_1, \dots, \beta_r\}$ be a set of commuting roots. The set $\{q^a F_{\beta_1}^{a_1} \cdots F_{\beta_r}^{a_r} | a_i \in \mathbb{N}, a \in \mathbb{Z}\}$ is an Ore subset of U_q.*

Proof. We will prove it by induction over r. $r = 1$ is Lemma 4.7.

Let $S_r = \{q^a F_{\beta_1}^{a_1} \cdots F_{\beta_r}^{a_r} | a_i \in \mathbb{N}, a \in \mathbb{Z}\}$. Let $a_1, \dots, a_r \in \mathbb{N}$, $a \in \mathbb{Z}$ and $u \in U_q$, then we need to show that

$$q^a F_{\beta_1}^{a_1} \cdots F_{\beta_r}^{a_r} U_q \cap u S_r \neq \emptyset. \tag{1}$$

and

$$U_q q^a F_{\beta_1}^{a_1} \cdots F_{\beta_r}^{a_r} \cap S_r u \neq \emptyset. \tag{2}$$

By Lemma 4.7 there exists $\widetilde{u} \in U_q$ and $b \in \mathbb{N}$ such that

$$F_{\beta_r}^{a_r} \widetilde{u} = u F_{\beta_r}^b. \tag{3}$$

By induction

$$q^a F_{\beta_1}^{a_1} \cdots F_{\beta_{r-1}}^{a_{r-1}} U_q \cap \widetilde{u} S_{r-1} \neq \emptyset$$

so

$$q^a F_{\beta_r}^{a_r} F_{\beta_1}^{a_1} \cdots F_{\beta_{r-1}}^{a_{r-1}} U_q \cap F_{\beta_r}^{a_r} \widetilde{u} S_{r-1} \neq \emptyset$$

Since Σ is a set of commuting roots $F_{\beta_r}^{a_r} F_{\beta_1}^{a_1} \cdots F_{\beta_{r-1}}^{a_{r-1}} = q^k F_{\beta_1}^{a_1} \cdots F_{\beta_{r-1}}^{a_{r-1}} F_{\beta_r}^{a_r}$ for some $k \in \mathbb{Z}$. Using this and (3) we get

$$\emptyset \neq q^{a+k} F_{\beta_1}^{a_1} \cdots F_{\beta_r}^{a_r} U_q \cap u F_{\beta_r}^{b} S_{r-1} \subset q^a F_{\beta_1}^{a_1} \cdots F_{\beta_r}^{a_r} U_q \cap u S_r$$

where $F_{\beta_r}^{b} S_{r-1} \subset S_r$ because F_{β_r} q-commutes with all the other root vectors.

(2) is shown similarly. $\square$

Lemma 4.18 *Let $\nu \in X$ and let $\Sigma = \{\beta_1, \dots, \beta_n\}$ be a basis of Q. Then there exists $\mathbf{b} = (b_1, \dots, b_n) \in (\mathbb{C}^*)^n$ such that*

$$\nu = b_1^{\beta_1} b_2^{\beta_2} \cdots b_n^{\beta_n}$$

and there are only finitely many different $\mathbf{b} \in (\mathbb{C}^*)^n$ satisfying this.

Proof. If $\gamma_1, \gamma_2 \in X$ satisfy $\gamma_1(K_{\beta_i}) = \gamma_2(K_{\beta_i})$ for $i = 1, \dots, n$ then $\gamma_1 = \gamma_2$ because $\{\beta_1, \dots, \beta_n\}$ is a basis of Q. Since for $a_1, \dots, a_n \in \mathbb{C}^*$, $a_1^{\beta_1} a_2^{\beta_2} \cdots a_n^{\beta_n}(K_{\beta_i}) = a_1^{(\beta_i|\beta_1)} a_2^{(\beta_i|\beta_2)} \cdots a_n^{(\beta_i|\beta_n)}$ we have to solve the system in n unknown variables $x_1, \dots, x_n$:

$$\begin{aligned}
x_1^{(\beta_1|\beta_1)} x_2^{(\beta_1|\beta_2)} \cdots x_n^{(\beta_1|\beta_n)} &= \nu(K_{\beta_1}) \\
x_1^{(\beta_2|\beta_1)} x_2^{(\beta_2|\beta_2)} \cdots x_n^{(\beta_2|\beta_n)} &= \nu(K_{\beta_2}) \\
&\vdots \\
x_1^{(\beta_n|\beta_1)} x_2^{(\beta_n|\beta_2)} \cdots x_n^{(\beta_n|\beta_n)} &= \nu(K_{\beta_n}).
\end{aligned}$$

Let $c_j \in \mathbb{C}$, $j = 1, \dots, n$ be such that $\nu(K_{\beta_j}) = e^{c_j}$. There is a choice here since any $c_j + 2k\pi i$, $k \in \mathbb{Z}$ could be chosen instead. Consider the linear system in n unknowns $X_1, \dots, X_n$

$$\begin{aligned}
(\beta_1|\beta_1)\, X_1 + (\beta_1|\beta_2)\, X_2 \cdots (\beta_1|\beta_n)\, X_n &= c_1 \\
(\beta_2|\beta_1)\, X_1 + (\beta_2|\beta_2)\, X_2 \cdots (\beta_2|\beta_n)\, X_n &= c_2 \\
&\vdots \\
(\beta_n|\beta_1)\, X_1 + (\beta_n|\beta_2)\, X_2 \cdots (\beta_n|\beta_n)\, X_n &= c_n.
\end{aligned}$$

This system has a unique solution $a_1, \dots, a_n \in \mathbb{C}$ since the matrix $((\beta_i|\beta_j))_{i,j}$ is invertible. So $x_i = e^{a_i}$ is a solution to the above system. Any other solution to the original system corresponds to making a different choice when taking the logarithm of $\nu(K_{\beta_i})$. So another solution would be of the form $x_i = e^{a_i + a_i'}$

where a'_i, $i = 1, \dots, n$ is a solution to a system of the form:

$$\begin{aligned}
(\beta_1|\beta_1)\, X_1 + (\beta_1|\beta_2)\, X_2 \cdots (\beta_1|\beta_n)\, X_n =& 2k_1\pi i\\
(\beta_2|\beta_1)\, X_1 + (\beta_2|\beta_2)\, X_2 \cdots (\beta_2|\beta_n)\, X_n =& 2k_2\pi i\\
&\vdots\\
(\beta_n|\beta_1)\, X_1 + (\beta_n|\beta_2)\, X_2 \cdots (\beta_n|\beta_n)\, X_n =& 2k_n\pi i.
\end{aligned}$$

for some $k_1, \dots, k_n \in \mathbb{Z}$. Since $A = ((\beta_i|\beta_j))_{i,j}$ is a matrix with only integer coefficients we have $A^{-1} = \frac{1}{\det A}\widetilde{A}$ for some $\widetilde{A}$ with only integer coefficients. So the solution to the system above is integer linear combinations in $\frac{2k_i\pi i}{\det A}$, $i = 1, \dots, n$ hence $\{(e^{a'_1}, \dots, e^{a'_n})|(a'_1, \dots, a'_n)$ is a solution to the above system$\}$ has fewer than $n \det A$ elements so it is a finite set. $\square$

In the next definition we would like to compose the φ's for different β. In particular let $\Sigma = \{\beta_1, \dots, \beta_n\}$ be a set of commuting roots and $F_{\beta_1}, \dots, F_{\beta_n}$ corresponding root vectors. Let $F_\Sigma := \{q^a F_{\beta_1}^{a_1} \cdots F_{\beta_n}^{a_n} | a_i \in \mathbb{N}, a \in \mathbb{Z}\}$ and let $U_{q(F_\Sigma)}$ be the Ore localization in F_Σ. For $i < j$ we have

$$F_{\beta_i}^{-k} F_{\beta_j} F_{\beta_i}^{k} = q^{-k(\beta_i|\beta_j)} F_{\beta_j}$$

or equivalently $\varphi_{F_{\beta_i}, q^k}(F_{\beta_j}) = \left(q^k\right)^{-(\beta_i|\beta_j)} F_{\beta_j}$. This implies $\varphi_{F_{\beta_i}, b}(F_{\beta_j}) = b^{-(\beta_i|\beta_j)} F_{\beta_j}$ for $b \in \mathbb{C}^*$ because $b \mapsto \varphi_{F_{\beta_i}, b}(F_{\beta_j})$ is Laurent polynomial. Similarly $\varphi_{F_{\beta_j}, b}(F_{\beta_i}) = b^{(\beta_i|\beta_j)} F_{\beta_i}$. This shows that we can define $\varphi_{F_\beta, b}(F_{\beta'}^{-1}) = \varphi_{F_\beta, b}(F_{\beta'})^{-1}$ for $\beta, \beta' \in \Sigma$ extending $\varphi_{F_\beta, b}$ to a homomorphism $U_{q(F_\Sigma)} \to U_{q(F_\Sigma)}$. Also note that the φ's commute because

$$\begin{aligned}
F_{\beta_i}^{-k_1} F_{\beta_j}^{-k_2} u F_{\beta_j}^{k_2} F_{\beta_i}^{k_1} =& q^{k_1 k_2 (\beta_i|\beta_j)} F_{\beta_j}^{-k_2} F_{\beta_i}^{-k_1} u q^{-k_1 k_2 (\beta_i|\beta_j)} F_{\beta_i}^{k_1} F_{\beta_j}^{k_2}\\
=& F_{\beta_j}^{-k_2} F_{\beta_i}^{-k_1} u F_{\beta_i}^{k_1} F_{\beta_j}^{k_2}
\end{aligned}$$

Definition 4.19 *Let $\Sigma = \{\beta_1, \dots, \beta_r\}$ be a set of commuting roots and let $F_{\beta_1}, \dots, F_{\beta_r}$ be corresponding root vectors such that $[F_{\beta_j}, F_{\beta_i}]_q = 0$ for $i < j$. Let $U_{q(F_\Sigma)}$ denote the Ore localization of U_q in the Ore set $F_\Sigma := \{q^a F_{\beta_1}^{a_1} \cdots F_{\beta_n}^{a_r} | a_i \in \mathbb{N}, a \in \mathbb{Z}\}$. Said in words we invert F_β for all $\beta \in \Sigma$.*

Let M be a U_q-module. We define M_{F_Σ} to be the $U_{q(F_\Sigma)}$-module $U_{q(F_\Sigma)} \otimes_{U_q} M$. Let $\mathbf{b} = (b_1, \dots, b_r) \in (\mathbb{C}^*)^r$. Then for a $U_{q(F_\Sigma)}$-module N we define $\varphi_{F_\Sigma, \mathbf{b}}.N$ to be the twist of the module by $\varphi_{F_{\beta_1}, b_1} \circ \cdots \circ \varphi_{F_{\beta_r}, b_r}$.

For $\mathbf{i} = (i_1, \dots, i_r) \in \mathbb{Z}^r$ define $q^{\mathbf{i}} = (q^{i_1}, \dots, q^{i_r}) \in (\mathbb{C}^*)^r$ and $q^{\mathbb{Z}^r} = \{q^{\mathbf{i}} | \mathbf{i} \in \mathbb{Z}^r\} \subset (\mathbb{C}^*)^r$.

For $\mathbf{b} = (b_1, \dots, b_r) \in (\mathbb{C}^*)^r$ we set $\mathbf{b}^\Sigma := b_1^{\beta_1} \cdots b_r^{\beta_r} \in X$. If Σ is a basis of Q then the map $\mathbf{b} \mapsto \mathbf{b}^\Sigma$ is surjective by Lemma 4.18 but not neccesarily injective.

Corollary 4.20 (to Lemma 4.12) *Let Σ be a set of commuting roots that is a $\mathbb{Z}$ basis of Q, let F_Σ be an Ore subset corresponding to Σ, let M be a $U_{q(F_\Sigma)}$-module and let $\mathbf{i} = (i_1, \dots, i_n) \in \mathbb{Z}^n$. Then*

$$\varphi_{F_\Sigma, q^{\mathbf{i}}}.M \cong M$$

as $U_{q(F_\Sigma)}$*-modules. Furthermore for* $\lambda \in \operatorname{wt} M$ *we have an isomorphism of* $(U_{q(F_\Sigma)})_0$*-modules:*

$$\varphi_{F_\Sigma, q^{\mathbf{i}}}.M_\lambda \cong M_{(q^{-\mathbf{i}})^\Sigma \lambda} = M_{q^{-\mu}\lambda}$$

where $\mu = \sum_{j=1}^n i_j \beta_j$*.*

Proof. The corollary follows from Lemma 4.12 because Σ is a $\mathbb{Z}$ basis of Q. $\square$

Definition 4.21 *Let* L *be an admissible module of degree* d*. The essential support of* L *is defined as*

$$\operatorname{Supp}_{\mathrm{ess}}(L) := \{\lambda \in \operatorname{wt} L \mid \dim L_\lambda = d\}$$

Lemma 4.22 *Let* M *be an admissible module. Let* $\Sigma \subset \Phi^+$ *be a set of commuting roots and* F_Σ *a corresponding Ore subset. Assume* $-\Sigma \subset T_M$*. Then for* $\lambda \in X$*:*

$$\dim(M_{F_\Sigma})_\lambda = \max_{\mu \in \mathbb{Z}\Sigma} \{\dim M_{q^\mu \lambda}\}$$

and if $\dim M_\lambda = \max_{\mu \in \mathbb{Z}\Sigma}\{\dim M_{q^\mu \lambda}\}$ *then* $(M_{F_\Sigma})_\lambda \cong M_\lambda$ *as* $(U_q)_0$*-modules.*

In particular if $\Sigma \subset T_M$ *as well then* $M_{F_\Sigma} \cong M$ *as* U_q*-modules.*

Compare to Lemma 4.4(ii) in [Mat00].

Proof. We have $\Sigma = \{\beta_1, \dots, \beta_r\}$ for some $\beta_1, \dots, \beta_r \in \Phi^+$ and corresponding root vectors $F_{\beta_1}, \dots, F_{\beta_r}$. Let $\lambda \in X$ and set $d = \max_{\mu \in \mathbb{Z}\Sigma}\{\dim M_{q^\mu \lambda}\}$. Let V be a finite dimensional subspace of $(M_{F_\Sigma})_\lambda$. Then there exists a homogenous element $s \in F_\Sigma$ such that $sV \subset M$. Let $\nu \in \mathbb{Z}\Sigma$ be the degree of s. So $sV \subset M_{q^\nu \lambda}$ hence $\dim sV \leq d$. Since s acts injectively on M_{F_Σ} we have $\dim V \leq d$. Now the first claim follows because $F_\beta^{\pm 1}$ acts injectively on M_{F_Σ} for all $\beta \in \Sigma$.

We have an injective U_q-homomorphism from M to M_{F_Σ} sending $m \in M$ to $1 \otimes m \in M_{F_\Sigma}$ that restricts to a $(U_q)_0$-homomorphism from M_λ to $(M_{F_\Sigma})_\lambda$. If $\dim M_\lambda = d$ then this is surjective as well. So it is an isomorphism. The last claim follow because $\pm\Sigma \subset T_M$ implies $\dim M_\lambda = \dim M_{q^\mu \lambda}$ for any $\mu \in \mathbb{Z}\Sigma$; so $M_\lambda \cong (M_{F_\Sigma})_\lambda$ for any $\lambda \in X$. Since M is a weight module this implies that $M \cong M_{F_\Sigma}$ as U_q-modules. $\square$

Lemma 4.23 *Let* L *be a simple infinite dimensional admissible module. Let* $\beta \in (T_L^s)^+$*. Then there exists a* $b \in \mathbb{C}^*$ *such that* $\varphi_{F_\beta, b}.L_{F_\beta}$ *contains a simple admissible* U_q*-submodule* L' *with* $T_{L'} \subset T_L$ *and* $\beta \notin T_L$*.*

Proof. Since $\beta \in T_L^s$ we have $L \cong L_{F_\beta}$ as U_q-modules by Lemma 4.22. So we will consider L as a $U_{q(F_\beta)}$-module via this isomorphism when taking twist etc.

Let E_β and F_β be root vectors corresponding to β. Let $\lambda \in \operatorname{wt} L$. Consider $F_\beta E_\beta$ as a linear operator on L_λ. Since $\mathbb{C}$ is algebraically closed $F_\beta E_\beta$ must have an eigenvalue c_β and an eigenvector $v \in L_\lambda$. By (the proof of) Lemma 4.9

$$F_\beta E_\beta \varphi_{F_\beta, b}.v = \varphi_{F_\beta, b}.(c_\beta - (q_\beta - q_\beta^{-1})^{-2}(b_\beta - b_\beta^{-1})(q_\beta b_\beta^{-1} \lambda(K_\beta) - q_\beta^{-1} b_\beta \lambda(K_\beta)^{-1})v.$$

The Laurent polynomial, in b, $c_\beta - (q - q^{-1})^{-2}(b_\beta - b_\beta^{-1})(b_\beta \lambda(K_\beta) - b_\beta^{-1} \lambda(K_\beta^{-1}))$ has a zero point $c \in \mathbb{C}^*$.

Thus $\varphi_{F_\beta,c}.L$ contains an element v' such that $F_\beta E_\beta v' = 0$ and since F_β acts injectively on $\varphi_{F_\beta,c}.L$, we have $E_\beta v' = 0$. Set $V = \{m \in \varphi_{F_\beta,c}.L | E_\beta^N m = 0, N >> 0\} = (\varphi_{F_\beta,c}.L)^{[\beta]}$. By Proposition 2.1 this is a U_q-submodule of the U_q-module $\varphi_{F_\beta,c}.L$. It is nonzero since $v' \in V$. By Lemma 4.6 V has a simple U_q-submodule L'.

We want to show that $T_{L'} \subset T_L$. Assume $\gamma \in T_{L'}$. Then $q^{\mathbb{N}\gamma} \operatorname{wt} L' \subset \operatorname{wt} L'$. But since $\operatorname{wt} L' \subset c^{-\beta} \operatorname{wt} L$ we get for some $\nu \in \operatorname{wt} L$, $q^{\mathbb{N}\gamma} c^{-\beta} \nu \subset c^{-\beta} \operatorname{wt} L$ or equivalently $q^{\mathbb{N}\gamma}\nu \subset \operatorname{wt} L$. But this shows that $\gamma \notin F_L$ and since L is a simple U_q-module this implies that $\gamma \in T_L$. By construction we have $\beta \notin T_{L'}$. $\square$

5 Coherent families

For a U_q-module $M \in \mathcal{F}$ define $\operatorname{Tr}^M : X \times (U_q)_0 \to \mathbb{C}$ by $\operatorname{Tr}^M(\lambda, u) = \operatorname{Tr} u|_{M_\lambda}$.

Lemma 5.1 *Let $M, N \in \mathcal{F}$ be semisimple U_q-modules. If $\operatorname{Tr}^M = \operatorname{Tr}^N$ then $M \cong N$.*

Proof. Theorem 7.19 in [Lam01] states that this is true for modules over a *finite dimensional* algebra. So we will reduce to the case of modules over a finite dimensional algebra. Let L be a composition factor of M and λ a weight of L. Then the multiplicity of the U_q-composition factor L in M is the multiplicity of the $(U_q)_0$-composition factor L_λ in M_λ by Theorem 2.7. M_λ is a finite dimensional $(U_q)_0$-module. Let I be the kernel of the homomorphism $(U_q)_0 \to \operatorname{End}_{\mathbb{C}}(M_\lambda)$ given by the action of $(U_q)_0$. Then $(U_q)_0/I$ is a finite dimensional $\mathbb{C}$ algebra and M_λ is a module over $(U_q)_0/I$. Furthermore since $\operatorname{Tr}^M(\lambda, u) = 0$ for all $u \in I$ the trace of an element $u \in (U_q)_0$ is the same as the trace of $u + I \in (U_q)_0/I$ on M_λ as a $(U_q)_0/I$-module. So if $\operatorname{Tr}^M = \operatorname{Tr}^N$ the multiplicity of L_λ in M_λ and N_λ are the same and hence the multiplicity of L in M is the same as in N. $\square$

We will use the Zariski topology on $(\mathbb{C}^*)^n$: V is a closed set if it is the zero-points of a Laurent polynomial $p \in \mathbb{C}[X_1^{\pm 1}, \ldots, X_n^{\pm 1}]$.

Proposition 5.2 *Let L be an infinite dimensional admissible simple module of degree d. Let Σ be a set of commuting roots that is a basis of Q and $w \in W$ such that $-\Sigma \subset w(T_L)$. Let F_Σ be a corresponding Ore subset. Let $\lambda \in \operatorname{Supp}_{\mathrm{ess}}(L)$. The set*

$$\{\mathbf{b} \in (\mathbb{C}^*)^n | \ {}^{\overline{w}}\left(\varphi_{F_\Sigma,\mathbf{b}}.\left(({}^{w}L)_{F_\Sigma}\right)_{w(\lambda)}\right) \text{ is a simple } (U_q)_0\text{-module}\}$$

is a Zariski open set of $(\mathbb{C}^*)^n$.

Proof. The $(U_q)_0$-module $V := {}^{\overline{w}}\left(\varphi_{F_\Sigma,\mathbf{b}}.\left(({}^{w}L)_{F_\Sigma}\right)_{w(\lambda)}\right)$ is simple if and only if the bilinear map $B_{\mathbf{b}}(u,v) \in (U_q)_0 \times (U_q)_0 \mapsto \operatorname{Tr}\left(uv|_{{}^{\overline{w}}\left(\varphi_{F_\Sigma,\mathbf{b}}.\left(({}^{w}L)_{F_\Sigma}\right)_{w(\lambda)}\right)}\right)$ has maximal rank d^2: The map factors through $\operatorname{End}_{\mathbb{C}}(V) \times \operatorname{End}_{\mathbb{C}}(V)$ given by the representation $(U_q)_0 \to \operatorname{End}_{\mathbb{C}}(V)$ on V. $B_{\mathbf{b}}$ has maximal rank d^2 if and only if the representation is surjective onto $\operatorname{End}_{\mathbb{C}}(V)$ which is equivalent to V being simple.

For any finite dimensional subspace $E \subset (U_q)_0$, the set Ω_E of all $\mathbf{b}$ such that $B_{\mathbf{b}}|_E$ has rank d^2 is either empty or the non-zero points of the Laurent polynomial $\det M$ for some $d^2 \times d^2$ minor M of the matrix $(B_{\mathbf{b}}(e_i, e_j))_{i,j}$ where $\{e_i\}$ is a basis of E. Therefore $\Omega = \cup_E \Omega_E$ is open. $\square$

For a module M that is a direct sum of modules of finite length we define M^{ss} to be the unique (up to isomorphism) semisimple module with the same composition factors as M.

Lemma 5.3 *Let L be an infinite dimensional simple admissible U_q-module of degree d, $w \in W$ and $\Sigma = \{\beta_1, \dots, \beta_n\} \subset \Phi^+$ a set of commuting roots that is a basis of Q such that $-\Sigma \subset w(T_L)$. Let F_Σ be a corresponding Ore subset to Σ. Let $\mathbf{c} \in (\mathbb{C}^*)^n$ and let L' be another infinite dimensional U_q-module such that L' is contained in $^{\overline{w}}\left(\varphi_{F_\Sigma,\mathbf{c}}.(^wL)_{F_\Sigma}\right)^{ss}$ (i.e. L' is a composition factor of $^{\overline{w}}\left(\varphi_{F_\Sigma,\mathbf{c}}.(^wL)_{F_\Sigma}\right)$). Assume that $\Sigma' = \{\beta'_1, \dots, \beta'_n\} \subset \Phi^+$ is another set of commuting roots that is a basis of Q and $w' \in W$ is such that $-\Sigma' \subset w'(T_{L'})$. Let $F_{\Sigma'}$ be a corresponding Ore subset.*

Define $a_{i,j} \in \mathbb{Z}$ by $w(w')^{-1}(\beta'_i) = \sum_{j=1}^n a_{i,j}\beta_j$ and define $f : (\mathbb{C}^*)^n \to (\mathbb{C}^*)^n$ by

$$f(b_1, \dots, b_n) = \left(\prod_{i=1}^n b_i^{a_{i,1}}, \dots, \prod_{i=1}^n b_i^{a_{i,n}}\right).$$

Then L' is admissible of degree d and

$$^{\overline{w'}}\left(\varphi_{F_{\Sigma'},\mathbf{b}}.(^{w'}L')_{F_{\Sigma'}}\right)^{ss} \cong {^{\overline{w}}}\left(\varphi_{F_\Sigma,f(\mathbf{b})\mathbf{c}}.(^wL)_{F_\Sigma}\right)^{ss}$$

Proof. We will show that $\operatorname{Tr}^{\overline{w'}\left(\varphi_{F_{\Sigma'},\mathbf{b}}.(^{w'}L')_{F_{\Sigma'}}\right)^{ss}} = \operatorname{Tr}^{\overline{w}\left(\varphi_{F_\Sigma,f(\mathbf{b})\mathbf{c}}.(^wL)_{F_\Sigma}\right)^{ss}}$.

Let $\lambda \in \operatorname{Supp}_{\text{ess}}(L)$. Then $w(\lambda) \in \operatorname{Supp}_{\text{ess}}(^wL)$. As a $(U_q)_0$-module we have $\left(^{\overline{w}}\left(\varphi_{F_\Sigma,\mathbf{c}}.(^wL)_{F_\Sigma}\right)\right)^{ss} \cong \bigoplus_{\mathbf{i}\in\mathbb{Z}^n} {^{\overline{w}}}\left(\varphi_{F_\Sigma,q^{\mathbf{i}}\mathbf{c}}.\left((^wL)_{F_\Sigma}\right)_{w(\lambda)}\right)^{ss}$ (Corollary 4.20). Let $\lambda' \in \operatorname{Supp}_{\text{ess}}(L')$. Then $L'_{\lambda'}$ is a $(U_q)_0$-submodule of $^{\overline{w}}\left(\varphi_{F_\Sigma,q^{\mathbf{j}}\mathbf{c}}.\left((^wL)_{F_\Sigma}\right)_{w(\lambda)}\right)^{ss}$ for some $\mathbf{j} \in \mathbb{Z}^n$. We can assume $\mathbf{j} = 0$ by replacing $\mathbf{c}$ with $q^{\mathbf{j}}\mathbf{c}$ (note that we have then $(\mathbf{c}^{-1})^\Sigma = w\left(\lambda'\lambda^{-1}\right)$). So $L'_{\lambda'}$ is a $(U_q)_0$-submodule of $^{\overline{w}}\left(\varphi_{F_\Sigma,\mathbf{c}}.\left((^wL)_{F_\Sigma}\right)_{w(\lambda)}\right)^{ss}$. For any other $\mu \in \operatorname{Supp}_{\text{ess}}(L')$ there is a unique $\mathbf{j}'_\mu \in \mathbb{Z}^n$ such that $\mu = (w')^{-1}\left(\left(q^{-\mathbf{j}'_\mu}\right)^{\Sigma'}\right)\lambda'$ and a unique $\mathbf{j}_\mu \in \mathbb{Z}^n$ such that $w^{-1}\left(\left(q^{-\mathbf{j}_\mu}\mathbf{c}^{-1}\right)^{\Sigma}\right)\lambda = \mu$. For such $\mathbf{j}_\mu$, L'_μ is a submodule of $^{\overline{w}}\left(\varphi_{F_\Sigma,q^{\mathbf{j}_\mu}\mathbf{c}}.\left((^wL)_{F_\Sigma}\right)_{w(\lambda)}\right)^{ss}$.

f is bijective, $f(q^{\mathbb{Z}^n}) = q^{\mathbb{Z}^n}$, $f(\mathbf{b})^\Sigma = w(w')^{-1}\left(\mathbf{b}^{\Sigma'}\right)$ for all $\mathbf{b} \in (\mathbb{C}^*)^n$ and for any $\mu \in \operatorname{Supp}_{\text{ess}}(L')$, $f(q^{\mathbf{j}'_\mu}) = q^{\mathbf{j}_\mu}$. For a Laurent polynomial p, $p \circ f$ is Laurent polynomial as well. Since $q^{\mathbb{N}^n}$ is Zariski dense in $(\mathbb{C}^*)^n$ (Lemma 4.8) and f is a Laurent polynomial the set $D = \{q^{\mathbf{j}_\mu}\mathbf{c} \in (\mathbb{C}^*)^n | \mu \in \operatorname{Supp}_{\text{ess}}(L')\}$ is Zariski dense. By Proposition 5.2 the $(U_q)_0$-module $^{\overline{w}}\left(\varphi_{F_\Sigma,\mathbf{b}}.\left((^wL)_{F_\Sigma}\right)_{w(\lambda)}\right)$ is simple for all $\mathbf{b} \in \Omega$ for some Zariski open set Ω of $(\mathbb{C}^*)^n$. Since D is dense and Ω is open $D \cap \Omega$ is nonempty. So there exists a $\mu_0 \in \operatorname{Supp}_{\text{ess}}(L')$ such that $^{\overline{w}}\left(\varphi_{F_\Sigma,q^{\mathbf{j}_{\mu_0}}\mathbf{c}}.\left((^wL)_{F_\Sigma}\right)_{w(\lambda)}\right)$ is simple and contains the nonzero simple $(U_q)_0$-module L'_{μ_0} as a submodule. Thus $L'_{\mu_0} \cong {^{\overline{w}}}\left(\varphi_{F_\Sigma,q^{\mathbf{j}_{\mu_0}}\mathbf{c}}.\left((^wL)_{F_\Sigma}\right)_{w(\lambda)}\right)$. We get now from Lemma 4.22 that L' is admissible of degree d and that for every $\mu \in \operatorname{Supp}_{\mathrm{ess}}(L')$,

$$\begin{aligned} L'_\mu &\cong {}^{\overline{w}}\left(\varphi_{F_\Sigma, q^{\mathbf{j}_\mu}\mathbf{c}} . \left(\left({}^{w}L\right)_{F_\Sigma}\right)_{w(\lambda)}\right) \\ &\cong {}^{\overline{w}}\left(\varphi_{F_\Sigma, f(q^{\mathbf{j}'_\mu})\mathbf{c}} . \left(\left({}^{w}L\right)_{F_\Sigma}\right)_{w(\lambda)}\right). \end{aligned}$$

By Lemma 4.22, Corollary 4.20 and the definition of $\mathbf{j}'_\mu$ we have for any $\mu \in \operatorname{Supp}_{\mathrm{ess}}(L')$

$$^{\overline{w'}}\left(\varphi_{F_{\Sigma'}, q^{\mathbf{j}'_\mu}} . \left(\left({}^{w'}L'\right)_{F_{\Sigma'}}\right)_{w'(\lambda')}\right) \cong L'_\mu.$$

Let $u \in (U_q)_0$. We see that for $\mathbf{b} = q^{\mathbf{j}'_\mu}$

$$\operatorname{Tr} u|_{{}^{\overline{w}}\left(\varphi_{F_\Sigma, f(\mathbf{b})\mathbf{c}} . \left(\left({}^{w}L\right)_{F_\Sigma}\right)_{w(\lambda)}\right)} = \operatorname{Tr} u|_{L'_\mu} = \operatorname{Tr} u|_{{}^{\overline{w'}}\left(\varphi_{F_{\Sigma'}, \mathbf{b}} . \left(\left({}^{w'}L'\right)_{F_{\Sigma'}}\right)_{w'(\lambda')}\right)}.$$

Since $\mathbf{b} \mapsto \operatorname{Tr} u|_{{}^{\overline{w}}\left(\varphi_{F_\Sigma, f(\mathbf{b})\mathbf{c}} . \left(\left({}^{w}L\right)_{F_\Sigma}\right)_{w(\lambda)}\right)^{ss}}$ and $\mathbf{b} \mapsto \operatorname{Tr} u|_{{}^{\overline{w'}}\left(\varphi_{F_{\Sigma'}, \mathbf{b}} . \left(\left({}^{w'}L'\right)_{F_{\Sigma'}}\right)_{w'(\lambda')}\right)^{ss}}$ are both Laurent polynomials and equal on the Zariski dense subset $\{q^{\mathbf{j}'_\mu} | \mu \in \operatorname{Supp}_{\mathrm{ess}}(L')\}$ they are equal for all $\mathbf{b} \in (\mathbb{C}^*)^n$. Thus by Lemma 5.1

$${}^{\overline{w}}\left(\varphi_{F_\Sigma, f(\mathbf{b})\mathbf{c}} . \left(\left({}^{w}L\right)_{F_\Sigma}\right)_{w(\lambda)}\right)^{ss} \cong {}^{\overline{w'}}\left(\varphi_{F_{\Sigma'}, \mathbf{b}} . \left(\left({}^{w'}L'\right)_{F_{\Sigma'}}\right)_{w'(\lambda')}\right)^{ss}$$

as $(U_q)_0$-modules. Since (by Corollary 4.20)

$${}^{\overline{w'}}\left(\varphi_{F_{\Sigma'}, \mathbf{b}} . \left({}^{w'}L'\right)_{F_{\Sigma'}}\right)^{ss} \cong \bigoplus_{\mathbf{i} \in \mathbb{Z}^n} {}^{\overline{w'}}\left(\varphi_{F_{\Sigma'}, q^{\mathbf{i}}\mathbf{b}} . \left(\left({}^{w'}L'\right)_{F_{\Sigma'}}\right)_{w'(\lambda')}\right)^{ss}$$

and

$${}^{\overline{w}}\left(\varphi_{F_\Sigma, f(\mathbf{b})\mathbf{c}} . \left({}^{w}L\right)_{F_\Sigma}\right)^{ss} \cong \bigoplus_{\mathbf{i} \in \mathbb{Z}^n} {}^{\overline{w}}\left(\varphi_{F_\Sigma, q^{\mathbf{i}}f(\mathbf{b})\mathbf{c}} . \left(\left({}^{w}L\right)_{F_\Sigma}\right)_{w(\lambda)}\right)^{ss}$$

we get

$$\begin{aligned} {}^{\overline{w'}}\left(\varphi_{F_{\Sigma'}, \mathbf{b}} . \left({}^{w'}L'\right)_{F_{\Sigma'}}\right)^{ss} &\cong \bigoplus_{\mathbf{i} \in \mathbb{Z}^n} {}^{\overline{w'}}\left(\varphi_{F_{\Sigma'}, q^{\mathbf{i}}\mathbf{b}} . \left(\left({}^{w'}L'\right)_{F_{\Sigma'}}\right)_{w'(\lambda')}\right)^{ss} \\ &\cong \bigoplus_{\mathbf{i} \in \mathbb{Z}^n} {}^{\overline{w}}\left(\varphi_{F_\Sigma, f(q^{\mathbf{i}}\mathbf{b})\mathbf{c}} . \left(\left({}^{w}L\right)_{F_\Sigma}\right)_{w(\lambda)}\right)^{ss} \\ &\cong \bigoplus_{\mathbf{i} \in \mathbb{Z}^n} {}^{\overline{w}}\left(\varphi_{F_\Sigma, q^{\mathbf{i}}f(\mathbf{b})\mathbf{c}} . \left(\left({}^{w}L\right)_{F_\Sigma}\right)_{w(\lambda)}\right)^{ss} \\ &\cong {}^{\overline{w}}\left(\varphi_{F_\Sigma, f(\mathbf{b})\mathbf{c}} . \left({}^{w}L\right)_{F_\Sigma}\right)^{ss} \end{aligned}$$

as $(U_q)_0$-modules. By Theorem 2.7 this implies they are isomorphic as U_q-modules as well. □

Corollary 4.20 tells us that twisting with an element of the form $q^{\mathbf{i}}$ gives us a module isomorphic to the original module. Thus it makes sense to write $\varphi_{F_\Sigma, t} . M$

for a $t \in (\mathbb{C}^*)^n/q^{\mathbb{Z}^n}$ and a $U_{q(F_\Sigma)}$-module M. Just choose a representative for t. Any representative gives the same $U_{q(F_\Sigma)}$-module up to isomorphism.

Let L be an admissible simple module. Assume for a $w \in W$ that $\Sigma \subset -w(T_L)$ is a set of commuting roots that is a basis of Q (it is always possible to find such w and Σ by Lemma 4.15 and Lemma 4.16) and let F_Σ be a corresponding Ore subset. Let $\nu \in X$. The U_q-module

$$^{\overline{w}}\left(\bigoplus_{\mathbf{b}\in(\mathbb{C}^*)^n:\,\mathbf{b}^\Sigma=\nu} \varphi_{F_\Sigma,\mathbf{b}}.\left({}^wL\right)_{F_\Sigma}\right)$$

has finite length by Lemma 4.18, Lemma 4.22 and Lemma 4.6.

We define

$$\mathcal{EXT}(L) = \left(\bigoplus_{t\in(\mathbb{C}^*)^n/q^{\mathbb{Z}^n}} {}^{\overline{w}}\left(\varphi_{F_\Sigma,t}.\left({}^wL\right)_{F_\Sigma}\right)\right)^{ss}.$$

The definition is independent (up to isomorphism) of the chosen w, Σ and F_Σ as suggested by the notation:

Lemma 5.4 *Let L be a simple admissible module. Let $w, w' \in W$ and assume $\Sigma \subset -w(T_L), \Sigma' \subset -w'(T_{L'})$ are sets of commuting roots that are both a basis of Q. Let $F_\Sigma, F'_{\Sigma'}$ be corresponding Ore subsets. Then*

$$\left(\bigoplus_{t\in(\mathbb{C}^*)^n/q^{\mathbb{Z}^n}} {}^{\overline{w}}\left(\varphi_{F_\Sigma,t}.\left({}^wL\right)_{F_\Sigma}\right)\right)^{ss} \cong \left(\bigoplus_{t\in(\mathbb{C}^*)^n/q^{\mathbb{Z}^n}} {}^{\overline{w'}}\left(\varphi_{F'_{\Sigma'},t}.\left({}^{w'}L\right)_{F'_{\Sigma'}}\right)\right)^{ss}$$

as U_q-modules.

Proof. Obviously L is a submodule of $\left({}^{\overline{w}}\left(\varphi_{F_\Sigma,\mathbf{1}}.\left({}^wL\right)_{F_\Sigma}\right)\right)^{ss}$ where $\mathbf{1} = (1,\dots,1)$. By Lemma 5.3 this implies that for $\mathbf{b} \in (\mathbb{C}^*)^n$

$$\left({}^{\overline{w'}}\left(\varphi_{F_{\Sigma'},\mathbf{b}}.\left({}^{w'}L\right)_{F_{\Sigma'}}\right)\right)^{ss} \cong \left({}^{\overline{w}}\left(\varphi_{F_\Sigma,f(\mathbf{b})}.\left({}^wL\right)_{F_\Sigma}\right)\right)^{ss}$$

for some f with the property that $f(q^{\mathbb{Z}^n}) = q^{\mathbb{Z}^n}$. So it makes sense to write $f(t)$ for $t \in (\mathbb{C}^*)^n/q^{\mathbb{Z}^n}$. Thus

$$\begin{aligned}\left(\bigoplus_{t\in(\mathbb{C}^*)^n/q^{\mathbb{Z}^n}} {}^{\overline{w'}}\left(\varphi_{F'_{\Sigma'},t}.\left({}^{w'}L\right)_{F'_{\Sigma'}}\right)\right)^{ss} &\cong \left(\bigoplus_{t\in(\mathbb{C}^*)^n/q^{\mathbb{Z}^n}} {}^{\overline{w}}\left(\varphi_{F_\Sigma,f(t)}.\left({}^wL\right)_{F_\Sigma}\right)\right)^{ss}\\ &\cong \left(\bigoplus_{t\in(\mathbb{C}^*)^n/q^{\mathbb{Z}^n}} {}^{\overline{w}}\left(\varphi_{F_\Sigma,t}.\left({}^wL\right)_{F_\Sigma}\right)\right)^{ss}\end{aligned}$$

since f is bijective. □

Proposition 5.5 *Let L be a simple infinite dimensional admissible module. For $x \in W$:*

$$\mathcal{EXT}({}^xL) \cong {}^x\left(\mathcal{EXT}(L)\right)$$

and

$$\mathcal{EXT}({}^{\overline{x}}L) \cong {}^{\overline{x}}\left(\mathcal{EXT}(L)\right).$$

Proof. Let $w \in W$ be such that $w(F_L \backslash F_L^s) \subset \Phi^+$ (exists by Lemma 4.15). Let Σ be a set of commuting roots that is a basis of Q such that $-\Sigma \subset w(T_L)$ (exists by Lemma 4.16) and let F_Σ be a corresponding Ore subset. First we will define $\mathcal{EXT}'(L) = \left(\bigoplus_{t \in (\mathbb{C}^*)^n / q^{\mathbb{Z}^n}} {}^{w^{-1}}\left(\varphi_{F_\Sigma, t} . ({}^{\overline{w^{-1}}} L)_{F_\Sigma}\right)\right)^{ss}$ and show that $\mathcal{EXT}'(L) \cong \mathcal{EXT}(L)$ as U_q-modules: Going through the proof of Lemma 5.3 and Lemma 5.4 and and replacing $T_{w^{-1}}$ and $T_{w^{-1}}^{-1}$ with T_w^{-1} and T_w respectively we get that $\mathrm{Tr}^{\mathcal{EXT}'(L)} = \mathrm{Tr}^{\mathcal{EXT}(L)}$ so they are isomorphic by Lemma 5.1.

We will show for any $\alpha \in \Pi$ that

$$\mathcal{EXT}({}^{s_\alpha} L) \cong {}^{s_\alpha}\left(\mathcal{EXT}(L)\right)$$

which implies the claim by induction over the length $l(x)$ of x (where $l(x)$ is the smallest number of simple reflections need to write x, i.e. there is a reduced expression $x = s_{i_1} \cdots s_{i_{l(x)}}$).

So let $\alpha \in \Pi$ and let w and Σ be defined as above. Let $w' = w s_\alpha$. Note that $w'(F_{{}^{s_\alpha} L} \backslash F^s_{{}^{s_\alpha} L}) \subset \Phi^+$ and $-\Sigma \subset T_{{}^{s_\alpha} L}$. We split into two cases: If $l(w') < l(w)$ then

$$\begin{aligned}
{}^{s_\alpha}\left(\mathcal{EXT}(L)\right) =& {}^{s_\alpha}\left(\left(\bigoplus_{t \in (\mathbb{C}^*)^n / q^{\mathbb{Z}^n}} {}^{\overline{w' s_\alpha}}\left(\varphi_{F_\Sigma, t} . ({}^{w' s_\alpha} L)_{F_\Sigma}\right)\right)^{ss}\right) \\
\cong& {}^{s_\alpha}\left(\left(\bigoplus_{t \in (\mathbb{C}^*)^n / q^{\mathbb{Z}^n}} {}^{\overline{s_\alpha}}\left({}^{\overline{w'}}\left(\varphi_{F_\Sigma, t} . ({}^{w'}({}^{s_\alpha} L))_{F_\Sigma}\right)\right)\right)^{ss}\right) \\
\cong& \left(\bigoplus_{t \in (\mathbb{C}^*)^n / q^{\mathbb{Z}^n}} {}^{\overline{w'}}\left(\varphi_{F_\Sigma, t} . ({}^{w'}({}^{s_\alpha} L))_{F_\Sigma}\right)\right)^{ss} \\
=& \mathcal{EXT}({}^{s_\alpha} L).
\end{aligned}$$

If $l(w') > l(w)$ we get

$$\begin{aligned}
{}^{s_\alpha}\left(\mathcal{EXT}(L)\right) \cong& {}^{s_\alpha}\left(\mathcal{EXT}'(L)\right) \\
=& {}^{s_\alpha}\left(\left(\bigoplus_{t \in (\mathbb{C}^*)^n / q^{\mathbb{Z}^n}} {}^{w^{-1}}\left(\varphi_{F_\Sigma, t} . ({}^{\overline{w^{-1}}} L)_{F_\Sigma}\right)\right)^{ss}\right) \\
\cong& \left(\bigoplus_{t \in (\mathbb{C}^*)^n / q^{\mathbb{Z}^n}} {}^{(w')^{-1}}\left(\varphi_{F_\Sigma, t} . ({}^{\overline{(w')^{-1}}}({}^{s_\alpha} L))_{F_\Sigma}\right)\right)^{ss} \\
=& \mathcal{EXT}({}^{s_\alpha} L).
\end{aligned}$$

The second claim is shown similarly. $\square$

Proposition 5.6 *Let L be an infinite dimensional admissible simple module of degree d. If L' is an infinite dimensional simple submodule of $\mathcal{EXT}(L)$ then L' is admissible of degree d and $\mathcal{EXT}(L) \cong \mathcal{EXT}(L')$.*

Proof. Let $w \in W$ and let Σ be a set of commuting roots that is a basis of Q such that $\Sigma \subset -w(T_L)$ (possible by Lemma 4.15 and Lemma 4.16). Then by definition

$$\mathcal{EXT}(L) = \left(\bigoplus_{t \in (\mathbb{C}^*)^n / q^{\mathbb{Z}^n}} {}^{\overline{w}}\left(\varphi_{F_\Sigma, t} \cdot \left({}^{w}L\right)_{F_\Sigma} \right) \right)^{ss}.$$

L' being a submodule of $\mathcal{EXT}(L)$ implies that L' must be a submodule of

$$\left({}^{\overline{w}}\left(\varphi_{F_\Sigma, \mathbf{c}} \cdot \left({}^{w}L\right)_{F_\Sigma} \right) \right)^{ss}$$

for some $\mathbf{c} \in (\mathbb{C}^*)^n$. Let $w' \in W$ and let Σ' be a set of commuting roots that is a basis of Q such that $\Sigma' \subset -w'(T_{L'})$. By Lemma 5.3 L' is admissible of degree d and there exists a bijective map $f : (\mathbb{C}^*)^n \to (\mathbb{C}^*)^n$ such that $f(q^{\mathbb{Z}^n}) = q^{\mathbb{Z}^n}$ and

$$\left({}^{\overline{w'}}\left(\varphi_{F_{\Sigma'}, \mathbf{b}} \cdot \left({}^{w'}L'\right)_{F_{\Sigma'}} \right) \right)^{ss} \cong \left({}^{\overline{w}}\left(\varphi_{F_\Sigma, f(\mathbf{b})\mathbf{c}} \cdot \left({}^{w}L\right)_{F_\Sigma} \right) \right)^{ss}.$$

Since $f(q^{\mathbb{Z}^n}) = q^{\mathbb{Z}^n}$ it makes sense to write $f(t)$ for $t \in (\mathbb{C}^*)^n / q^{\mathbb{Z}^n}$. So writing $t_{\mathbf{c}} = q^{\mathbb{Z}^n}\mathbf{c} \in (\mathbb{C}^*)^n / q^{\mathbb{Z}^n}$ we get

$$\begin{aligned}
\mathcal{EXT}(L') &= \left(\bigoplus_{t \in (\mathbb{C}^*)^n / q^{\mathbb{Z}^n}} {}^{\overline{w'}}\left(\varphi_{F_\Sigma, t} \cdot \left({}^{w'}(L')\right)_{F_{\Sigma'}} \right) \right)^{ss} \\
&\cong \left(\bigoplus_{t \in (\mathbb{C}^*)^n / q^{\mathbb{Z}^n}} {}^{\overline{w}}\left(\varphi_{F_\Sigma, f(t)t_{\mathbf{c}}} \cdot \left({}^{w}L\right)_{F_\Sigma} \right) \right)^{ss} \\
&\cong \left(\bigoplus_{t \in (\mathbb{C}^*)^n / q^{\mathbb{Z}^n}} {}^{\overline{w}}\left(\varphi_{F_\Sigma, t} \cdot \left({}^{w}L\right)_{F_\Sigma} \right) \right)^{ss} \\
&= \mathcal{EXT}(L)
\end{aligned}$$

since the assignment $t \mapsto f(t)t_{\mathbf{c}}$ is bijective. □

Lemma 5.7 *Let $f \in \mathbb{C}[X_1^{\pm 1}, \dots, X_n^{\pm 1}]$ be a nonzero Laurent polynomial. There exists $b_1, \dots, b_n \in \mathbb{C}^*$ such that for all $i_1, \dots, i_n \in \mathbb{Z}$*

$$f(q^{i_1}b_1, \dots, q^{i_n}b_n) \neq 0.$$

Proof. Assume $f = X_1^{-N_1} \cdots X_n^{-N_n} g$ with $g \in \mathbb{C}[X_1, \dots, X_n]$. g has coefficients in some finitely generated (over $\mathbb{Q}$) subfield k of $\mathbb{C}$. Let $b_1, \dots b_n$ be generators of n disjoint extensions of k of degree $> \deg g$. The monomials $b_1^{m_1} \cdots b_n^{m_n}$, $0 \leq m_i \leq \deg g$ are all linearly independent over k. Since $q^i \neq 0$ for $i \in \mathbb{Z}$ the same is true for the monomials $(q^{i_1}b_1)^{m_1} \cdots (q^{i_n}b_n)^{m_n}$. So $g(q^{i_1}b_1, \dots, q^{i_n}b_n) \neq 0$, hence $f(q^{i_1}b_1, \dots, q^{i_n}b_n) \neq 0$. □

Theorem 5.8 *Let L be an infinite dimensional admissible simple modules of degree d. Then $\mathcal{EXT}(L)$ contains at least one simple torsion free module.*

Proof. Let $\lambda \in w(\operatorname{wt} L)$. Then as a $(U_q)_0$-module

$$\mathcal{EXT}(L) = \left({}^{\overline{w}}\left(\bigoplus_{\mathbf{b} \in (\mathbb{C}^*)^n} \varphi_{F_\Sigma,\mathbf{b}}.\left(({}^wL)_{F_\Sigma}\right)_\lambda \right) \right)^{ss}$$

for some $w \in W$ and some Ore subset F_Σ corresponding to a set of commuting roots Σ that is a basis of Q. Let $u \in (U_q)_0$. Then the map $\mathbf{b} \mapsto \det u|_{{}^{\overline{w}}\left(\varphi_{F_\Sigma,\mathbf{b}}.\left(({}^wL)_{F_\Sigma}\right)_\lambda\right)} = \det \varphi_{F_\Sigma,\mathbf{b}}(T_w^{-1}(u))|_{\left(({}^wL)_{F_\Sigma}\right)_\lambda}$ is Laurent polynomial. Let $p(\mathbf{b}) = \prod_{\beta\in\Sigma} \det E_\beta F_\beta|_{{}^{\overline{w}}\left(\varphi_{F_\Sigma,\mathbf{b}}.\left(({}^wL)_{F_\Sigma}\right)_\lambda\right)}$. p is a Laurent polynomial by the above. By Lemma 5.7 there exists a $\mathbf{c} \in (\mathbb{C}^*)^n$ such that $p(\mathbf{b}) \neq 0$ for all $\mathbf{b} \in q^{\mathbb{Z}^n}\mathbf{c}$ which implies that $E_\beta F_\beta$ acts injectively on the module $L' := {}^{\overline{w}}\left(\varphi_{F_\beta,\mathbf{c}}.({}^wL)_{F_\Sigma}\right)$ for all $\beta \in \Sigma$. Since F_β acts injectively on the module by construction this implies that E_β acts injectively as well. So we have $\pm\Sigma \subset T_{L'}$. Any simple submodule V of L' is admissible of degree d by Lemma 5.3 and since F_β and E_β act injectively we get $\dim V_\lambda = d = \dim L'_\lambda$ for any $\lambda \in \operatorname{wt} L'$ thus $V = L'$. So L' is a simple module. Using Proposition 2.5 it is easy to see that L' is torsion free since $\pm\Sigma \subset T_{L'}$ and Σ is a basis of Q. □

Proposition 5.9 *Let L be an infinite dimensional admissible simple module. Let $\beta \in \Phi^+$. If $-\beta \in T_L$ then $\mathcal{EXT}(L)$ contains $\left(\bigoplus_{t\in\mathbb{C}^*/q^{\mathbb{Z}}} \varphi_{F_\beta,t}.L_{F_\beta}\right)^{ss}$ as a U_q-submodule.*

Proof. Let $w \in W$ and $\Sigma = \{\beta_1, \dots, \beta_n\}$ be such that Σ is a set of commuting roots that is a basis of Q and $-\Sigma \subset w(T_L)$ and F_Σ a corresponding Ore subset (always possible by Lemma 4.15 and Lemma 4.16).

We have $w(\beta) = \sum_{i=1}^n a_i\beta_i$ for some $a_i \in \mathbb{Z}$. Set $x = F_{\beta_1}^{a_1} \cdots F_{\beta_n}^{a_n} \in U_{q(F_\Sigma)}$. Let $U_{q(x)}$ be the U_q-subalgebra generated by x in $U_{q(F_\Sigma)}$. x is playing the role of F_β and that is why the notation resembles the notation for Ohre localization. The Ohre localization of U_q in x does not neccesarily make sense though because x is not neccesarily an element of U_q.

Let V be the $U_{q(x)}$-submodule of $({}^wL)_{F_\Sigma}$ generated by $1 \otimes {}^wL$. For any $t \in \mathbb{C}^*/q^{\mathbb{Z}}$

$${}^{\overline{w}}\left(\varphi_{F_\Sigma,(t^{a_1},\dots,t^{a_n})}.V\right) := \left\{\varphi_{F_\Sigma,(t^{a_1},\dots,t^{a_n})}.v \in {}^{\overline{w}}\left(\varphi_{F_\Sigma,(t^{a_1},\dots,t^{a_n})}.({}^wL)_{F_\Sigma}\right) | v \in V\right\}$$

is a $U_{q(x)}$-submodule of ${}^{\overline{w}}\left(\varphi_{F_\Sigma,(t^{a_1},\dots,t^{a_n})}.({}^wL)_{F_\Sigma}\right)$: To show this we show that for $u \in U_{q(x)}$ and $c \in \mathbb{C}^*$, $\varphi_{F_\Sigma,(c^{a_1},\dots,c^{a_n})}(u) \in U_{q(x)}$. We know that $\varphi_{F_\Sigma,(c^{a_1},\dots,c^{a_n})}(u) \in U_{q(F_\Sigma)}[c^{\pm 1}]$ and we also see by construction that for $c = q^i$, $i \in \mathbb{Z}$, we have $\varphi_{F_\Sigma,(c^{a_1},\dots,c^{a_n})}(u) = x^{-i}ux^i \in U_{q(x)}$. Choose a vector space basis of $U_{q(x)}$, $\{u_i\}_{i\in I}$ and extend to a basis $\{u_i, u'_j\}_{i\in I, j\in J}$ of $U_{q(F_\Sigma)}$ where I and J are some index sets. Then for $u \in U_{q(x)}$ we have $\varphi_{F_\Sigma,(c^{a_1},\dots,c^{a_n})}(u) = \sum_{i\in I'} u_ip_i(c) + \sum_{j\in J'} u'_jp'_j(c)$ for some finite $I' \subset I$ and $J' \subset J$ and some $p_i, p'_j \in \mathbb{C}[X^{\pm 1}]$. We see that for $j \in J'$, $p'_j(q^i) = 0$ for all $i \in \mathbb{Z}$ so $p'_j = 0$. Hence $\varphi_{F_\Sigma,(c^{a_1},\dots,c^{a_n})}(u) = \sum_{i\in I'} u_ip_i(c) \in U_{q(x)}$. This shows that ${}^{\overline{w}}\left(\varphi_{F_\Sigma,(t^{a_1},\dots,t^{a_n})}.V\right)$ is a submodule of ${}^{\overline{w}}\left(\varphi_{F_\Sigma,(t^{a_1},\dots,t^{a_n})}.({}^wL)_{F_\Sigma}\right)$. Set

$$\mathcal{V} = \left(\bigoplus_{t\in\mathbb{C}^*/q^{\mathbb{Z}}} {}^{\overline{w}}\left(\varphi_{F_\Sigma,(t^{a_1},\dots,t^{a_n})}.V\right) \right)^{ss}.$$

Clearly $\mathcal{V}$ is a U_q-submodule of $\mathcal{EXT}(L)$. We claim that $\mathcal{V} \cong \left(\bigoplus_{t\in\mathbb{C}^*/q^{\mathbb{Z}}} \varphi_{F_\beta,t}.L_{F_\beta}\right)^{ss}$ as U_q-modules. We will show this using Lemma 5.1.

Note that for $\lambda \in \operatorname{wt} V$ and $i \in \mathbb{Z}$ we have

$$\overline{w}\left(\varphi_{F_\Sigma,((q^i)^{a_1},\dots,(q^i)^{a_n})}.V_\lambda\right) \cong \overline{w}\left(V_{q^{-i\sum_{k=1}^n a_k\beta_k}\lambda}\right)$$

as a $(U_q)_0$-module by Corollary 4.20.

We have $\operatorname{wt}\mathcal{V} = (\mathbb{C}^*)^\beta \operatorname{wt} L = \operatorname{wt}\left(\bigoplus_{t\in\mathbb{C}^*/q^{\mathbb{Z}}} \varphi_{F_\beta,t}.L_{F_\beta}\right)^{ss}$. Let $\lambda \in \operatorname{wt} L$ be such that $\dim L_\lambda = \max_{i\in\mathbb{Z}}\{\dim L_{q^{i\beta}\lambda}\}$ then $V_{w(\lambda)} \cong ({}^wL)_{w(\lambda)} \cong {}^w(L_\lambda)$ as a $(U_q)_0$-module by Lemma 4.22 and we have for $\nu \in (\mathbb{C}^*)^\beta\lambda$:

$$\mathcal{V}_\nu = \left(\bigoplus_{c\in\mathbb{C}^*:c^{w(\beta)}=w(\nu^{-1}\lambda)} \overline{w}\left(\varphi_{F_\Sigma,(c^{a_1},\dots,c^{a_n})}.V_{w(\lambda)}\right)\right)^{ss}$$

so for $u \in (U_q)_0$:

$$\operatorname{Tr} u|_{\mathcal{V}_\nu} = \sum_{c\in\mathbb{C}^*:c^\beta=\nu^{-1}\lambda} \operatorname{Tr}\left(\varphi_{F_\Sigma,(c^{a_1},\dots,c^{a_n})}(T_w^{-1}(u))\right)|_{V_{w(\lambda)}}$$

(note that $c^{w(\beta)} = w(\nu^{-1}\lambda)$ if and only if $c^\beta = \nu^{-1}\lambda$ since $c^{w(\beta)} = w(c^\beta)$).

Set $p(c) = \operatorname{Tr}\left(\varphi_{F_\Sigma,(c^{a_1},\dots,c^{a_n})}(T_w^{-1}(u))\right)|_{V_{w(\lambda)}}$. p is Laurent polynomial in c and $p(q^i) = \operatorname{Tr} u|_{L_{q^{-i\beta}\lambda}}$ for $i \in \mathbb{N}$.

On the other hand we can show similarly that

$$\begin{aligned}&\operatorname{Tr} u|_{\left(\left(\bigoplus_{t\in\mathbb{C}^*/q^{\mathbb{Z}}} \varphi_{F_\beta,t}.L_{F_\beta}\right)^{ss}\right)_\nu}\\ &= \sum_{c\in\mathbb{C}^*:c^\beta=\nu^{-1}\lambda} \operatorname{Tr}\left(\varphi_{F_\beta,c}(u)\right)|_{(L_{F_\beta})_\lambda}.\end{aligned}$$

Similarly $\operatorname{Tr}\left(\varphi_{F_\beta,c}(u)\right)|_{(L_{F_\beta})_\lambda}$ is Laurent polynomial in c and equal to $\operatorname{Tr} u|_{L_{q^{-i\beta}\lambda}}$ for $c = q^i$, $i \in \mathbb{N}$. So $\operatorname{Tr}\left(\varphi_{F_\beta,c}(u)\right)|_{(L_{F_\beta})_\lambda} = p(c)$. We conclude that $\operatorname{Tr}^{\mathcal{V}} = \operatorname{Tr}^{\left(\left(\bigoplus_{t\in\mathbb{C}^*/q^{\mathbb{Z}}} \varphi_{F_\beta,t}.L_{F_\beta}\right)^{ss}\right)}$ so $\mathcal{V} \cong \left(\bigoplus_{t\in\mathbb{C}^*/q^{\mathbb{Z}}} \varphi_{F_\beta,t}.L_{F_\beta}\right)^{ss}$ as U_q-modules by Lemma 5.1.$\square$

For any $\lambda \in X$ there is a unique simple highest weight module which we call $L(\lambda)$. It is the unique simple quotient of the Verma module $M(\lambda) := U_q \otimes_{U_q^{\geq 0}} \mathbb{C}_\lambda$ where $\mathbb{C}_\lambda$ is the 1-dimensional $U_q^{\geq 0}$-module with U_q^+ acting trivially and U_q^0 acting like λ. Let $\rho = \frac{1}{2}\sum_{\beta\in\Phi^+}\beta$. In the following we use the dot action on X. For $w \in W$, $w.\lambda := q^{-\rho}w(q^\rho\lambda)$.

Proposition 5.10 *Let $\lambda \in X$ be such that $L(\lambda)$ is admissible. Let $\alpha \in \Pi$. Assume $\lambda(K_\alpha) \notin \pm q^{\mathbb{N}}$. Let $a = \frac{2}{(\alpha|\alpha)}$. If $a = \frac{1}{2}$ choose a squareroot $\lambda(K_\alpha)^{\frac{1}{2}}$ of $\lambda(K_\alpha)$. Then*

- *$-\alpha \in T_{L(\lambda)}$.*
- *$L(s_\alpha.\lambda)$ is admissible.*
- *${}^{s_\alpha}L(s_\alpha.\lambda)$ is a subquotient of the U_q-module $L(\lambda)_{F_\alpha}$.*
- *$L(s_\alpha.\lambda)$ and ${}^{s_\alpha}L(\lambda)$ are subquotients of the U_q-module. $\varphi_{F_\alpha,\lambda(K_\alpha)^a}.L(\lambda)_{F_\alpha}$.*

Proof. $\lambda(K_\alpha) \notin \pm q_\alpha^{\mathbb{N}}$ implies that $-\alpha \subset T_{L(\lambda)}$ since for $i \in \mathbb{N}$:

$$E_\alpha^{(i)} F_\alpha^{(i)} v_\lambda = \prod_{j=1}^{i} \frac{q_\alpha^{j-1}\lambda(K_\alpha) - q_\alpha^{1-j}\lambda(K_\alpha)^{-1}}{q_\alpha^j - q_\alpha^{-j}} v_\lambda .$$

This is only zero for an $i \in \mathbb{N}$ when $\lambda(K_\alpha) \in \pm q_\alpha^{\mathbb{N}}$.

Let $v_\lambda \in L(\lambda)$ be a highest weight vector. Denote the vector $\varphi_{F_\alpha,\lambda(K_\alpha)^a}.F_\alpha v_\lambda \in \varphi_{F_\alpha,\lambda(K_\alpha)}.L(\lambda)_{F_\Sigma}$ as $v_{s_\alpha.\lambda}$. This is a highest weight vector of weight $s_\alpha.\lambda$: For $\mu \in Q$:

$$\begin{aligned} K_\mu v_{s_\alpha.\lambda} =& K_\mu \varphi_{F_\alpha,\lambda(K_\alpha)^a}.F_\alpha v_\lambda \\ =& \varphi_{F_\alpha,q\lambda(K_\alpha)^a}.\left(\left(q\lambda(K_\alpha)^a\right)^{-(\mu|\alpha)} \lambda(K_\mu) F_\alpha v_\lambda \right) \\ =& q^{-(\mu|\alpha)} \lambda\left(K_\alpha^{-\langle \mu,\alpha^\vee\rangle} K_\mu \right) \varphi_{F_\alpha,q_\alpha\lambda(K_\alpha)}.F_\alpha v_\lambda \\ =& q^{-(\mu|\alpha)} (s_\alpha\lambda)(K_\mu) v_{s_\alpha.\lambda} \\ =& s_\alpha.\lambda(K_\mu) v_{s_\alpha.\lambda}. \end{aligned}$$

For $\alpha' \in \Pi \backslash \{\alpha\}$

$$E_{\alpha'} \varphi_{F_\alpha,\lambda(K_\alpha)^a}.v_\lambda = \varphi_{F_\alpha,\lambda(K_\alpha)^a}.E_{\alpha'} v_\lambda$$

and for $\alpha' = \alpha$ we have by the formula in the proof of Lemma 4.9

$$\begin{aligned} & E_\alpha \varphi_{F_\alpha,\lambda(K_\alpha)^a}.F_\alpha v_\lambda \\ & \qquad = \varphi_{F_\alpha,\lambda(K_\alpha)^a}.F_\alpha \varphi_{F_\alpha,q\lambda(K_\alpha)^a}(E_\alpha) v_\lambda \\ & \qquad = \varphi_{F_\alpha,\lambda(K_\alpha)^a}.F_\alpha \left(E_\alpha + F_\alpha^{-1} \frac{q_\alpha (q_\alpha\lambda(K_\alpha))^{-1} K_\alpha - q_\alpha^{-1} q_\alpha \lambda(K_\alpha) K_\alpha^{-1}}{(q_\alpha - q_\alpha^{-1})^2} \right) v_\lambda \\ & \qquad = 0. \end{aligned}$$

So $v_{s_\alpha.\lambda}$ is a highest weight vector of weight $s_\alpha.\lambda$ hence $L(s_\alpha.\lambda)$ is a subquotient of $\varphi_{F_\alpha,\lambda(K_\alpha)^a}.L(\lambda)_{F_\alpha}$. Since $L(s_\alpha.\lambda)$ is a subquotient of $\varphi_{F_\alpha,\lambda(K_\alpha)^a}.L(\lambda)_{F_\alpha}$ it is admissible by Lemma 4.22.

Consider ${}^{\overline{s_\alpha}}\left(\varphi_{F_\alpha,\lambda(K_\alpha)^a}.L(\lambda)_{F_\alpha} / (U_q v_{s_\alpha.\lambda}) \right)$ and the vector

$$v' = F_\alpha^{-1} v_{s_\alpha.\lambda} + U_q v_{s_\alpha.\lambda} \in {}^{\overline{s_\alpha}}\left(\varphi_{F_\alpha,\lambda(K_\alpha)^a}.L(\lambda)_{F_\alpha} / (U_q v_{s_\alpha.\lambda}) \right).$$

Then $E_\beta v' = 0$ for all $\beta \in \Pi$: First of all

$$\begin{aligned} E_\alpha \cdot v' =& T_{s_\alpha}^{-1}(E_\alpha) v' \\ =& -K_\alpha F_\alpha v' \\ =& -K_\alpha v_{s_\alpha.\lambda} + U_q v_{s_\alpha.\lambda} \\ =& 0. \end{aligned}$$

For $\beta \in \Pi \backslash \{\alpha\}$

$$\begin{aligned}
E_\beta \cdot v' =& T_{s_\alpha}^{-1}(E_\beta)v'. \\
=& \sum_{i=0}^{-\langle \beta,\alpha^\vee \rangle} (-1)^i q_\alpha^{-i} E_\alpha^{(i)} E_\beta E_\alpha^{(-\langle \beta,\alpha^\vee \rangle - i)} v' \\
=& (-1)^{\langle \beta,\alpha^\vee \rangle} q_\alpha^{\langle \beta,\alpha^\vee \rangle} E_\alpha^{(-\langle \beta,\alpha^\vee \rangle)} E_\beta v' \\
=& (-1)^{\langle \beta,\alpha^\vee \rangle} q_\alpha^{\langle \beta,\alpha^\vee \rangle} E_\alpha^{(-\langle \beta,\alpha^\vee \rangle)} F_\alpha^{-1} E_\beta v_{s_\alpha.\lambda} + U_q v_{s_\alpha.\lambda} \\
=& 0
\end{aligned}$$

since $E_\alpha v' = 0$ and $E_\beta v_{s_\alpha.\lambda} = 0$ by the above.

So v' is a highest weight vector and v' has weight λ: For $\mu \in Q$:

$$\begin{aligned}
K_\mu \cdot v' =& K_{s_\alpha \mu} v' \\
=& K_{s_\alpha \mu} F_\alpha^{-1} v_{s_\alpha.\lambda} + U_q v_{s_\alpha.\lambda} \\
=& q^{(s_\alpha(\mu)|\alpha)} s_\alpha.\lambda(K_{s_\alpha \mu}) F_\alpha^{-1} v_{s_\alpha.\lambda} + U_q v_{s_\alpha.\lambda} \\
=& \lambda(K_\mu) F_\alpha^{-1} v_{s_\alpha.\lambda} + U_q v_{s_\alpha.\lambda}.
\end{aligned}$$

So $L(\lambda)$ is a subquotient of ${}^{\overline{s_\alpha}}(\varphi_{F_\alpha,\lambda(K_\alpha)^a}.L(\lambda)_{F_\alpha})$ hence ${}^{s_\alpha}L(\lambda)$ is a subquotient of $\varphi_{F_\alpha,\lambda(K_\alpha)^a}.L(\lambda)_{F_\alpha}$. Consider the vector

$$v'' = F_\alpha^{-1} v_\lambda + U_q v_\lambda \in {}^{\overline{s_\alpha}} \left(L(\lambda)_{F_\alpha} / (U_q v_\lambda) \right).$$

By an argument analog to above we get $E_\beta \cdot v'' = 0$ for all $\beta \in \Pi \backslash \{\alpha\}$ since E_β and F_α^{-1} commutes and v_λ is a highest weight vector. We get $E_\alpha \cdot v'' = 0$ by the following:

$$\begin{aligned}
E_\alpha \cdot v'' =& T_{s_\alpha}^{-1}(E_\alpha) v'' \\
=& - K_\alpha F_\alpha v'' \\
=& - q^{-2} F_\alpha K_\alpha F_\alpha^{-1} v_\lambda + U_q v_\lambda \\
=& 0.
\end{aligned}$$

So v'' is a highest weight vector in ${}^{\overline{s_\alpha}} \left(L(\lambda)_{F_\alpha} / (U_q v_\lambda) \right)$. v'' has weight $s_\alpha.\lambda$: For $\mu \in Q$:

$$\begin{aligned}
K_\mu \cdot v'' =& K_{s_\alpha \mu} v'' \\
=& K_{s_\alpha \mu} F_\alpha^{-1} v_\lambda + U_q v_\lambda \\
=& q^{(s_\alpha(\mu)|\alpha)} \lambda(K_{s_\alpha \mu}) v'' \\
=& (q^{-\alpha} s_\alpha \lambda)(K_\mu) v''.
\end{aligned}$$

Hence $L(s_\alpha.\lambda)$ is a subquotient of ${}^{\overline{s_\alpha}}L(\lambda)_{F_\Sigma}$ and therefore ${}^{s_\alpha}L(s_\alpha.\lambda)$ is a subquotient of $L(\lambda)_{F_\Sigma}$. ☐

Lemma 5.11 *Let $\lambda \in X$ be such that $L(\lambda)$ is an infinite dimensional admissible module of degree d. Let $\alpha \in \Pi$. Then*

$$\mathcal{EXT}(L(\lambda)) \cong \mathcal{EXT}({}^{s_\alpha}L(\lambda))$$

and if $\lambda(K_\alpha) \notin \pm q_\alpha^{\mathbb{N}}$ then $\mathcal{EXT}(L(\lambda))$ contains $L(s_\alpha.\lambda)$ and ${}^{s_\alpha}L(s_\alpha.\lambda)$ as U_q-submodules, where $s_\alpha.\lambda := q^{-\rho} s_\alpha(q^\rho \lambda) = q^{-\alpha} s_\alpha \lambda$.

Proof. Assume first that $\lambda(K_\alpha) \notin \pm q_\alpha^{\mathbb{N}}$. By Proposition 5.10 the U_q-module $\bigoplus_{t\in\mathbb{C}^*/q^{\mathbb{Z}}} \varphi_{F_\alpha,t}.L(\lambda)_{F_\alpha}$ contains $L(s_\alpha.\lambda)$, ${}^{s_\alpha}L(\lambda)$ and ${}^{s_\alpha}L(s_\alpha.\lambda)$ as subquotients. By Proposition 5.9 and Proposition 5.6 this finishes the proof of the claim when $\lambda(K_\alpha) \notin \pm q_\alpha^{\mathbb{N}}$.

Assume now that $\lambda(K_\alpha) = \pm q_\alpha^k$ for some $k \in \mathbb{N}$: If $\lambda(K_\alpha) = q_\alpha^k$ it is easy to prove that $L(\lambda) \cong {}^{s_\alpha}L(\lambda)$. Assume from now on that $\lambda(K_\alpha) = -q_\alpha^k$. We have

$$\mathcal{EXT}(L(\lambda)) = \left(\bigoplus_{t\in(\mathbb{C}^*)/q^{\mathbb{Z}^n}} \varphi_{F_\Sigma,t}.L(\lambda)_{F_\Sigma} \right)^{ss}$$

for some set of commuting roots $\Sigma = \{\beta_1,\dots,\beta_n\}$ that is a basis of Q with $-\Sigma \subset T_{L(\lambda)}$. Since Σ is a basis of Q there exists $a_1,\dots,a_n \in \mathbb{Z}$ such that $\alpha = \sum_{i=1}^n a_i\beta_i$. Let v_λ be a highest weight vector in $L(\lambda)$. We will show that $v_0 := \varphi_{F_\Sigma,((-1)^{a'_1},\dots,(-1)^{a'_n})}.F_\alpha^i v_\lambda \in {}^{\overline{s_\alpha}}\mathcal{EXT}(L(\lambda))$ is a highest weight vector of weight λ where $a'_i = \frac{2a_i}{(\alpha|\alpha)}$. This will imply $\mathcal{EXT}({}^{s_\alpha}L(\lambda)) \cong \mathcal{EXT}(L(\lambda))$ by Proposition 5.6. The weight of v_0: Let $\mu \in Q$:

$$\begin{aligned}
K_\mu \cdot v_0 =& K_{s_\alpha(\mu)}\varphi_{F_\Sigma,((-1)^{a'_1},\dots,(-1)^{a'_n})}.F_\alpha^i v_\lambda \\
=&(-1)^{\left(\sum_{i=1}^n a'_i\beta_i|\mu\right)} q^{i(\alpha|\mu)} \lambda(K_\mu K_\alpha^{-\langle\mu,\alpha^\vee\rangle})\varphi_{F_\Sigma,((-1)^{a'_1},\dots,(-1)^{a'_n})}.F_\alpha^i v_\lambda \\
=&(-1)^{\langle\mu,\alpha^\vee\rangle} q_\alpha^{i\langle\mu,\alpha^\vee\rangle}(-q_\alpha^i)^{-\langle\mu,\alpha^\vee\rangle}\lambda(K_\mu)v_0 \\
=&\lambda(K_\mu)v_0.
\end{aligned}$$

By Proposition 4.10 $\varphi_{F_\beta,(-1)^{\frac{2}{(\beta|\beta)}}}(E_{\alpha'}) = E_{\alpha'}$ and $\varphi_{F_\beta,(-1)^{\frac{2}{(\beta|\beta)}}}(F_{\alpha'}) = \pm F_{\alpha'}$ for any $\alpha' \in \Pi$ and any $\beta \in \Phi^+$. So $\varphi_{F_\Sigma,((-1)^{a'_1},\dots,(-1)^{a'_n})}(E_\beta)$, $\beta \in \Pi\backslash\{\alpha\}$ and $\varphi_{F_\Sigma,((-1)^{a'_1},\dots,(-1)^{a'_n})}(F_\alpha)$ kills $F_\alpha^i v_\lambda \in L(\lambda)$ because E_β and F_α does. Hence E_β, $\beta \in \Pi$ kills v_0 by the same argument as in the proof of Proposition 5.10 when proving that v' is a highest weight vector. $\square$

Theorem 5.12 *Let L be an infinite dimensional admissible simple module of degree d. Then the U_q-module $\mathcal{EXT}(L)$ contains an infinite dimensional admissible simple highest weight module $L(\lambda)$ of degree d for some weight $\lambda \in X$. Furthermore for any $x \in W$:*

$$^{x}\mathcal{EXT}(L) \cong \mathcal{EXT}(L).$$

Proof. Let $w \in W$ be such that $w(F_L\backslash F_L^s) \subset \Phi^+$ and $w(T_L\backslash T_L^s) \subset \Phi^-$. Set $L' = {}^{\overline{w^{-1}}}L$ (then ${}^{w^{-1}}L' = L$). We will show the result first for L' by induction on $|T_{L'}^+|$. If $|T_{L'}^+| = 0$ then L' is itself a highest weight module. Assume $|T_{L'}^+| > 0$. Let $\beta \in T_{L'}^+$. Then $\beta \in T_{L'}^s$ since $T_{L'}\backslash T_{L'}^s \subset \Phi^-$. So $-\beta \in T_{L'}$. Then by Lemma 4.23 there exists a $b \in \mathbb{C}^*$ such that $\varphi_{F_\beta,b}.L'_{F_\beta}$ contains a U_q-submodule L'' with $T_{L''} \subset T_{L'}$ and $\beta \notin T_{L''}$. By Proposition 5.9 and Proposition 5.6 $\mathcal{EXT}(L') \cong \mathcal{EXT}(L'')$ as U_q-modules. By induction $\mathcal{EXT}(L'')$ contains an infinite dimensional admissible simple highest weight module $L(\lambda)$ for some λ. So $\mathcal{EXT}(L') \cong \mathcal{EXT}(L(\lambda))$ by Proposition 5.6. Choose a reduced expression

$s_{i_r}\cdots s_{i_1}$ for w^{-1}. By Proposition 5.5 and Lemma 5.11

$$\begin{aligned}\mathcal{EXT}(L) \cong& \mathcal{EXT}({}^{w^{-1}}L')\\ \cong& {}^{w^{-1}}\mathcal{EXT}(L')\\ \cong& {}^{w^{-1}}\mathcal{EXT}(L(\lambda))\\ \cong& {}^{s_{i_r}\cdots s_{i_2}}\mathcal{EXT}({}^{s_{i_1}}L(\lambda))\\ \cong& {}^{s_{i_r}\cdots s_{i_2}}\mathcal{EXT}(L(\lambda))\\ &\vdots\\ \cong& \mathcal{EXT}(L(\lambda)).\end{aligned}$$

So $\mathcal{EXT}(L)$ contains a simple highest weight module $L(\lambda)$. For any $x \in W$ we can do as above to show ${}^{x}\mathcal{EXT}(L) \cong \mathcal{EXT}({}^{x}L(\lambda)) \cong \mathcal{EXT}(L(\lambda)) \cong \mathcal{EXT}(L)$.□

Corollary 5.13 *Let L be a simple torsion free module. Then there exists a set of commuting roots Σ that is a basis of Q with corresponding Ore subset F_Σ, a $\lambda \in X$ and $\mathbf{b} \in (\mathbb{C}^*)^n$ such that $-\Sigma \subset T_{L(\lambda)}$ and $L \cong \varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$*

Proof. By Theorem 5.12 $\mathcal{EXT}(L) \cong \mathcal{EXT}(L(\lambda))$ for some $\lambda \in X$. So L is a U_q-submodule of $\mathcal{EXT}(L(\lambda))$. Let Σ be a set of commuting roots such that $-\Sigma \subset L(\lambda)$ (exists by Lemma 4.16 by setting $w = e$, the neutral element in W) then

$$\mathcal{EXT}(L(\lambda)) = \left(\bigoplus_{t\in(\mathbb{C}^*)^n/q^{\mathbb{Z}^n}} \varphi_{F_\Sigma,t}.L(\lambda)_{F_\Sigma} \right)^{ss}.$$

Since L is simple we must have that L is a submodule of $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ for some $\mathbf{b} \in (\mathbb{C}^*)^n$. By Proposition 5.6 and Lemma 4.22 $\dim\left(\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}\right)_\lambda = \dim L_\lambda$ for all $\lambda \in \operatorname{wt} L$ so we have $L \cong \varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$. □

So to classify torsion free simple modules we need to classify the admissible infinite dimensional simple highest weight modules $L(\lambda)$ and then we need to determine the $t \in (\mathbb{C}^*)^n/q^{\mathbb{Z}^n}$ such that $\varphi_{F_\Sigma,t}.L(\lambda)_{F_\Sigma}$ is simple. Furthermore we have that if there exists an admissible infinite dimensional simple module then there exists a torsion free simple module. In the classical case torsion free modules only exists if $\mathfrak{g}$ is of type A or C so we expect the same to be true in the quantum group case. We show this in Section 8.5.

6 Classification of simple torsion free $U_q(\mathfrak{sl}_2)$-modules

In this section let $\mathfrak{g} = \mathfrak{sl}_2$. In this case there is a single simple root α. It is natural to identify X with $\mathbb{C}^*$ via $\lambda \mapsto \lambda(K_\alpha)$. We define $F = F_\alpha$, $E = E_\alpha$ and $K^{\pm 1} = K_\alpha^{\pm 1}$. Let $\lambda \in \mathbb{C}^*\backslash\{\pm q^{\mathbb{N}}\}$ and consider the simple highest weight module $L(\lambda)$. Let $0 \neq v_0 \in L(\lambda)_\lambda$. $\operatorname{wt} L = q^{-2\mathbb{N}}\lambda$ so $L(\lambda)$ is an admissible infinite dimensional highest weight module. Thus $\mathcal{EXT}(L(\lambda))$ contains a torsion free module by Theorem 5.8. Let $b \in \mathbb{C}^*$. We will describe the action on the module $\varphi_{F,b}.L(\lambda)_{(F)}$ and determine exactly for which b's $\varphi_{F,b}.L(\lambda)_{(F)}$ is torsion free.

Let $v_i = F^i\varphi_{F,b}.v_0$ for all $i \in \mathbb{Z}$. Then we have for $i \in \mathbb{Z}$

$$\begin{aligned} Fv_i =& v_{i+1} \\ K^{\pm 1}v_i =& q^{-2i}b^{\mp 2}\lambda v_i \\ Ev_i =& \frac{(q^i b - q^{-i}b^{-1})(q^{1-i}b^{-1}\lambda - q^{i-1}b\lambda^{-1})}{(q-q^{-1})^2} v_{i-1}. \end{aligned}$$

We see that unless $b = \pm q^i$ or $b = \pm q^i\lambda$ for some $i \in \mathbb{Z}$ then $\varphi_{F,b}.L(\lambda)_{(F)}$ is torsion free. In this case we see that $\varphi_{F,-b} = \varphi_{F,b}$ since for all $u \in U_q(\mathfrak{sl}_2)$, $\varphi_{F,b}(u)$ is Laurent polynomial in b^2.

So in this case $\mathcal{EXT}(L(\lambda))$ contains a maximum of four different simple submodules which are *not* torsion free: We have $(\varphi_{F,\pm q^i}.L(\lambda)_{(F)})^{ss} \cong (L(\lambda)_{(F)})^{ss} \cong L(\lambda) \oplus {}^{s_\alpha}L(s_\alpha.\lambda)$ (which can be seen directly from the calculations but also follows from Corollary 4.20 and the fact that $\varphi_{F,-b} = \varphi_{F,b}$) and $(\varphi_{F,\pm q^i\lambda}.L(\lambda)_{(F)})^{ss} \cong (L(s_\alpha.\lambda)_{(F)})^{ss} \cong L(s_\alpha.\lambda) \oplus {}^{s_\alpha}L(\lambda)$ if $\lambda \notin \pm q^{\mathbb{Z}}$.

The weights of $\varphi_{F,b}.L(\lambda)_{(F)}$ are $b^{-\alpha}\operatorname{wt} L(\lambda)_{(F)} = q^{2\mathbb{Z}}b^{-2}\lambda$. Suppose we want to find a torsion free $U_q(\mathfrak{sl}_2)$-modules with integral weights. Then we just need to find $\lambda, b \in \mathbb{C}^*$ such that $\lambda \notin \pm q^{\mathbb{Z}_{\geq 0}}$, $b \notin \pm q^{\mathbb{Z}}$ and $b \notin \pm q^{\mathbb{Z}}\lambda$ such that $b^{-2}\lambda \in q^{\mathbb{Z}}$. For example choose a square root $q^{1/2}$ of q and set $\lambda = q^{-1}$ and $b = q^{1/2}$. Then we have a torsion free module $L = \operatorname{span}_{\mathbb{C}}\{v_i | i \in \mathbb{Z}\}$ with action given by:

$$\begin{aligned} Fv_i =& v_{i+1} \\ Kv_i =& q^{-2i-2}v_i \\ Ev_i =& \frac{(q^{1/2+i} - q^{-1/2-i})(q^{-1/2-i} - q^{i+1/2})}{(q-q^{-1})^2} v_{i-1} \\ =& \frac{q(q^{-i-1} - q^i)^2}{(q-q^{-1})^2} v_{i-1}. \end{aligned}$$

In this paper we only focus on quantized enveloping algebras over $\mathbb{C}$ but note that we can define, for a general field $\mathbb{F}$ with $q \in \mathbb{F}\backslash\{0\}$ a non-root of unity, a simple torsion free $U_{\mathbb{F}}(\mathfrak{sl}_2)$-module with integral weights by the above formulas (here $U_{\mathbb{F}}(\mathfrak{sl}_2) = U_A \otimes_A \mathbb{F}$ where $\mathbb{F}$ is considered an A-algebra by sending v to q).

7 An example for $U_q(\mathfrak{sl}_3)$

In this section we will show how we can construct a specific torsion free simple module for $U_q(\mathfrak{sl}_3)$. In Section 9 we classify all torsion free $U_q(\mathfrak{sl}_n)$-modules with $n \geq 3$ so this example is of course included there. If you are only interested in the general classification you can skip this section but the calculations in this section gives a taste of the calculations needed in the general case in Section 9 and they show a phenomona that does not happen in the classical case.

Let α_1 and α_2 be the two simple roots of the root system. We will consider the set of commuting roots $\Sigma = \{\beta_1, \beta_2\}$ where $\beta_1 = \alpha_1$ and $\beta_2 = \alpha_1 + \alpha_2$. Set $F_{\beta_1} := F_{\alpha_1}$ and $F_{\beta_2} := T_{s_1}(F_{\alpha_2}) = F_{\alpha_2}F_{\alpha_1} - qF_{\alpha_1}F_{\alpha_2} = [F_{\alpha_2}, F_{\alpha_1}]_q$. We have $(\beta_1|\beta_2) = 1$ and $0 = [F_{\beta_2}, F_{\beta_1}]_q = F_{\beta_2}F_{\beta_1} - q^{-1}F_{\beta_1}F_{\beta_2}$ or equivalently $F_{\beta_1}F_{\beta_2} = qF_{\beta_2}F_{\beta_1}$. Let $\lambda \in X$ be determined by $\lambda(K_{\alpha_1}) = q^{-1}$ and $\lambda(K_{\alpha_2}) = 1$. Then $M(s_{\alpha_1}.\lambda)$ is a submodule of $M(\lambda)$ and $L(\lambda) = M(\lambda)/M(s_{\alpha_2}.\lambda) =$

$M(\lambda)/M(q^{-\alpha_2}\lambda)$ is admissible of degree 1. Let $\xi = e^{2\pi i/3}$. We will show that $\varphi_{F_\Sigma,(\xi,\xi)}.L(\lambda)_{F_\Sigma}$ is a torsion free module. We have here a phenomona that does not happen in the classical case: $\operatorname{wt} L(\lambda)_{F_\Sigma} = \operatorname{wt} \varphi_{F_\Sigma,(\xi,\xi)}.L(\lambda)_{F_\Sigma}$ but $L(\lambda)_{F_\Sigma} \not\cong \varphi_{F_\Sigma,(\xi,\xi)}.L(\lambda)_{F_\Sigma}$ as U_q-modules since one is simple and torsion free and the other isn't (compare to [Mat00, Section 10] where Mathieu classifies the torsion free simple modules by determining for a coherent family $\mathcal{M}$ for which cosets $t \in \mathfrak{h}^*/Q$, $\mathcal{M}[t]$ is torsion free).

We will show that E_{α_1} and E_{α_2} act injectively on the module $\varphi_{F_\Sigma,(\xi,\xi)}.L(\lambda)_{F_\Sigma}$. So we need to calculate $\varphi_{F_\Sigma,(\xi,\xi)}(E_{\alpha_1})$ and $\varphi_{F_\Sigma,(\xi,\xi)}(E_{\alpha_2})$. $\varphi_{F_\Sigma,(\xi,\xi)} = \varphi_{F_{\beta_1},\xi} \circ \varphi_{F_{\beta_2},\xi}$. We have

$$\begin{aligned}[E_{\alpha_1}, F_{\beta_2}] =& F_{\alpha_2}[E_{\alpha_1}, F_{\alpha_1}] - q[E_{\alpha_1}, F_{\alpha_1}]F_{\alpha_2} \\ =& F_{\alpha_2}\frac{K_{\alpha_1} - K_{\alpha_1}^{-1}}{q-q^{-1}} - qF_{\alpha_2}\frac{qK_{\alpha_1} - q^{-1}K_{\alpha_1}^{-1}}{q-q^{-1}} \\ =& F_{\alpha_2}\frac{K_{\alpha_1} - q^2K_{\alpha_1}}{q-q^{-1}} \\ =& -F_{\alpha_2}q\frac{q-q^{-1}}{q-q^{-1}}K_{\alpha_1} \\ =& -qF_{\alpha_2}K_{\alpha_1}.\end{aligned}$$

We can show by induction that

$$[E_{\alpha_1}, F_{\beta_2}^j] = -q^{2-j}[j]F_{\beta_2}^{j-1}F_{\alpha_2}K_{\alpha_1}$$

for any $j \in \mathbb{N}$. Using that $\varphi_{F_{\beta_2},b}(E_{\alpha_1})$ is Laurent polynomial and equal to $F_{\beta_2}^{-j}E_{\alpha_1}F_{\beta_2}^j$ for $b = q^j$ we get

$$\varphi_{F_{\beta_2},b}(E_{\alpha_1}) = E_{\alpha_1} - q^2b^{-1}\frac{b-b^{-1}}{q-q^{-1}}F_{\beta_2}^{-1}F_{\alpha_2}K_{\alpha_1}.$$

We have $F_{\beta_2}F_{\beta_1} = q^{-1}F_{\beta_1}F_{\beta_2}$ so $F_{\beta_1}^{-i}F_{\beta_2}F_{\beta_1}^i = q^{-i}F_{\beta_2}$ thus $\varphi_{F_{\beta_1},b}(F_{\beta_2}^{-1}) = bF_{\beta_2}^{-1}$. We have

$$\begin{aligned}\varphi_{F_{\alpha_1},b}(F_{\alpha_2}) =& bF_{\alpha_2} - \frac{b-b^{-1}}{q-q^{-1}}F_{\alpha_1}^{-1}(qF_{\alpha_1}F_{\alpha_2} - F_{\alpha_2}F_{\alpha_1}) \\ =& bF_{\alpha_2} + \frac{b-b^{-1}}{q-q^{-1}}F_{\alpha_1}^{-1}F_{\beta_2}\end{aligned}$$

and

$$\begin{aligned}
&\varphi_{F_{\beta_1},b_1}(\varphi_{F_{\beta_2},b_2}(E_{\alpha_1}))\\
&\quad=\varphi_{F_{\alpha_1},b_1}\left(E_{\alpha_1}-q^2b_2^{-1}\frac{b_2-b_2^{-1}}{q-q^{-1}}F_{\beta_2}^{-1}F_{\alpha_2}K_{\alpha_1}\right)\\
&\quad=E_{\alpha_1}+F_{\alpha_1}^{-1}\frac{(b_1-b_1^{-1})(qb_1^{-1}K_{\alpha_1}-q^{-1}b_1K_{\alpha_1}^{-1})}{(q-q^{-1})^2}\\
&\qquad-q^2b_2^{-1}\frac{b_2-b_2^{-1}}{q-q^{-1}}b_1F_{\beta_2}^{-1}\left(b_1F_{\alpha_2}+\frac{b_1-b_1^{-1}}{q-q^{-1}}F_{\alpha_1}^{-1}F_{\beta_2}\right)b_1^{-2}K_{\alpha_1}\\
&\quad=E_{\alpha_1}+F_{\alpha_1}^{-1}\frac{(b_1-b_1^{-1})(qb_1^{-1}K_{\alpha_1}-q^{-1}b_1K_{\alpha_1}^{-1})}{(q-q^{-1})^2}\\
&\qquad-q^2b_2^{-1}\frac{b_2-b_2^{-1}}{q-q^{-1}}F_{\beta_2}^{-1}F_{\alpha_2}K_{\alpha_1}\\
&\qquad-qb_1^{-1}b_2^{-1}\frac{(b_2-b_2^{-1})(b_1-b_1^{-1})}{(q-q^{-1})^2}F_{\alpha_1}^{-1}K_{\alpha_1}\\
&\quad=E_{\alpha_1}+b_2^{-1}F_{\alpha_1}^{-1}\frac{(b_1-b_1^{-1})(qb_1^{-1}b_2^{-1}K_{\alpha_1}-q^{-1}b_1b_2K_{\alpha_1}^{-1})}{(q-q^{-1})^2}\\
&\qquad-q^2b_2^{-1}\frac{b_2-b_2^{-1}}{q-q^{-1}}F_{\beta_2}^{-1}F_{\alpha_2}K_{\alpha_1}.
\end{aligned}$$

Let v'_λ be a highest weight vector in $L(\lambda)$ and set $v_\lambda = 1\otimes v'_\lambda \in L(\lambda)_{F_\Sigma}$. We have $F_{\alpha_2}v_\lambda = 0$ by construction so we have

$$\varphi_{F_\Sigma,(b_1,b_2)}(E_{\alpha_1})v_\lambda = b_2^{-1}\frac{(b_1-b_1^{-1})(b_1^{-1}b_2^{-1}-b_1b_2)}{(q-q^{-1})^2}F_{\alpha_1}^{-1}v_\lambda.$$

$\varphi_{F_\Sigma,(c_1,c_2)}.L(\lambda)_{F_\Sigma}$ is spanned by the elements $F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(c_1,c_2)}.v_\lambda$, $i,j\in\mathbb{Z}$ because every weight space is one-dimensional and $F_{\beta_1}^iF_{\beta_2}^j$ acts injectively. Since

$$\begin{aligned}
F_{\beta_2}^{-j}F_{\beta_1}^{-i}E_{\alpha_1}F_{\beta_1}^iF_{\beta_2}^j &= F_{\beta_1}^{-i}F_{\beta_2}^{-j}E_{\alpha_1}F_{\beta_2}^jF_{\beta_1}^i\\
&=\varphi_{F_{\beta_1},q^i}(\varphi_{F_{\beta_2},q^j}(E_{\alpha_1}))\\
&=\varphi_{F_\Sigma,(q^i,q^j)}(E_{\alpha_1})
\end{aligned}$$

we have

$$\begin{aligned}
&E_{\alpha_1}F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(c_1,c_2)}.v_\lambda\\
&\qquad=F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(q^i,q^j)}(E_{\alpha_1})\varphi_{F_\Sigma,(c_1,c_2)}.v_\lambda\\
&\qquad=F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(c_1,c_2)}.\varphi_{F_\Sigma,(q^ic_1,q^jc_2)}(E_{\alpha_1})v_\lambda\\
&\qquad=q^{-j}c_2^{-1}\frac{(q^ic_1-q^{-i}c_1^{-1})(q^{-i-j}c_1^{-1}c_2^{-1}-q^{i+j}c_1c_2)}{(q-q^{-1})^2}F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(c_1,c_2)}.F_{\alpha_1}^{-1}v_\lambda\\
&\qquad=\frac{(q^ic_1-q^{-i}c_1^{-1})(q^{-i-j}c_1^{-1}c_2^{-1}-q^{i+j}c_1c_2)}{(q-q^{-1})^2}F_{\beta_1}^{i-1}F_{\beta_2}^j\varphi_{F_\Sigma,(c_1,c_2)}.v_\lambda.
\end{aligned}$$

This is only zero when $c_1=\pm q^{-i}$ or $c_1c_2=\pm q^{-i-j}$. Set $c_1=c_2=e^{2\pi i/3}=:\xi$. Then we have shown that E_{α_1} acts injectively on $\varphi_{F_\Sigma,(\xi,\xi)}.L(\lambda)_{F_\Sigma}$.

Now we will show that E_{α_2} acts injectively on $F_{\Sigma,(\xi,\xi)}.L(\lambda)_{F_\Sigma}$. We can show by induction that

$$[E_{\alpha_2}, F_{\beta_2}^j] =[j]F_{\alpha_1}F_{\beta_2}^{j-1}K_{\alpha_2}^{-1}$$

so $\varphi_{F_{\beta_2},b}(E_{\alpha_2}) = E_{\alpha_2} + b\frac{b-b^{-1}}{q-q^{-1}}F_{\alpha_1}F_{\beta_2}^{-1}K_{\alpha_2}^{-1}$ and

$$\begin{aligned}\varphi_{F_\Sigma,(b_1,b_2)}(E_{\alpha_2}) =&\varphi_{F_{\beta_1},b_1}(\varphi_{F_{\beta_2},b_2}(E_{\alpha_2}))\\ =&E_{\alpha_2} + b_2\frac{b_2-b_2^{-1}}{q-q^{-1}}F_{\alpha_1}F_{\beta_2}^{-1}K_{\alpha_2}^{-1}.\end{aligned}$$

Thus

$$\begin{aligned}E_{\alpha_2}F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(c_1,c_2)}.v_\lambda =&F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(q^i,q^j)}(E_{\alpha_2})\varphi_{F_\Sigma,(c_1,c_2)}.v_\lambda\\ =&F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(c_1,c_2)}.\varphi_{F_\Sigma,(q^ic_1,q^jc_2)}(E_{\alpha_2})v_\lambda\\ =&F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(c_1,c_2)}.c_2\frac{q^jc_2-q^{-j}c_2^{-1}}{q-q^{-1}}F_{\alpha_1}F_{\beta_2}^{-1}K_{\alpha_2}^{-1}v_\lambda\\ =&q^{-j-1}c_2\frac{q^jc_2-q^{-j}c_2^{-1}}{q-q^{-1}}F_{\beta_1}^{i+1}F_{\beta_2}^{j-1}\varphi_{F_\Sigma,(c_1,c_2)}.v_\lambda.\end{aligned}$$

We see that this is nonzero only if $c_2 = \pm q^{-j}$ so again setting $c_1 = c_2 = \xi$ ensures that this is nonzero.

We have shown that the U_q-module $\varphi_{F_\Sigma,(\xi,\xi)}.L(\lambda)_{F_\Sigma}$ has a basis $F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(\xi,\xi)}.v_\lambda$, $i,j\in\mathbb{Z}$ and we have

$$\begin{aligned}F_{\beta_1}F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(\xi,\xi)}.v_\lambda =&F_{\beta_1}^{i+1}F_{\beta_2}^{j-1}\varphi_{F_\Sigma,(\xi,\xi)}.v_\lambda\\ F_{\beta_2}F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(\xi,\xi)}.v_\lambda =&q^{-j}F_{\beta_1}^iF_{\beta_2}^{j+1}\varphi_{F_\Sigma,(\xi,\xi)}.v_\lambda\\ E_{\alpha_1}F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(\xi,\xi)}.v_\lambda =&C_1F_{\beta_1}^{i-1}F_{\beta_2}^j\varphi_{F_\Sigma,(\xi,\xi)}.v_\lambda\\ E_{\alpha_1}E_{\alpha_2}F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(\xi,\xi)}.v_\lambda =&C_2F_{\beta_1}^iF_{\beta_2}^{j-1}\varphi_{F_\Sigma,(\xi,\xi)}.v_\lambda\end{aligned}$$

for some nonzero constants $C_1, C_2 \in \mathbb{C}^*$. We see that any of the basis vectors $F_{\beta_1}^iF_{\beta_2}^j\varphi_{F_\Sigma,(\xi,\xi)}.v_\lambda$ can be mapped injectively to any other basis vector $F_{\beta_1}^{i'}F_{\beta_2}^{j'}\varphi_{F_\Sigma,(\xi,\xi)}.v_\lambda$ by elements of U_q so $\varphi_{F_\Sigma,(\xi,\xi)}.L(\lambda)_{F_\Sigma}$ is a simple module. The module is torsion free by Proposition 2.5.

8 Classification of admissible simple highest weight modules

8.1 Preliminaries

In this section we prove some preliminary results with the goal to classify all admissible simple highest weight modules. We will only focus on non-integral weights since we have the following theorem from [AM15]:

Theorem 8.1 *Assume $q \in \mathbb{C}\backslash\{0\}$ is transcendental. Let $\lambda : U_q^0 \to \mathbb{C}$ be a weight such that $\lambda(K_\alpha) = q_\beta^i$ for some $i \in \mathbb{Z}$ for every $\alpha \in \Pi$ - i.e. $\lambda \in q^Q$. Say $\lambda = q^\mu$, $\mu \in Q$. Let $L_\mathbb{C}(\mu)$ denote the simple highest weight $\mathfrak{g}$-module of highest weight μ. Then the character of $L(\lambda)$ and $L_\mathbb{C}(\mu)$ are equal - i.e. for any $\nu \in Q$, $\dim L(\lambda)_{q^\nu\lambda} = \dim L_\mathbb{C}(\mu)_{\nu+\mu}$.*

Proof. [AM15, Corollary 6.3]. □

Extending to modules which are not of type 1 is done in the usual way (cf. e.g. [Jan96, Section 5.1–5.2]). The above theorem implies that the integral admissible simple highest weight modules can be classified from the classification of the classical admissible simple highest weight modules when q is transcendental. Hence we need only to consider weights $\lambda \in X$ such that $\lambda(K_\alpha) \notin \pm q^{\mathbb{Z}}$ for at least one $\alpha \in \Pi$ in this case. *So in the rest of the paper we will restrict our attention to the case when q is transcendental.* If a similar theorem is true for any non-root-of-unity q then the results in this paper extend to all non-root-of-unities but the author is not aware of any such result.

Theorem 8.2 *Let $\lambda \in X$. Then there exists a filtration of $M(\lambda)$, $M(\lambda) \supset M_1 \supset \cdots \supset M_r$ such that M_1 is the unique maximal submodule of $M(\lambda)$ and*

$$\sum_{i=1}^{r} \operatorname{ch} M_i = \sum_{\substack{\beta \in \Phi^+ \\ q^\rho \lambda(K_\beta) \in \pm q_\beta^{\mathbb{Z}>0}}} \operatorname{ch} M(s_\beta . \lambda)$$

The filtration is called the Jantzen filtration and the formula is called the Jantzen sum formula.

Proof. This is proved in [Jos95, Section 4.1.2-4.1.3]. A proof using twisting functors can also be found in [Ped15b, Theorem 6.3]. □

Definition 8.3 *Let $\lambda \in X$.*

$$A(\lambda) = \{\alpha \in \Pi | \lambda(K_\alpha) \notin \pm q_\alpha^{\mathbb{N}}\}.$$

Let $\gamma \in \Pi$.

$$D(\gamma) = \{\beta \in \Phi^+ | \beta = \sum_{\alpha \in \Pi} m_\alpha \alpha,\, m_\gamma > 0\}.$$

Lemma 8.4 *Let $\lambda \in X$. Let $\gamma \in \Pi$ be such that $\gamma \in A(\lambda)$. Then $-D(\gamma) \subset T_{L(\lambda)}$.*

Proof. Let $\beta = \sum_{\alpha \in \Pi} m_\alpha \alpha \in D(\gamma)$. We prove by induction over $\operatorname{ht} \beta = \sum_{\alpha \in \Pi} m_\alpha$ that $-\beta \in T_{L(\lambda)}$. If $\operatorname{ht} \beta = 1$ then $\beta = \gamma$ and $-\gamma \in T_{L(\lambda)}$ by Proposition 5.10.

Assume $\operatorname{ht} \beta > 1$. Then $\beta - \alpha \in \Phi^+$ for some $\alpha \in \Pi$. We have either $\alpha = \gamma$ or $\beta - \alpha \in D(\gamma)$. In either case we get $\beta = \beta' + \beta''$ for some $\beta', \beta'' \in \Phi^+$ with $\beta' \in D(\gamma)$ and $\operatorname{ht} \beta' < \operatorname{ht} \beta$. By induction $-\beta' \in T_{L(\lambda)}$. If $-\beta \in F_{L(\lambda)}$ then $-\beta' = -\beta + \beta'' \in F_{L(\lambda)}$ since $\Phi^+ \subset F_{L(\lambda)}$ and $F_{L(\lambda)}$ is closed (Proposition 2.6). A contradiction. So $-\beta \in T_{L(\lambda)}$. □

Lemma 8.5 *Let $\gamma \in \Pi$. $D(\gamma)$ generates Q.*

Proof. Let $\langle D(\gamma) \rangle$ be the subgroup of Q generated by $D(\gamma)$. Assume $\Pi \cap \langle D(\gamma) \rangle \neq \Pi$. Let $\alpha \notin \langle D(\gamma) \rangle$ be a simple root that is connected to an $\alpha' \in \langle D(\gamma) \rangle$ (possible since the Dynkin diagram of a simple Lie algebra is connected). Then $\alpha + \alpha' \in \langle D(\gamma) \rangle$. But then $\alpha = \alpha + \alpha' - \alpha' \in \langle D(\gamma) \rangle$. A contradiction. So $\langle D(\gamma) \rangle = Q$. □

Lemma 8.6 *Let $\lambda \in X$ be a non-integral weight. Assume that $L(\lambda)$ is admissible. Then $A(\lambda)$ is connected and $|A(\lambda)| \leq 2$.*

Proof. Assume $|A(\lambda)| \geq 2$. Let $\alpha, \alpha' \in A(\lambda)$ be two distinct elements. We will show that α and α' are connected. So assume $(\alpha|\alpha') = 0$ to reach a contradiction. $L(\lambda)$ is admissible of some degree d. By Lemma 5.11 and Proposition 5.6 ${}^{s_\alpha}L(s_\alpha.\lambda)$ is admissible of the same degree d ($L(s_\alpha.\lambda)$ is infinite dimensional since $s_\alpha.\lambda(K_{\alpha'}) = \lambda(K_{\alpha'}) \notin \pm q_{\alpha'}^{\mathbb{N}}$). Let Σ be a set of commuting roots that is a basis of Q such that $\alpha \in \Sigma$ and $-\Sigma \subset T_{L(\lambda)}$ (Lemma 4.16). By Proposition 5.10 ${}^{s_\alpha}L(s_\alpha.\lambda)$ is a subquotient of $L(\lambda)_{F_\Sigma}$. We claim that $\operatorname{Supp}_{\text{ess}}(L(\lambda)) \cap \operatorname{Supp}_{\text{ess}}({}^{s_\alpha}L(s_\alpha.\lambda)) \neq \emptyset$. If this is true then we have for $\nu \in \operatorname{Supp}_{\text{ess}}(L(\lambda)) \cap \operatorname{Supp}_{\text{ess}}({}^{s_\alpha}L(s_\alpha.\lambda))$, $L(\lambda)_\nu \cong (L(\lambda)_{F_\Sigma})_\nu \cong ({}^{s_\alpha}L(s_\alpha.\lambda))_\nu$ as $(U_q)_0$-modules by Lemma 4.22. But then by Theorem 2.7 $L(\lambda) \cong {}^{s_\alpha}L(s_\alpha.\lambda)$ which is clearly a contradiction by looking at the weights of the modules. So we will prove the claim that $\operatorname{Supp}_{\text{ess}}(L(\lambda)) \cap \operatorname{Supp}_{\text{ess}}({}^{s_\alpha}L(s_\alpha.\lambda)) \neq \emptyset$:

We have $-D(\alpha') \subset T_{L(\lambda)}$ and $-D(\alpha') \subset T_{{}^{s_\alpha}L(s_\alpha.\lambda)} = s_\alpha(T_{L(s_\alpha.\lambda)})$ by Lemma 8.4 and the fact that $(\alpha|\alpha') = 0$. So $-D(\alpha') \subset C(L(\lambda)) \cap C({}^{s_\alpha}L(s_\alpha.\lambda))$ thus $C(L(\lambda)) \cap C({}^{s_\alpha}L(s_\alpha.\lambda))$ generate Q by Lemma 8.5. This implies that $C(L(\lambda)) - C({}^{s_\alpha}L(s_\alpha.\lambda)) = Q$. The weights of $L(\lambda)$ and ${}^{s_\alpha}L(s_\alpha.\lambda)$ are contained in $q^Q\lambda$ so a weight in the essential support of $L(\lambda)$ (resp. ${}^{s_\alpha}L(s_\alpha.\lambda)$) is of the form $q^{\mu_1}\lambda$ (resp. $q^{\mu_2}\lambda$) for some $\mu_1, \mu_2 \in Q$. By the above $q^{C(L(\lambda))+\mu_1}\lambda \cap q^{C({}^{s_\alpha}L(s_\alpha.\lambda))+\mu_2}\lambda \neq \emptyset$. Since $q^{C(L(\lambda))+\mu_1}\lambda \subset \operatorname{Supp}_{\text{ess}}(L(\lambda))$ and $q^{C({}^{s_\alpha}L(s_\alpha.\lambda))+\mu_2}\lambda \subset \operatorname{Supp}_{\text{ess}}({}^{s_\alpha}L(s_\alpha.\lambda))$ we have proved the claim.

So we have proved that any two roots of $A(\lambda)$ are connected. Since there are no cycles in the Dynkin diagram of a simple Lie algebra we get $A(\lambda) = 2$.□

8.2 Rank 2 calculations

Following the procedure in [Mat00, Section 7] we classify admissible simple highest weight modules in rank 2 in order to classify the modules in higher ranks. We only consider non-integral weights because of Theorem 8.1. We assume that q is transcendental over $\mathbb{Q}$.

Lemma 8.7 *Assume $\mathfrak{g} = \mathfrak{sl}_3$. Let $\lambda \in X$ be a non-integral weight. The module $L(\lambda)$ is admissible if and only if $q^\rho\lambda(K_\beta) \in \pm q^{\mathbb{Z}_{>0}}$ for at least one root $\beta \in \Phi^+$.*

Proof. It is easy to show that the Verma module $M(\lambda)$ is not admissible. So $q^\rho\lambda(K_\beta) \in \pm q^{\mathbb{Z}_{>0}}$ for at least one root $\beta \in \Phi^+$ by Theorem 8.2. On the other hand suppose $q^\rho\lambda(K_\beta) \in \pm q^{\mathbb{Z}_{>0}}$ for at least one root $\beta \in \Phi^+$. If $q^\rho\lambda(K_\alpha) \in \pm q^{\mathbb{Z}_{>0}}$ for a simple root $\alpha \in \Pi$ then by easy calculations we see that $M(s_\alpha.\lambda)$ is a submodule of $M(\lambda)$. If $q^\rho\lambda(K_\alpha) \notin \pm q^{\mathbb{Z}_{>0}}$ for both simple roots $\alpha \in \Pi$ then we get that $M(s_\beta.\lambda)$ is a submodule by Theorem 8.2. So in both cases we have a submodule $M(s_\beta.\lambda)$ of $M(\lambda)$. Since $L(\lambda)$ is the unique simple quotient of $M(\lambda)$, $L(\lambda)$ is a subquotient of $M(\lambda)/M(s_\beta.\lambda)$. Since $M(\lambda)/M(s_\beta.\lambda)$ is admissible we see that $L(\lambda)$ is admissible as well. □

Lemma 8.8 *Assume $\mathfrak{g}$ is of type C_2 (i.e. $\mathfrak{g} = \mathfrak{sp}(4)$). Let $\Pi = \{\alpha_1, \alpha_2\}$ where α_1 is short and α_2 is long. Let $\lambda \in X$ be a non-integral weight. The module $L(\lambda)$ is infinite dimensional and admissible if and only if $q^\rho\lambda(K_{\alpha_1}), q^\rho\lambda(K_{\alpha_1+\alpha_2}) \in \pm q^{\mathbb{Z}_{>0}}$ and $\lambda(K_{\alpha_2}), \lambda(K_{2\alpha_1+\alpha_2}) \in \pm q^{1+2\mathbb{Z}} (= \pm q_{\alpha_2}^{1/2+\mathbb{Z}} = \pm q_{2\alpha_1+\alpha_2}^{1/2+\mathbb{Z}})$.*

Proof. Theorem 8.2 implies that $q^\rho\lambda(K_\beta) \in q_\beta^{\mathbb{Z}_{>0}}$ for at least two $\beta \in \Phi^+$ because otherwise $L(\lambda) = M(\lambda)/M(s_\beta.\lambda)$ for some $\beta \in \Phi^+$. But $M(\lambda)/M(s_\beta.\lambda)$ is not admissible. Since λ is not integral we know $q^\rho\lambda(K_\alpha) \notin q_\alpha^{\mathbb{Z}}$ for some $\alpha \in \Pi$. Suppose $\lambda(K_{\alpha_1}) \notin \pm q_{\alpha_1}^{\mathbb{Z}}$. We split into cases and arrive at a contradiction in both cases: If $\lambda(K_{\alpha_2}) \notin \pm q_{\alpha_2}^{\mathbb{Z}_{>0}}$ then by the above $q^\rho\lambda(K_{\alpha_1+\alpha_2}) \in \pm q_{\alpha_1+\alpha_2}^{\mathbb{Z}_{>0}} = \pm q^{\mathbb{Z}_{>0}}$ and $q^\rho\lambda(K_{2\alpha_1+\alpha_2}) \in \pm q_{2\alpha_1+\alpha_2}^{\mathbb{Z}_{>0}} = \pm q^{2\mathbb{Z}_{>0}}$ which implies that $q^\rho\lambda(K_{\alpha_1}) = q^\rho\lambda(K_{2\alpha_1+\alpha_2}K_{\alpha_1+\alpha_2}^{-1}) \in \pm q^{\mathbb{Z}} = \pm q_{\alpha_1}^{\mathbb{Z}}$. A contradiction.

The other case is $q^\rho\lambda(K_{\alpha_2}) \in \pm q_{\alpha_2}^{\mathbb{Z}_{>0}} = \pm q^{2\mathbb{Z}_{>0}}$: In this case we get $\lambda(K_{\alpha_1+\alpha_2}) \notin \pm q^{\mathbb{Z}} = \pm q_{\alpha_1+\alpha_2}^{\mathbb{Z}}$ so the last root, $2\alpha_1 + \alpha_2$, must satisfy that $q^\rho\lambda(K_{2\alpha_1+\alpha_2}) \in \pm q_{2\alpha_1+\alpha_2}^{\mathbb{Z}_{>0}} = \pm q^{2\mathbb{Z}_{>0}}$. But this implies that $\lambda(K_{\alpha_1})^2 = \lambda(K_{2\alpha_1+\alpha_2}K_{\alpha_2}^{-1}) \in \pm q^{2\mathbb{Z}}$ which implies that $\lambda(K_{\alpha_1}) \in \pm q^{\mathbb{Z}}$. A contradiction.

So $\lambda(K_{\alpha_1}) \in \pm q^{\mathbb{Z}}$. Since λ is not integral we get $\lambda(K_{\alpha_2}) \notin \pm q_{\alpha_2}^{\mathbb{Z}} = \pm q^{2\mathbb{Z}}$. This implies that $\lambda(K_{2\alpha_1+\alpha_2}) \notin \pm q^{2\mathbb{Z}} = \pm q_{2\alpha_1+\alpha_2}^{\mathbb{Z}}$. Since $q^\rho\lambda(K_\beta) \in \pm q_\beta^{\mathbb{Z}_{>0}}$ for at least two $\beta \in \Phi^+$ we get $q^\rho\lambda(K_{\alpha_1}) \in \pm q^{\mathbb{Z}_{>0}}$ and $q^\rho\lambda(K_{\alpha_1+\alpha_2}) \in \pm q^{\mathbb{Z}_{>0}}$. This in turn implies that $\lambda(K_{\alpha_2}) = \lambda(K_{\alpha_1+\alpha_2}K_{\alpha_1}^{-1}) \in \pm q^{\mathbb{Z}}$. Since $\lambda(K_{\alpha_2}) \notin \pm q^{2\mathbb{Z}}$ we get $\lambda(K_{\alpha_2}) \in \pm q^{1+2\mathbb{Z}}$. Similarly $\lambda(K_{2\alpha_1+\alpha_2}) = \lambda(K_{\alpha_1+\alpha_2}K_{\alpha_1}) \in \pm q^{1+2\mathbb{Z}}$. So we have shown the only if part.

Assume λ is as required in the lemma. We will show that $L(\lambda)$ is admissible. By Theorem 8.2 we see that the composition factors of $M(s_{\alpha_1}.\lambda)$ are $L(s_{\alpha_1}.\lambda)$ and $L(s_{\alpha_1+\alpha_2}s_{\alpha_1}.\lambda) = M(w_0.\lambda)$ and the composition factors of $M(s_{\alpha_1+\alpha_2})$ are $L(s_{\alpha_1+\alpha_2})$ and $L(s_{\alpha_1}s_{\alpha_1+\alpha_2}) = M(w_0.\lambda)$. So

$$\sum_{\substack{\beta\in\Phi^+ \\ q^\rho\lambda(K_\beta)\in\pm q_\beta^{\mathbb{Z}_{>0}}}} \operatorname{ch} M(s_\beta.\lambda) = \operatorname{ch} L(s_{\alpha_1}.\lambda) + \operatorname{ch} L(s_{\alpha_1+\alpha_2}.\lambda) + 2\operatorname{ch} L(w_0.\lambda).$$

So the composition factors of the maximal submodule of $M(\lambda)$ are $L(s_{\alpha_1}.\lambda)$, $L(s_{\alpha_1+\alpha_2}.\lambda)$ and $L(w_0.\lambda)$. The worst case scenario being multiplicity one. In this case the character of $L(\lambda)$ is

$$\begin{aligned}\operatorname{ch} M(\lambda) - \operatorname{ch} L(s_{\alpha_1}.\lambda) - \operatorname{ch} L(s_{\alpha_1+\alpha_2}) - \operatorname{ch} L(w_0.\lambda) &= \\ = \operatorname{ch} M(\lambda) - \operatorname{ch} M(s_{\alpha_1}.\lambda) - \operatorname{ch} M(s_{\alpha_1+\alpha_2}.\lambda) + \operatorname{ch} M(w_0.\lambda)\end{aligned}$$

The character of Verma modules are known and by an easy calculation it is seen that this would imply $L(\lambda)$ is admissible (cf. the proof of Lemma 7.2 in [Mat00]). ☐

8.3 Type A, D, E

In this section we complete the classification of all simple admissible highest weight modules when the Dynkin diagram of $\mathfrak{g}$ is simply laced. In particular we show that $\mathfrak{g}$ does not admit infinite dimensional simple admissible modules when $\mathfrak{g}$ is of type D and E. In Section 8.5 we show that the same is the case when $\mathfrak{g}$ is of type B or F. Combining this and Section 8.5 we get that $\mathfrak{g}$ admits infinite dimensional simple admissible modules if and only if $\mathfrak{g}$ is of type A or C. Remember that we restrict our attention to transcendental q and to non-integral weights because of Theorem 8.1.

Definition 8.9 *Let $\lambda : U_q^0 \to \mathbb{C}$ be a weight. In the Dynkin diagram of $\mathfrak{g}$ let any node corresponding to $\alpha \in \Pi \cap A(\lambda)$ be written as $\circ$ and every other as $\bullet$.*

e.g. if $\mathfrak{g} = \mathfrak{sl}_3$ *and* $|A(\lambda)| = 1$ *then the graph corresponding to* λ *would look like this:*

$$\bullet \text{ ——— } \circ$$

We call this the colored Dynkin diagram corresponding to λ.

In this way we get a 'coloring' of the Dynkin diagram for every λ.

Lemma 8.10 *Let* $\lambda \in X$ *be a non-integral weight such that* $L(\lambda)$ *is admissible. If the colored Dynkin diagram of* λ *contains*

$$\overset{\alpha'}{\circ} \text{ ——— } \overset{\alpha}{\circ}$$

as a subdiagram then $q^{\rho}\lambda(K_{\alpha+\alpha'}) \in \pm q_{\alpha'+\alpha}^{\mathbb{Z}_{>0}}$.

Proof. Let v_λ be a highest weight vector of $L(\lambda)$. Let $\mathfrak{s}$ be the Lie algebra $\mathfrak{sl}_3$ with α and α' as simple roots. Let U be the subalgebra of U_q generated by $F_\alpha, F_{\alpha'}, K_\alpha^{\pm 1}, K_{\alpha'}^{\pm 1}, E_\alpha, E_{\alpha'}$. Then $U \cong U_{q_\alpha}(\mathfrak{s})$ as algebras and Uv_λ contains the simple highest weight $U_{q_\alpha}(\mathfrak{s})$-module $L(\lambda, \mathfrak{s})$ of highest weight λ (restricted to $U_{q_\alpha}^0(\mathfrak{s})$) as a subquotient. Since $L(\lambda)$ is admissible so is Uv_λ hence $L(\lambda, \mathfrak{s})$ is admissible. Then Lemma 8.7 implies that $q^{\rho}\lambda(K_{\alpha+\alpha'}) \in \pm q_\alpha^{\mathbb{Z}_{>0}}$. $\square$

Lemma 8.11 *Let* $\lambda \in X$ *be a non-integral weight such that* $L(\lambda)$ *is admissible. If the colored Dynkin diagram of* λ *contains*

$$\overset{\alpha'}{\circ} \text{ ——— } \overset{\alpha}{\circ} \text{ ——— } \overset{\alpha''}{\bullet}$$

as a subdiagram then $L(s_\alpha.\lambda)$ *is admissible and the colored Dynkin diagram corresponding to* $s_\alpha.\lambda$ *contains*

$$\overset{\alpha'}{\bullet} \text{ ——— } \overset{\alpha}{\circ} \text{ ——— } \overset{\alpha''}{\circ}$$

i.e. we can 'move' $\circ$ ——— $\circ$ *and still get an admissible module.*

Proof. $L(s_\alpha.\lambda)$ is admissible by Proposition 5.10. It is easy to see that $q^{\rho}s_\alpha.\lambda(K_\alpha) \notin \pm q^{\mathbb{Z}}$ (follows by Lemma 8.10 since λ is non-integral), that $q^{\rho}s_\alpha.\lambda(K_{\alpha''}) \notin \pm q^{\mathbb{Z}}$ and that $q^{\rho}s_\alpha.\lambda(K_{\alpha'}) \in \pm q^{\mathbb{Z}_{>0}}$ (by Lemma 8.10) $\square$

Lemma 8.12 *Assume* $\mathfrak{g} \neq \mathfrak{sl}_2$. *Let* $\lambda \in X$ *be a non-integral weight such that* $L(\lambda)$ *is admissible.*

If $A(\lambda) = \{\alpha\}$ *then* α *is only connected to one other simple root* α', $L(s_\alpha.\lambda)$ *is admissible and the corresponding colored Dynkin diagram of* $s_\alpha.\lambda$ *contains*

$$\overset{\alpha}{\circ} \text{ ——— } \overset{\alpha'}{\circ}$$

as a subdiagram.

On the other hand if the colored Dynkin diagram of λ *contains*

$$\overset{\alpha}{\circ} \text{ ——— } \overset{\alpha'}{\circ}$$

and α' is the only root connected to α then the colored Dynkin diagram of $s_\alpha.\lambda$ contains



as a subdiagram.

Proof. Since $\alpha \in A(\lambda)$, $L(s_\alpha.\lambda)$ is admissible by Proposition 5.10. First assume $A(\lambda) = \{\alpha\}$. If α is connected to two distinct roots α' and α'' then it is easily seen that $\alpha', \alpha'' \in A(s_\alpha.\lambda)$ contradicting the fact that $A(s_\alpha.\lambda)$ is connected (Lemma 8.6). It is easily seen that $q^\rho s_\alpha.\lambda(K_\alpha) \notin \pm q^{\mathbb{Z}_{>0}}$ (since λ is non integral) and $q^\rho s_\alpha.\lambda(K_{\alpha'}) \notin q^{\mathbb{Z}_{>0}}$.

On the other hand if $A(\lambda) = \{\alpha, \alpha'\}$ then $q^\rho s_\alpha.\lambda(K_{\alpha'}) = q^\rho \lambda(K_{\alpha+\alpha'}) \in \pm q^{\mathbb{Z}_{>0}}$ by Lemma 8.10. $\square$

Now we can eliminate the types that are not type A by the following theorem:

Theorem 8.13 *Assume $\mathfrak{g}$ is a simple Lie algebra of simply laced type. If there exists an infinite dimensional admissible simple module then $\mathfrak{g}$ is of type A.*

Proof. Suppose there exists an infinite dimensional admissible simple module then by Theorem 5.12 there exists a $\lambda \in X$ such that $L(\lambda)$ is an infinite admissible simple highest weight module. By Theorem 8.1 and the classification in [Mat00] there exists no highest weight simple admissible modules with integral weights unless $\mathfrak{g}$ is of type A. We need to show the same for non-integral weights.

If the Dynkin diagram is simply laced and not of type A then the Dynkin diagram contains



as a subdiagram.

By Lemma 8.12 we can assume without loss of generality that $|A(\lambda)| = 2$ and by Lemma 8.11 we can assume that the colored Dynkin diagram corresponding to λ contains the following:



But then $L(s_\gamma.\lambda)$ is admissible as well by Proposition 5.10 and the colored Dynkin diagram for $s_\gamma.\lambda$ contains



contradicting the fact that $A(\lambda)$ is connected. $\square$

Combining all the above results we get

Theorem 8.14 *Let* $\mathfrak{g} = \mathfrak{sl}_{n+1}$*,* $n \geq 2$ *with simple roots* $\alpha_1, \dots, \alpha_n$ *such that* $(\alpha_i|\alpha_{i+1}) = -1$*,* $i = 1, \dots, n$*. Let* $\lambda \in X$ *be a non-integral weight.*

$L(\lambda)$ is admissible if and only if the colored Dynkin diagram of λ is of one of the following types:



Proof. By the above results these are the only possibilites. To show that $L(\lambda)$ is admissible when the colored Dynkin diagram is of the above form use the fact that by Lemma 8.11 and Lemma 8.12 we can assume λ has colored Dynkin diagram as follows:



Let $\beta_i = \alpha_1 + \alpha_2 + \cdots + \alpha_i$, $i = 1, \dots, n$. We see easily that $T_{L(\lambda)} = -\{\beta_1, \beta_2, \dots, \beta_n\}$ and $F_L = \Phi^+ \cup \Phi_{\{\alpha_2, \dots, \alpha_n\}}$. Let $\mathfrak{l}$, $\mathfrak{u}$, $\mathfrak{p}$ etc. be defined as in Section 2 of [Ped15a]. By [Ped15a, Theorem 2.23] $N := L(\lambda)^{\mathfrak{u}}$ is a simple finite dimensional $U_q(\mathfrak{l})$-module and $L(\lambda)$ is the unique simple quotient of $\mathcal{M}(N) = U_q \otimes_{U_q(\mathfrak{p})} N$. Since the vectors $\beta_1, \dots, \beta_n$ are linearly independent $\mathcal{M}(N)$ is admissible. This implies that $L(\lambda)$ is admissible since it is a quotient of $\mathcal{M}(N)$. $\square$

We can now make Corollary 5.13 more specific in type A:

Corollary 8.15 *Let* $\mathfrak{g} = \mathfrak{sl}_{n+1}$*,* $n \geq 2$ *with simple roots* $\alpha_1, \dots, \alpha_n$ *such that* $(\alpha_i|\alpha_{i+1}) = -1$*,* $i = 1, \dots, n$*. Let* $\beta_j = \alpha_1 + \cdots + \alpha_j$*,* $j = 1, \dots, n$ *and* $\Sigma = \{\beta_1, \dots, \beta_n\}$*. Let* $F_{\beta_j} = T_{s_1} \cdots T_{s_{j-1}}(F_{\alpha_j})$ *and let* $F_\Sigma = \{q^a F_{\beta_1}^{a_1} \cdots F_{\beta_n}^{a_n} | a_i \in \mathbb{N}, a \in \mathbb{Z}\}$ *be the corresponding Ore subset. Then* Σ *is a set of commuting roots that is a basis of* Q *with corresponding Ore subset* F_Σ*.*

Let $\beta'_j = \alpha_n + \cdots + \alpha_{n-j}$*,* $j = 1, \dots, n$ *and* $\Sigma = \{\beta'_1, \dots, \beta'_n\}$*. Let* $F'_{\beta'_j} = T_{s_n} \cdots T_{s_{n-j+1}}(F_{\alpha_{n-j}})$ *and let* $F_{\Sigma'} = \{q^a (F'_{\beta'_1})^{a_1} \cdots (F'_{\beta'_n})^{a_n} | a_i \in \mathbb{N}, a \in \mathbb{Z}\}$ *be the corresponding Ore subset. Then* Σ' *is a set of commuting roots that is a basis of* Q *with corresponding Ore subset* $F_{\Sigma'}$*.*

Let L *be a simple torsion free module then one of the two following claims hold*

- *There exists a* $\lambda \in X$ *with* $\lambda(K_{\alpha_1}) \notin \pm q^{\mathbb{N}}$*,* $\lambda(K_{\alpha_i}) \in \pm q^{\mathbb{N}}$*,* $i = 2, \dots, n$ *and* $\mathbf{b} \in (\mathbb{C}^*)^n$ *such that*
$$L \cong \varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}.$$
- *There exists a* $\lambda \in X$ *with* $\lambda(K_{\alpha_n}) \notin \pm q^{\mathbb{N}}$*,* $\lambda(K_{\alpha_i}) \in \pm q^{\mathbb{N}}$*,* $i = 1, \dots, n-1$ *and* $\mathbf{b} \in (\mathbb{C}^*)^n$ *such that*
$$L \cong \varphi_{F_{\Sigma'}, \mathbf{b}}.L(\lambda)_{F_{\Sigma'}}.$$

Proof. By Theorem 5.12 $\mathcal{EXT}(L) \cong \mathcal{EXT}(L(\lambda'))$ for some $\lambda' \in X$. If λ' is non-integral then by Theorem 8.14, Lemma 8.11, Lemma 5.11 and Proposition 5.6 there exists a λ such that $\lambda(K_{\alpha_1}) \notin \pm q^{\mathbb{N}}$, $\lambda(K_{\alpha_i}) \in \pm q^{\mathbb{N}}$, $i = 2, \ldots, n$ and such that $\mathcal{EXT}(L(\lambda')) \cong \mathcal{EXT}(L(\lambda))$. By Lemma 5.4 we can choose Σ as the commuting set of roots that is used in the definition of $\mathcal{EXT}(L(\lambda))$.

If λ' is integral we see by Theorem 8.1, Lemma 5.11, Proposition 5.6 and the classification in [Mat00, Section 8] that $\mathcal{EXT}(L(\lambda')) \cong \mathcal{EXT}(L(\lambda))$ for a λ such that $A(\lambda) = \{\alpha_1\}$ or $A(\lambda) = \{\alpha_n\}$ (cf. e.g. [Mat00, Proposition 8.5]).

Now the result follows just like in the proof of Corollary 5.13. $\square$

In Section 9 we determine all $\mathbf{b} \in (\mathbb{C}^*)^n$ such that $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ is torsion free with Σ as above in Corollary 8.15 and λ such that $\lambda(K_{\alpha_1}) \notin \pm q^{\mathbb{N}}$, $\lambda(K_{\alpha_i}) \in \pm q^{\mathbb{N}}$, $i = 2, \ldots, n$. By symmetry of the Dynkin diagram and Corollary 8.15 this classifies all simple torsion free modules.

8.4 Quantum Shale-Weil representation

In this section we assume $\mathfrak{g}$ is of type C_n. Let $\alpha_1, \ldots, \alpha_n$ be the simple roots such that α_i is connected to α_{i+1} and α_1 is long. We will describe a specific admissible module V and show that $V = L(\omega^+) \oplus L(\omega^-)$ for some weights $\omega^\pm$ with the purpose of classifying the admissible simple highest weight modules, see Theorem 8.17. Let $V = \mathbb{C}[X_1, \ldots, X_n]$. We describe an action of the simple root vectors on V: For $i \in \{2, \ldots, n\}$

$$\begin{aligned}
E_{\alpha_1} X_1^{a_1} X_2^{a_2} \cdots X_n^{a_n} =& - \frac{[a_1][a_1-1]}{[2]} X_1^{a_1-2} X_2^{a_2} \cdots X_n^{a_n} \\
F_{\alpha_1} X_1^{a_1} X_2^{a_2} \cdots X_n^{a_n} =& \frac{1}{[2]} X_1^{a_1+2} X_2^{a_2} \cdots X_n^{a_n} \\
E_{\alpha_i} X_1^{a_1} \cdots X_n^{a_n} =& [a_i] X_1^{a_1} \cdots X_{i-1}^{a_{i-1}+1} X_i^{a_i-1} \cdots X_n^{a_n} \\
F_{\alpha_i} X_1^{a_1} \cdots X_n^{a_n} =& [a_{i-1}] X_1^{a_1} \cdots X_{i-1}^{a_{i-1}-1} X_i^{a_i+1} \cdots X_n^{a_n} \\
K_{\alpha_1}^{\pm 1} X_1^{a_1} X_2^{a_2} \cdots X_n^{a_n} =& q^{\mp(2a_1+1)} X_1^{a_1} X_2^{a_2} \cdots X_n^{a_n} \\
K_{\alpha_i}^{\pm 1} X_1^{a_1} X_2^{a_2} \cdots X_n^{a_n} =& q^{\pm(a_{i-1}-a_i)} X_1^{a_1} X_2^{a_2} \cdots X_n^{a_n}.
\end{aligned}$$

We check that this is an action of U_q by checking the generating relations. These are tedious and kind of long calculations but just direct calculations. We refer to the generating relations as (R1) to (R6) like in [Jan96, Section 4.3].

(R1) is clear. (R2) and (R3): Let $j \in \{1, \ldots, n\}$

$$\begin{aligned}
K_{\alpha_j} E_{\alpha_1} X_1^{a_1} \cdots X_n^{a_n} =& \begin{cases} -q^{-2a_1+3} \frac{[a_1][a_1-1]}{[2]} X_1^{a_1-2} X_2^{a_2} \cdots X_n^{a_n} & \text{if } j = 1 \\ -q^{a_1-2-a_2} \frac{[a_1][a_1-1]}{[2]} X_1^{a_1-2} X_2^{a_2} \cdots X_n^{a_n} & \text{if } j = 2 \\ -q^{a_{j-1}-a_j} \frac{[a_1][a_1-1]}{[2]} X_1^{a_1-2} X_2^{a_2} \cdots X_n^{a_n} & \text{if } j > 2 \end{cases} \\
=& q^{(\alpha_1|\alpha_j)} E_{\alpha_1} K_{\alpha_j} X_1^{a_1} X_2^{a_2} \cdots X_n^{a_n}.
\end{aligned}$$

Similar for $K_{\alpha_j}F_{\alpha_1}$. For $i \in \{2,\dots,n\}$

$$\begin{aligned}
K_{\alpha_j}E_{\alpha_i}X_1^{a_1}\cdots X_n^{a_n} =& \begin{cases} q^{a_{j-1}-a_j}[a_i]X_1^{a_1}\cdots X_{i-1}^{a_{i-1}+1}X_i^{a_i-1}\cdots X_n^{a_n} & \text{if } |j-i|>1 \\ q^{a_{j-1}-a_j-1}[a_i]X_1^{a_1}\cdots X_{i-1}^{a_{i-1}+1}X_i^{a_i-1}\cdots X_n^{a_n} & \text{if } j=i-1 \\ q^{a_{j-1}+1-a_j+1}[a_i]X_1^{a_1}\cdots X_{i-1}^{a_{i-1}+1}X_i^{a_i-1}\cdots X_n^{a_n} & \text{if } j=i \\ q^{a_{j-1}-1-a_j}[a_i]X_1^{a_1}\cdots X_{i-1}^{a_{i-1}+1}X_i^{a_i-1}\cdots X_n^{a_n} & \text{if } j=i+1 \end{cases} \\
=& q^{(\alpha_i|\alpha_j)}E_{\alpha_1}K_{\alpha_j}X_1^{a_1}X_2^{a_2}\cdots X_n^{a_n}.
\end{aligned}$$

Similarly for $K_{\alpha_j}F_{\alpha_i}$.

(R4):

$$\begin{aligned}
[E_{\alpha_1},F_{\alpha_1}]X_1^{a_1}X_2^{a_2}\cdots X_n^{a_n} =& E_{\alpha_1}\frac{1}{[2]}X_1^{a_1+2}X_2^{a_2}\cdots X_n^{a_n} + F_{\alpha_1}\frac{[a_1][a_1-1]}{[2]}X_1^{a_1-2}X_2^{a_2}\cdots X_n^{a_n} \\
=& \left(-\frac{[a_1+2][a_1+1]}{[2][2]} + \frac{[a_1][a_1-1]}{[2][2]}\right)X_1^{a_1}\cdots X_n^{a_n} \\
=& \frac{q^{-2a_1-1}-q^{2a_1+1}}{q^2-q^{-2}}X_1^{a_1}\cdots X_n^{a_n} \\
=& \frac{K_{\alpha_1}-K_{\alpha_1}^{-1}}{q^2-q^{-2}}X_1^{a_1}\cdots X_n^{a_n}.
\end{aligned}$$

$$\begin{aligned}
[E_{\alpha_1},F_{\alpha_2}]X_1^{a_1}\cdots X_n^{a_n} =& [a_1]E_{\alpha_1}X_1^{a_1-1}X_2^{a_2+1}\cdots X_n^{a_n} + \frac{[a_1][a_1-1]}{[2]}F_{\alpha_2}X_1^{a_1-2}X_2^{a_2}\cdots X_n^{a_n} \\
=& -\frac{[a_1][a_1-1][a_1-2]}{[2]}X_1^{a_1-3}X_2^{a_2+1}\cdots X_n^{a_n} \\
& + \frac{[a_1][a_1-1][a_1-2]}{[2]}X_1^{a_1-3}X_2^{a_2+1}\cdots X_n^{a_n} \\
=& 0.
\end{aligned}$$

For $i>2$ clearly $[E_{\alpha_1},F_{\alpha_i}]X_1^{a_1}\cdots X_n^{a_n}=0$. For $i,j\in\{2,\dots,n\}$: If $|i-j|>1$ clearly $[E_{\alpha_i},F_{\alpha_j}]X_1^{a_1}\cdots X_n^{a_n}=0$.

$$\begin{aligned}
[E_{\alpha_i},F_{\alpha_{i+1}}]X_1^{a_n}\cdots X_n^{a_n} =& [a_i]E_{\alpha_i}X_1^{a_1}\cdots X_i^{a_i-1}X_{i+1}^{a_{i+1}+1}\cdots X_n^{a_n} \\
& - [a_i]F_{\alpha_{i+1}}X_1^{a_1}\cdots X_{i-1}^{a_{i-1}+1}X_i^{a_i-1}\cdots X_n^{a_n} \\
=& [a_i][a_i-1]X_1^{a_1}\cdots X_{i-1}^{a_{i-1}+1}X_i^{a_i-2}X_{i+1}^{a_{i+1}+1}\cdots X_n^{a_n} \\
& - [a_i][a_i-1]X_1^{a_1}\cdots X_{i-1}^{a_{i-1}+1}X_i^{a_i-2}X_{i+1}^{a_{i+1}+1}\cdots X_n^{a_n} \\
=& 0.
\end{aligned}$$

$$\begin{aligned}
[E_{\alpha_2},F_{\alpha_1}]X_1^{a_1}\cdots X_n^{a_n} =& E_{\alpha_2}\frac{1}{[2]}X_1^{a_1+2}X_2^{a_2}\cdots X_n^{a_n} - [a_2]F_{\alpha_1}X_1^{a_1+1}X_2^{a_2-1}\cdots X_n^{a_n} \\
=& \frac{[a_2]}{[2]}X_1^{a_1+3}X_2^{a_2-1}\cdots X_n^{a_n} - \frac{[a_2]}{[2]}X_1^{a_1+3}X_2^{a_2-1}\cdots X_n^{a_n} \\
=& 0.
\end{aligned}$$

For $i > 2$:

$$\begin{aligned}
[E_{\alpha_i}, F_{\alpha_{i-1}}]X_1^{a_n} \cdots X_n^{a_n} =& [a_{i-2}]E_{\alpha_i}X_1^{a_1} \cdots X_{i-2}^{a_{i-2}-1}X_{i-1}^{a_{i-1}+1} \cdots X_n^{a_n} \\
&- [a_i]F_{\alpha_{i-1}}X_1^{a_1} \cdots X_{i-1}^{a_{i-1}+1}X_i^{a_i-1} \cdots X_n^{a_n} \\
=& [a_{i-2}][a_i]X_1^{a_1} \cdots X_{i-2}^{a_{i-2}-1}X_{i-1}^{a_{i-1}+2}X_i^{a_i-1} \cdots X_n^{a_n} \\
&- [a_i][a_{i-2}]F_{\alpha_{i-1}}X_1^{a_1} \cdots X_{i-2}^{a_{i-2}-1}X_{i-1}^{a_{i-1}+2}X_i^{a_i-1} \cdots X_n^{a_n} \\
=& 0.
\end{aligned}$$

For $i > 1$:

$$\begin{aligned}
[E_{\alpha_i}, F_{\alpha_i}]X_1^{a_1} \cdots X_n^{a_n} =& [a_{i-1}]E_{\alpha_i}X_1^{a_1} \cdots X_{i-1}^{a_{i-1}+1}X_i^{a_i-1} \cdots X_n^{a_n} \\
&- [a_i]F_{\alpha_i}X_1^{a_1} \cdots X_{i-1}^{a_{i-1}-1}X_i^{a_i+1} \cdots X_n^{a_n} \\
=& ([a_{i-1}][a_i - 1] - [a_i][a_{i-1} - 1])X_1^{a_1} \cdots X_{i-1}^{a_{i-1}}X_i^{a_i} \cdots X_n^{a_n} \\
=& [a_{i-1} - a_i]X_1^{a_1} \cdots X_n^{a_n} \\
=& \frac{K_{\alpha_i} - K_{\alpha_i}^{-1}}{q - q^{-1}}X_1^{a_1} \cdots X_n^{a_n}.
\end{aligned}$$

Finally we have the relations (R5) and (R6): Clearly $[E_{\alpha_i}, E_{\alpha_j}]X_1^{a_1} \cdots X_n^{a_n} = 0$ and $[F_{\alpha_i}, F_{\alpha_j}]X_1^{a_1} \cdots X_n^{a_n} = 0$ when $|j - i| > 1$.

$$\begin{aligned}
&(E_{\alpha_2}^3 E_{\alpha_1} - [3]E_{\alpha_2}^2 E_{\alpha_1} E_{\alpha_2} + [3]E_{\alpha_2} E_{\alpha_1} E_{\alpha_2}^2 - E_{\alpha_1} E_{\alpha_2}^3)X_1^{a_1} \cdots X_n^{a_n} \\
&= \frac{1}{[2]}\Big(- [a_1][a_1 - 1][a_2][a_2 - 1][a_2 - 2] \\
&\quad + [3][a_1 + 1][a_1][a_2][a_2 - 1][a_2 - 2] \\
&\quad - [3][a_1 + 2][a_1 + 1][a_2][a_2 - 1][a_2 - 2] \\
&\quad + [a_1 + 3][a_1 + 2][a_2][a_2 - 1][a_2 - 2]\Big)X_1^{a_1+1}X_2^{a_2-3} \cdots X_n^{a_n} \\
&= \frac{[a_2][a_2 - 1][a_2 - 2]}{[2]}\Big(- [a_1][a_1 - 1] + [3][a_1 + 1][a_1] \\
&\quad - [3][a_1 + 2][a_1 + 1] + [a_1 + 3][a_1 + 2]\Big)X_1^{a_1+1}X_2^{a_2-3} \cdots X_n^{a_n} \\
&= 0.
\end{aligned}$$

$$\begin{aligned}
(E_{\alpha_1}^2 E_{\alpha_2} - [2]_{\alpha_1} E_{\alpha_1} E_{\alpha_2} E_{\alpha_1} + E_{\alpha_2} E_{\alpha_1}^2)X_1^{a_1} \cdots X_n^{a_n} \\
=& \frac{[a_2]}{[2][2]}\big([a_1 + 1][a_1][a_1 - 1][a_1 - 2] \\
&- [2]_{\alpha_1}[a_1][a_1 - 1][a_1 - 1][a_1 - 2] \\
&+ [a_1][a_1 - 1][a_1 - 2][a_1 - 3]\big)X_1^{a_1+3}X_2^{a_2-1} \cdots X_n^{a_n} \\
=& \frac{[a_2][a_1][a_1 - 1][a_1 - 2]}{[2][2]}\Big([a_1 + 1] - [2]_{\alpha_1}[a_1 - 1] \\
&+ [a_1 - 3]\Big)X_1^{a_1+3}X_2^{a_2-1} \cdots X_n^{a_n} \\
=& 0.
\end{aligned}$$

For $i > 1$:

$$\begin{aligned}
(E_{\alpha_i}^2 E_{\alpha_{i+1}} - &[2]E_{\alpha_i}E_{\alpha_{i+1}}E_{\alpha_i} + E_{\alpha_{i+1}}E_{\alpha_i}^2)X_1^{a_1}\cdots X_n^{a_n}\\
&=[a_{i+1}][a_i]([a_i+1]-[2][a_i]+[a_i-1])X_1^{a_1}\cdots X_{i-1}^{a_{i-1}+2}X_i^{a_i-1}X_{i+1}^{a_{i+1}-1}\cdots X_n^{a_n}\\
&=0.
\end{aligned}$$

$$\begin{aligned}
(E_{\alpha_{i+1}}^2 E_{\alpha_i} - &[2]E_{\alpha_{i+1}}E_{\alpha_i}E_{\alpha_{i+1}} + E_{\alpha_i}E_{\alpha_{i+1}}^2)X_1^{a_1}\cdots X_n^{a_n}\\
&=[a_{i+1}][a_{i+1}-1]([a_i]-[2][a_i+1]+[a_i+2])X_1^{a_1}\cdots X_{i-1}^{a_{i-1}+1}X_i^{a_i+1}X_{i+1}^{a_{i+1}-2}\cdots X_n^{a_n}\\
&=0.
\end{aligned}$$

$$\begin{aligned}
(F_{\alpha_1}^2F_{\alpha_2} - [2]_{\alpha_1}F_{\alpha_1}F_{\alpha_2}F_{\alpha_1} + F_{\alpha_2}F_{\alpha_1}^2)X_1^{a_1}\cdots X_n^{a_n}&\\
=&\frac{1}{[2][2]}([a_1]-[2]_{\alpha_1}[a_1+2]+[a+4])X_1^{a_1+3}X_2^{a_2+1}\cdots X_n^{a_n}\\
=&0.
\end{aligned}$$

$$\begin{aligned}
(F_{\alpha_2}^3F_{\alpha_1} - [3]F_{\alpha_2}^2F_{\alpha_1}F_{\alpha_2} + [3]F_{\alpha_2}F_{\alpha_1}F_{\alpha_2}^2 - F_{\alpha_1}F_{\alpha_2}^3)X_1^{a_1}\cdots X_n^{a_n}&\\
=&\frac{1}{[2]}\Big([a_1+2][a_1+1][a_1]-[3][a_1][a_1+1][a_1]\\
&+[3][a_1][a_1-1][a_1]\\
&-[a_1][a_1-1][a_1-2]\Big)X_1^{a_1-1}X_2^{a_2+3}\cdots X_n^{a_n}\\
=&\frac{[a_1]}{[2]}\Big([a_1+2][a_1+1]-[3][a_1+1][a_1]\\
&+[3][a_1][a_1-1]\\
&-[a_1-1][a_1-2]\Big)X_1^{a_1-1}X_2^{a_2+3}\cdots X_n^{a_n}\\
=&0.
\end{aligned}$$

For $i > 1$:

$$\begin{aligned}
(F_{\alpha_i}^2 F_{\alpha_{i+1}} - &[2]F_{\alpha_i}F_{\alpha_{i+1}}F_{\alpha_i} + F_{\alpha_{i+1}}F_{\alpha_i}^2)X_1^{a_1}\cdots X_n^{a_n}\\
&=[a_{i-1}][a_{i-1}-1]([a_i]-[2][a_i+1]+[a_i+2])X_1^{a_1}\cdots X_{i-1}^{a_{i-1}-2}X_i^{a_i+1}X_{i+1}^{a_{i+1}+1}\cdots X_n^{a_n}\\
&=0.
\end{aligned}$$

$$\begin{aligned}
(F_{\alpha_{i+1}}^2 F_{\alpha_i} - &[2]F_{\alpha_{i+1}}F_{\alpha_i}F_{\alpha_{i+1}} + F_{\alpha_i}F_{\alpha_{i+1}}^2)X_1^{a_1}\cdots X_n^{a_n}\\
&=[a_{i-1}][a_i]([a_i+1]-[2][a_i]+[a_i-1])X_1^{a_1}\cdots X_{i-1}^{a_{i-1}-1}X_i^{a_i-1}X_{i+1}^{a_{i+1}+2}\cdots X_n^{a_n}\\
&=0.
\end{aligned}$$

So we have shown that V is a $U_q(\mathfrak{g})$-module. Note that V is admissible of degree 1 and $V = V^{even} \oplus V^{odd}$ where V^{even} are even degree polynomials and

V^{odd} are odd degree polynomials. Furthermore we see that $V^{even} = L(\omega^+)$ and $V^{odd} = L(\omega^-)$ where $\omega^\pm$ are the weights defined by $\omega^+(K_{\alpha_1}) = q^{-1}$, $\omega^+(K_{\alpha_i}) = 1$, $i > 1$ and $\omega^-(K_{\alpha_1}) = q^{-3}$, $\omega^-(K_{\alpha_2}) = q^{-1}$, $\omega^-(K_{\alpha_i}) = 1$, $i > 2$. V^{even} is generated by 1 and V^{odd} is generated by X_1. We will use the fact that $L(\omega^+)$ is admissible in Theorem 8.17 in the next section.

8.5 Type B, C, F

In this section we classify the simple highest weight admissible modules when $\mathfrak{g}$ is of type B, C or F. Remember that we have assumed that q is transcendental.

Theorem 8.16 *Let $\mathfrak{g}$ be a simple Lie algebra not of type G_2. Suppose there exists an infinite dimensional admissible simple $U_q(\mathfrak{g})$-module. Then $\mathfrak{g}$ is of type A or C.*

Proof. If $\mathfrak{g}$ is simply laced then Theorem 8.14 gives that $\mathfrak{g}$ is of type A. So assume $\mathfrak{g}$ is not of simply laced type. Theorem 8.1 and the classification in the classical case tells us that no admissible infinite dimensional simple highest weight modules exists with integral weights when $\mathfrak{g}$ is not simply laced (cf. [Mat00, Lemma 9.1]).

We have assumed that $\mathfrak{g}$ is not of type G_2 so the remaining non-simply laced types are B, C or F. We will show that the Dynkin diagram of $\mathfrak{g}$ can't contain the subdiagram



Assume the Dynkin diagram contains the above as a subdiagram. If there exists a simple admissible infinite dimensional module L then there exists a non-integral $\lambda \in X$ such that $L(\lambda)$ is infinite dimensional and admissible (Theorem 5.12). Let $\lambda \in X$ be a non-integral weight such that $L(\lambda)$ is admissible. Then by Lemma 8.8, $q^\rho\lambda(K_{\alpha_1}) \in \pm q_{\alpha_1}^{\mathbb{Z}} = \pm q^{\mathbb{Z}}$. By Lemma 8.11 and Lemma 8.12 we can assume without loss of generality that the colored Dynkin diagram of λ is of the form



Let $\mathfrak{s}$ be the simple rank 3 Lie algebra of type B_3. Let U be the subalgebra of U_q generated by $E_{\alpha_i}, F_{\alpha_i}, K_{\alpha_i}^{\pm 1}$, $i = 1, 2, 3$. Then $U \cong U_q(\mathfrak{s})$. Let $Q_{\mathfrak{s}} := \mathbb{Z}\{\alpha_1, \alpha_2, \alpha_3\} \subset Q$. Let v_λ be a highest weight vector of $L(\lambda)$. Then Uv_λ contains the simple highest weight $U_q(\mathfrak{s})$-module $L(\lambda, \mathfrak{s})$ of highest weight λ (restricted to $U_q^0(\mathfrak{s})$) as a subquotient. Since $L(\lambda)$ is admissible so is $L(\lambda, \mathfrak{s})$.

Like in the proof of Lemma 8.6 we get a contradiction if we can show that $T_{L(\lambda,\mathfrak{s})} \cap T_{{}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s})}$ generates $Q_{\mathfrak{s}}$. It is easily seen that $\{-\alpha_1-\alpha_2, -\alpha_3, -2\alpha_1-\alpha_2\} \subset T_{L(\lambda,\mathfrak{s})} \cap T_{{}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s})}$, so $T_{L(\lambda,\mathfrak{s})} \cap T_{{}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s})}$ generates $Q_{\mathfrak{s}}$. So $C(L(\lambda,\mathfrak{s})) \cap C({}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s}))$ generates $Q_{\mathfrak{s}}$. Therefore $C(L(\lambda,\mathfrak{s})) - C({}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s})) = Q_{\mathfrak{s}}$. The weights of $L(\lambda,\mathfrak{s})$ and ${}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s})$ are contained in $q^{Q_{\mathfrak{s}}}\lambda$ so a weight in the essential support of $L(\lambda,\mathfrak{s})$ (resp. ${}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s})$) is of the form $q^{\mu_1}\lambda$ (resp. $q^{\mu_2}\lambda$) for some $\mu_1, \mu_2 \in Q_{\mathfrak{s}}$. By the above $q^{C(L(\lambda,\mathfrak{s}))+\mu_1}\lambda \cap q^{C({}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s}))+\mu_2}\lambda \neq \emptyset$. Since $q^{C(L(\lambda,\mathfrak{s}))+\mu_1}\lambda \subset \operatorname{Supp}_{\mathrm{ess}}(L(\lambda))$ and $q^{C({}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s}))+\mu_2}\lambda \subset \operatorname{Supp}_{\mathrm{ess}}({}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s}))$ we have proved that $\operatorname{Supp}_{\mathrm{ess}}({}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s})) \cap \operatorname{Supp}_{\mathrm{ess}}(L(\lambda,\mathfrak{s})) \neq \emptyset$. By Proposition 5.10 $L(\lambda,\mathfrak{s})$ and ${}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s})$ are subquotients of $L(\lambda,\mathfrak{s})_{F_{\alpha_2}}$. Let $\nu \in \operatorname{Supp}_{\mathrm{ess}}({}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s})) \cap \operatorname{Supp}_{\mathrm{ess}}(L(\lambda,\mathfrak{s}))$. Then by Lemma 4.22 $L(\lambda,\mathfrak{s})_\nu \cong$

$(L(\lambda,\mathfrak{s})_{F_{\alpha_2}})_\nu \cong ({}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s}))_\nu$ so by Theorem 2.7 $L(\lambda,\mathfrak{s}) \cong {}^{s_{\alpha_2}}L(s_{\alpha_2}.\lambda,\mathfrak{s})$. This is a contradiction by looking at weights of the modules. $\square$

Theorem 8.17 *Let $\mathfrak{g}$ be a simple Lie algebra of type C_n (i.e. $\mathfrak{g} = \mathfrak{sp}(2n)$). Let $\alpha_1,\dots,\alpha_n$ be the simple roots such that α_i is connected to α_{i+1} and α_1 is long – i.e. the Dynkin diagram of C_n is*



Let $\lambda \in X$. $L(\lambda)$ is infinite dimensional and admissible if and only if

- $\lambda(K_{\alpha_i}) \in \pm q^{\mathbb{N}}$ *for* $1 < i \leq n$
- $\lambda(K_{\alpha_1}) \in \pm q_{\alpha_1}^{1/2+\mathbb{Z}} = \pm q^{1+2\mathbb{Z}}$
- $\lambda(K_{\alpha_1+\alpha_2}) \in \pm q^{\mathbb{Z}_{\geq -2}}$

or equivalently $q^\rho\lambda(K_\beta) \in \pm q^{\mathbb{Z}_{>0}}$ for every short root $\beta \in \Phi^+$ and $\lambda(K_{\beta'}) \in \pm q^{1+2\mathbb{Z}}$ for every long root $\beta' \in \Phi^+$.

Proof. Assume $\lambda(K_{\alpha_i}) \notin \pm q^{\mathbb{N}}$ for some $i > 1$. Then by Lemma 8.11 there exists a λ' such that $L(\lambda')$ is admissible and such that $\lambda'(K_{\alpha_2}) \notin q^{\mathbb{N}}$. Let $\mathfrak{s}$ be the Lie algebra $\mathfrak{sp}(4)$ with simple roots α_2 and α_1. Let U be the subalgebra of U_q generated by $F_{\alpha_1}, F_{\alpha_2}, K_{\alpha_1}, K_{\alpha_2}, E_{\alpha_1}, E_{\alpha_2}$. Then $U \cong U_q(\mathfrak{s})$ as algebras and $Uv_{\lambda'}$ contains the simple highest weight $U_q(\mathfrak{s})$-module $L(\lambda',\mathfrak{s})$ of highest weight λ' (restricted to $U^0_{q_\alpha}(\mathfrak{s})$) as a subquotient. Since $L(\lambda')$ is admissible so is $Uv_{\lambda'}$ hence $L(\lambda',\mathfrak{s})$ is admissible. So $\lambda'(K_{\alpha_2}) \in \pm q^{\mathbb{N}}$ by Lemma 8.8. A contradiction. So we have proven that $\lambda(K_{\alpha_i}) \in \pm q^{\mathbb{N}}$ for $1 < i \leq n$ is a neccesary condition. We get also from Lemma 8.8 that $\lambda(K_{\alpha_1}) \in q^{1+2\mathbb{Z}}$ and $q^3\lambda(K_{\alpha_1+\alpha_2}) = q^\rho\lambda(K_{\alpha_1+\alpha_2}) \in \pm q^{\mathbb{Z}_{>0}}$ which shows that the two other conditions are neccesary.

Now assume we have a weight $\lambda \in X$ that satisfies the above. So $\lambda(K_{\alpha_1}) = q^{-1+r}$ for some $r \in 2\mathbb{Z}$. We can assume $r \in \mathbb{N}$ by Lemma 5.11 and Proposition 5.6 (if $r < 0$ replace λ with $s_1.\lambda$, $L(\lambda)$ is admissible if and only if $L(s_1.\lambda)$ is). We have $\lambda = \omega^+\lambda_0$ for some dominant integral weight λ_0 and $L(\lambda)$ is a subquotient of $L(\omega^+) \otimes L(\lambda_0)$. Since $L(\omega^+)$ is admissible and $L(\lambda_0)$ is finite dimensional $L(\omega^+) \otimes L(\lambda_0)$ is admissible and since $L(\lambda)$ is a subquotient of $L(\omega^+) \otimes L(\lambda_0)$, $L(\lambda)$ is admissible as well. $\square$

Corollary 8.18 *Let $\mathfrak{g}$ be a simple Lie algebra of type C_n (i.e. $\mathfrak{g} = \mathfrak{sp}(2n)$). Let $\alpha_1,\dots,\alpha_n$ be the simple roots such that α_i is connected to α_{i+1} and α_1 is long.*

Let $\beta_j = \alpha_1 + \cdots + \alpha_j$, $j = 1,\dots,n$ and $\Sigma = \{\beta_1,\dots,\beta_n\}$. Let $F_{\beta_j} = T_{s_1}\cdots T_{s_{j-1}}(F_{\alpha_j})$ and let $F_\Sigma = \{q^a F_{\beta_1}^{a_1}\cdots F_{\beta_n}^{a_n} | a_i \in \mathbb{N}, a \in \mathbb{Z}\}$ be the corresponding Ore subset. Then Σ is a set of commuting roots that is a basis of Q with corresponding Ore subset F_Σ.

Let L be a simple torsion free module. Then there exists a $\lambda \in X$ with $\lambda(K_\beta) \in \pm q^{\mathbb{N}}$ for all short $\beta \in \Phi^+$ and $\lambda(K_\gamma) \in \pm q^{1+2\mathbb{Z}}$ for all long $\gamma \in \Phi^+$ and a $\mathbf{b} \in (\mathbb{C}^*)^n$ such that

$$L \cong \varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$$

Proof. By Theorem 5.12 there exists a $\lambda \in X$ such that $\mathcal{EXT}(L) \cong \mathcal{EXT}(L(\lambda))$. By Proposition 5.6 $L(\lambda)$ is admissible and by Theorem 8.17 λ is as described in the statement of the corollary. Now the result follows just like in the proof of Corollary 5.13. $\square$

In Section 10 we determine all $\mathbf{b} \in (\mathbb{C}^*)^n$ such that $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ is torsion free (with Σ and λ as above in Corollary 8.18). By the corollary this classifies all simple torsion free modules for type C.

9 Classification of simple torsion free modules. Type A.

In this section we assume $\mathfrak{g} = \mathfrak{sl}_{n+1}$ with $n \geq 2$. Let $\Pi = \{\alpha_1, \dots, \alpha_n\}$ denote the simple roots such that $(\alpha_i|\alpha_{i+1}) = -1$, $i = 1, \dots, n-1$. Set $\beta_j = s_1 \cdots s_{j-1}(\alpha_j) = \alpha_1 + \cdots + \alpha_j$, then $\Sigma = \{\beta_1, \dots, \beta_n\}$ is a set of commuting roots with corresponding root vectors $F_{\beta_j} = T_{s_1} \cdots T_{s_{j-1}}(F_{\alpha_j})$. We will show some commutation formulas and use these to calculate $\varphi_{F_\Sigma,\mathbf{b}}$ on all simple root vectors. This will allow us to determine exactly for which $\mathbf{b} \in (\mathbb{C}^*)^n$, $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ is torsion free, see Theorem 9.8.

Choose a reduced expression of w_0 starting with $s_1 \cdots s_n$ and define roots $\gamma_1, \dots, \gamma_N$ and root vectors $F_{\gamma_1}, \dots, F_{\gamma_N}$ from this expression. Note that $F_{\beta_i} = F_{\gamma_i}$ for $i = 1, \dots, n$.

Proposition 9.1 *Let $i \in \{2, \dots, n\}$ and $j \in \{1, \dots, n\}$.*

$$[F_{\alpha_i}, F_{\beta_j}]_q = \begin{cases} F_{\beta_i}, & \textit{if } j = i-1 \\ 0, & \textit{otherwise} \end{cases}$$

and

$$[E_{\alpha_i}, F_{\beta_j}] = \begin{cases} F_{\beta_{i-1}} K_{\alpha_i}^{-1}, & \textit{if } j = i \\ 0, & \textit{otherwise.} \end{cases}$$

Proof. We will show the proposition for the F's first and then for the E's.

Assume first that $j < i-1$. Then clearly $[F_{\alpha_i}, F_{\beta_j}]_q = [F_{\alpha_i}, F_{\beta_j}] = 0$ since α_i is not connected to any of the simple roots $\alpha_1, \dots, \alpha_j$ appearing in β_j.

Then assume $j \geq i$. We must have $\alpha_i = \gamma_k$ for some $k > n$ since $\{\gamma_1, \dots, \gamma_N\} = \Phi^+$. By Theorem 3.2 $[F_{\alpha_i}, F_{\beta_j}]_q$ is a linear combination of monomials of the form $F_{\gamma_{j+1}}^{a_{j+1}} \cdots F_{\gamma_{k-1}}^{a_{k-1}}$. For a monomial of this form to appear with nonzero coefficient we must have

$$\sum_{h=j+1}^{k-1} a_h \gamma_h = \alpha_i + \beta_j = \alpha_1 + \cdots + \alpha_{i-1} + 2\alpha_i + \alpha_{i+1} + \dots \alpha_j.$$

For this to be possible one of the positive roots γ_s, $j < s < k$ must be equal to $\alpha_1 + \alpha_2 + \cdots + \alpha_m$ for some $m \leq j$ but $\alpha_1 + \alpha_2 + \cdots + \alpha_m = \gamma_m$ by construction and $m \leq j < s$ so $m \neq s$. We conclude that this is not possible.

Finally we investigate the case when $j = i - 1$. We have

$$\begin{aligned}[F_{\alpha_i}, F_{\beta_{i-1}}]_q =&[T_{s_1} \cdots T_{s_{i-2}}(F_{\alpha_i}), T_{s_1} \cdots T_{s_{i-2}}(F_{\alpha_{i-1}})]_q \\ =&T_{s_1} \cdots T_{s_{i-2}} \big([F_{\alpha_i}, F_{\alpha_{i-1}}]_q\big) \\ =&T_{s_1} \cdots T_{s_{i-2}} T_{s_{i-1}}(F_{\alpha_i}) \\ =&F_{\beta_i}.\end{aligned}$$

For the E's: Assume first $j < i$: Since F_{β_j} is a polynomial in $F_{\alpha_1}, \ldots, F_{\alpha_j}$, E_{α_i} commutes with F_{β_j} when $j < i$.

Assume then $j = i$: We have by the above

$$F_{\beta_i} = [F_{\alpha_i}, F_{\beta_{i-1}}]_q$$

so

$$\begin{aligned}[E_{\alpha_i}, F_{\beta_i}] =&[E_{\alpha_i}, (F_{\alpha_i} F_{\beta_{i-1}} - q^{-(\beta_{i-1}|\alpha_i)} F_{\beta_{i-1}} F_{\alpha_i})] \\ =&[E_{\alpha_i}, F_{\alpha_i}] F_{\beta_{i-1}} - q F_{\beta_{i-1}} [E_{\alpha_i}, F_{\alpha_i}] \\ =&\frac{K_{\alpha_i} - K_{\alpha_i}^{-1}}{q - q^{-1}} F_{\beta_{i-1}} - q F_{\beta_{i-1}} \frac{K_{\alpha_i} - K_{\alpha_i}^{-1}}{q - q^{-1}} \\ =&F_{\beta_{i-1}} \frac{q K_{\alpha_i} - q^{-1} K_{\alpha_i}^{-1} - q K_{\alpha_i} + q K_{\alpha_i}^{-1}}{q - q^{-1}} \\ =&F_{\beta_{i-1}} K_{\alpha_i}^{-1}.\end{aligned}$$

Finally assume $j > i$: Observe first that we have

$$T_{s_{i+1}} \cdots T_{s_{j-1}} F_{\alpha_j} = \sum_{s=1}^{m} u_s F_{\alpha_{i+1}} u_s'$$

for some $m \in \mathbb{N}$ and some u_s, u_s' that are polynomials in $F_{\alpha_{i+2}}, \ldots F_{\alpha_j}$. Note that $T_{s_i}(u_s) = u_s$ and $T_{s_i}(u_s') = u_s'$ for all s since α_i is not connected to any of the simple roots $\alpha_{i+2}, \ldots \alpha_j$. So

$$\begin{aligned}T_{s_i} T_{s_{i+1}} \cdots T_{s_{j-1}} F_{\alpha_j} =&T_{s_i}\left(\sum_{s=1}^{m} u_s F_{\alpha_{i+1}} u_s'\right) \\ =&\sum_{s=1}^{m} u_s T_{s_i}(F_{\alpha_{i+1}}) u_s' \\ =&\sum_{s=1}^{m} u_s (F_{\alpha_{i+1}} F_{\alpha_i} - q F_{\alpha_i} F_{\alpha_{i+1}}) u_s' \\ =&\sum_{s=1}^{m} u_s F_{\alpha_{i+1}} u_s' F_{\alpha_i} - q F_{\alpha_i} \sum_{s=1}^{m} u_s F_{\alpha_{i+1}} u_s' \\ =&T_{s_{i+1}} \cdots T_{s_{j-1}}(F_{\alpha_j}) F_{\alpha_i} - q F_{\alpha_i} T_{s_{i+1}} \cdots T_{s_{j-1}}(F_{\alpha_j}).\end{aligned}$$

Thus we see that

$$\begin{aligned}F_{\beta_j} =&T_{s_1} \ldots T_{s_i} \cdots T_{s_{j-1}}(F_{\alpha_j}) \\ =&T_{s_{i+1}} \cdots T_{s_{j-1}}(F_{\alpha_j}) T_{s_1} \cdots T_{s_{i-1}}(F_{\alpha_i}) - q T_{s_1} \cdots T_{s_{i-1}}(F_{\alpha_i}) T_{s_{i+1}} \cdots T_{s_{j-1}}(F_{\alpha_j}) \\ =&T_{s_{i+1}} \cdots T_{s_{j-1}}(F_{\alpha_j}) F_{\beta_i} - q F_{\beta_i} T_{s_{i+1}} \cdots T_{s_{j-1}}(F_{\alpha_j})\end{aligned}$$

and therefore

$$\begin{aligned}[E_{\alpha_i}, F_{\beta_j}] =& T_{s_{i+1}} \cdots T_{s_{j-1}}(F_{\alpha_j})[E_{\alpha_i}, F_{\beta_i}] - q[E_{\alpha_i}, F_{\beta_i}]T_{s_{i+1}} \cdots T_{s_{j-1}}(F_{\alpha_j})\\ =& T_{s_{i+1}} \cdots T_{s_{j-1}}(F_{\alpha_j})F_{\beta_{i-1}}K_{\alpha_i}^{-1} - qF_{\beta_{i-1}}K_{\alpha_i}^{-1}T_{s_{i+1}} \cdots T_{s_{j-1}}(F_{\alpha_j})\\ =& F_{\beta_{i-1}}T_{s_{i+1}} \cdots T_{s_{j-1}}(F_{\alpha_j})K_{\alpha_i}^{-1} - F_{\beta_{i-1}}T_{s_{i+1}} \cdots T_{s_{j-1}}(F_{\alpha_j})K_{\alpha_i}^{-1}\\ =& 0.\end{aligned}$$

□

Proposition 9.2 *Let $i \in \{2, \dots, n\}$. Let $a \in \mathbb{Z}_{>0}$. Then*

$$[F_{\alpha_i}, F_{\beta_{i-1}}^a]_q = [a]F_{\beta_{i-1}}^{a-1}F_{\beta_i}$$

and for $b \in \mathbb{C}^*$

$$\varphi_{F_{\beta_{i-1}},b}(F_{\alpha_i}) = bF_{\alpha_i} + \frac{b - b^{-1}}{q - q^{-1}}F_{\beta_{i-1}}^{-1}F_{\beta_i}.$$

Proof. The first claim is proved by induction over a. $a = 1$ is shown in Proposition 9.1. The induction step:

$$\begin{aligned}F_{\alpha_i}F_{\beta_{i-1}}^{a+1} =& \left(q^aF_{\beta_{i-1}}^aF_{\alpha_i} + [a]F_{\beta_{i-1}}^{a-1}F_{\beta_i}\right)F_{\beta_{i-1}}\\ =& q^{a+1}F_{\beta_{i-1}}^{a+1}F_{\alpha_i} + q^aF_{\beta_{i-1}}^aF_{\beta_i} + q^{-1}[a]F_{\beta_{i-1}}^aF_{\beta_i}\\ =& q^{a+1}F_{\beta_{i-1}}^{a+1}F_{\alpha_i} + [a+1]F_{\beta_{i-1}}^aF_{\beta_i}.\end{aligned}$$

So we have proved the first claim. We get then for $a \in \mathbb{Z}_{>0}$

$$\varphi_{F_{\beta_{i-1}},q^a}(F_{\alpha_i}) = F_{\beta_{i-1}}^{-a}F_{\alpha_i}F_{\beta_{i-1}}^a = q^aF_{\alpha_i} + \frac{q^a - q^{-a}}{q - q^{-1}}F_{\beta_{i-1}}^{-1}F_{\beta_i}.$$

Using the fact that $\varphi_{F_{\beta_{i-1}},b}(F_{\alpha_i})$ is Laurent polynomial in b we get the second claim of the proposition. □

Proposition 9.3 *Let $i \in \{2, \dots, n\}$. Let $a \in \mathbb{Z}_{>0}$. Then*

$$[E_{\alpha_i}, F_{\beta_i}^a] = q^{a-1}[a]F_{\beta_i}^{a-1}F_{\beta_{i-1}}K_{\alpha_i}^{-1}$$

and for $b \in \mathbb{C}^*$

$$\varphi_{F_{\beta_i},b}(E_{\alpha_i}) = E_{\alpha_i} + q^{-1}b\frac{b - b^{-1}}{q - q^{-1}}F_{\beta_i}^{-1}F_{\beta_{i-1}}K_{\alpha_i}^{-1}.$$

Proof. The first claim is proved by induction over a. $a = 1$ is shown in Proposition 9.1. The induction step:

$$\begin{aligned}E_{\alpha_i}F_{\beta_i}^{a+1} =& \left(F_{\beta_i}^aE_{\alpha_i} + q^{a-1}[a]F_{\beta_i}^{a-1}F_{\beta_{i-1}}K_{\alpha_i}^{-1}\right)F_{\beta_i}\\ =& F_{\beta_i}^{a+1}E_{\alpha_i} + F_{\beta_i}^aF_{\beta_{i-1}}K_{\alpha_i}^{-1} + q^{a+1}[a]F_{\beta_i}^aF_{\beta_{i-1}}K_{\alpha_i}^{-1}\\ =& F_{\beta_i}^{a+1}E_{\alpha_i} + q^a(q^{-a} + q[a])F_{\beta_i}^aF_{\beta_{i-1}}K_{\alpha_i}^{-1}\\ =& F_{\beta_i}^{a+1}E_{\alpha_i} + q^a[a+1]F_{\beta_i}^aF_{\beta_{i-1}}K_{\alpha_i}^{-1}.\end{aligned}$$

This proves the first claim. We get then for $a \in \mathbb{Z}_{>0}$

$$\varphi_{F_{\beta_i},q^a}(E_{\alpha_i}) = F_{\beta_i}^{-a} E_{\alpha_i} F_{\beta_i}^a = E_{\alpha_i} + q^{-1} q^a \frac{q^a - q^{-a}}{q - q^{-1}} F_{\beta_i}^{-1} F_{\beta_{i-1}} K_{\alpha_i}^{-1}.$$

Using the fact that $\varphi_{F_{\beta_i},b}(E_{\alpha_i})$ is Laurent polynomial in b we get the second claim of the proposition. ☐

In our classification we don't need to calculate $\varphi_{F_\Sigma,\mathbf{b}}(E_{\alpha_1})$ but for completeness we show what it is in this case in Proposition 9.5. To do this we need the following proposition:

Proposition 9.4 *Let $j \in \{2, \dots, n\}$. Then*

$$[E_{\alpha_1}, F_{\beta_j}] = -q T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) K_{\alpha_1},$$

for $a \in \mathbb{Z}_{>0}$:

$$[E_{\alpha_1}, F_{\beta_j}^a] = -q^{2-a}[a] F_{\beta_j}^{a-1} T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) K_{\alpha_1}$$

and for $b \in \mathbb{C}^*$:

$$\varphi_{F_{\beta_j},b}(E_{\alpha_1}) = E_{\alpha_1} - q^2 b \frac{b - b^{-1}}{q - q^{-1}} F_{\beta_j}^{-1} T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) K_{\alpha_1}.$$

Proof. Like in the proof of Proposition 9.1 we see that

$$T_{s_2} \cdots T_{s_{j-1}} F_{\alpha_j} = \sum_{s=1}^m u_s F_{\alpha_2} u_s'$$

for some $m \in \mathbb{N}$ and some u_s, u_s' that are polynomials in $F_{\alpha_3}, \dots F_{\alpha_j}$. Note that $T_{s_1}(u_s) = u_s$ and $T_{s_1}(u_s') = u_s'$ for all s since α_1 is not connected to any of the simple roots $\alpha_3, \dots \alpha_j$. So

$$\begin{aligned}
T_{s_1} T_{s_2} \cdots T_{s_{j-1}} F_{\alpha_j} =& T_{s_1}\left(\sum_{s=1}^m u_s F_{\alpha_2} u_s'\right) \\
=& \sum_{s=1}^m u_s T_{s_1}(F_{\alpha_2}) u_s' \\
=& \sum_{s=1}^m u_s (F_{\alpha_2} F_{\alpha_1} - q F_{\alpha_1} F_{\alpha_2}) u_s' \\
=& \sum_{s=1}^m u_s F_{\alpha_2} u_s' F_{\alpha_1} - q F_{\alpha_1} \sum_{s=1}^m u_s F_{\alpha_2} u_s' \\
=& T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) F_{\alpha_1} - q F_{\alpha_1} T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}).
\end{aligned}$$

Thus

$$\begin{aligned}
[E_{\alpha_1}, F_{\beta_j}] =& T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) [E_{\alpha_1}, F_{\alpha_1}] - q [E_{\alpha_1}, F_{\alpha_1}] T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) \\
=& T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) \frac{K_{\alpha_1} - K_{\alpha_1}^{-1}}{q - q^{-1}} - q \frac{K_{\alpha_1} - K_{\alpha_1}^{-1}}{q - q^{-1}} T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) \\
=& T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) \frac{K_{\alpha_1} - K_{\alpha_1}^{-1} - q^2 K_{\alpha_1} + K_{\alpha_1}^{-1}}{q - q^{-1}} \\
=& - q T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) K_{\alpha_1}.
\end{aligned}$$

Note that $T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j})$ is a polynomial in $F_{\alpha_2}, \dots, F_{\alpha_j}$. By Proposition 9.1 $[F_{\alpha_i}, F_{\beta_j}]_q = [F_{\alpha_i}, F_{\beta_j}] = 0$ for $1 < i < j$ and $[F_{\alpha_j}, F_{\beta_j}]_q = F_{\alpha_j} F_{\beta_j} - q^{-1} F_{\beta_j} F_{\alpha_j} = 0$ so

$$\begin{aligned} T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) F_{\beta_j} - q^{-1} F_{\beta_j} T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) =& [T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}), F_{\beta_j}]_q \\ =& 0. \end{aligned}$$

The second claim is by induction on a:

$$\begin{aligned} E_{\alpha_1} F_{\beta_j}^{a+1} =& \left(F_{\beta_j}^a E_{\alpha_1} - q^{2-a}[a] F_{\beta_j}^{a-1} T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) K_{\alpha_1} \right) F_{\beta_j} \\ =& F_{\beta_j}^{a+1} E_{\alpha_1} - q F_{\beta_j}^a T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) K_{\alpha_1} \\ & - q^{-a}[a] F_{\beta_j}^a T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) K_{\alpha_1} \\ =& F_{\beta_j}^{a+1} E_{\alpha_1} - q^{1-a} \left(q^a + q^{-1}[a] \right) F_{\beta_j}^a T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) K_{\alpha_1} \\ =& F_{\beta_j}^{a+1} E_{\alpha_1} - q^{1-a}[a+1] F_{\beta_j}^a T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) K_{\alpha_1}. \end{aligned}$$

So we get for $a \in \mathbb{Z}_{>0}$:

$$\varphi_{F_{\beta_j}, q^a}(E_{\alpha_1}) = F_{\beta_j}^{-a} E_{\alpha_1} F_{\beta_j}^a = E_{\alpha_1} - q^2 q^{-a} \frac{q^a - q^{-a}}{q - q^{-1}} T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) K_{\alpha_1}.$$

Using the fact that $\varphi_{F_{\beta_j}, b}(E_{\alpha_1})$ is Laurent polynomial in b we get the third claim of the proposition. $\square$

We can combine the above propositions in the following proposition

Proposition 9.5 *Let $i \in \{2, \dots, n\}$. For $\mathbf{b} = (b_1, \dots, b_n) \in (\mathbb{C}^*)^n$*

$$\begin{aligned} \varphi_{F_\Sigma, \mathbf{b}}(F_{\alpha_i}) =& b_i^{-1} b_{i+1}^{-1} \cdots b_n^{-1} \varphi_{F_{\beta_{i-1}}, b_{i-1}}(F_{\alpha_i}) \\ =& b_i^{-1} b_{i+1}^{-1} \cdots b_n^{-1} (b_{i-1} F_{\alpha_i} + \frac{b_{i-1} - b_{i-1}^{-1}}{q - q^{-1}} F_{\beta_{i-1}}^{-1} F_{\beta_i}) \\ \varphi_{F_\Sigma, \mathbf{b}}(E_{\alpha_i}) =& \varphi_{F_{\beta_i}, b_i}(E_{\alpha_i}) = E_{\alpha_i} + q^{-1} b_i \frac{b_i - b_i^{-1}}{q - q^{-1}} F_{\beta_i}^{-1} F_{\beta_{i-1}} K_{\alpha_i}^{-1}. \end{aligned}$$

Furthermore

$$\varphi_{F_\Sigma, \mathbf{b}}(F_{\alpha_1}) = b_2 \cdots b_n F_{\alpha_1}$$

and

$$\begin{aligned} \varphi_{F_\Sigma, \mathbf{b}}(E_{\alpha_1}) =& E_{\alpha_1} - q^2 \sum_{j=2}^n b_j b_{j+1}^{-1} \cdots b_n^{-1} \frac{b_j - b_j^{-1}}{q - q^{-1}} F_{\beta_j}^{-1} T_{s_2} \cdots T_{s_{j-1}}(F_{\alpha_j}) K_{\alpha_1} \\ & + b_2^{-1} \cdots b_n^{-1} F_{\beta_1}^{-1} \frac{(b_1 - b_1^{-1})(q b_1^{-1} \cdots b_n^{-1} K_{\alpha_1} - q^{-1} b_1 \cdots b_n K_{\alpha_1}^{-1})}{(q - q^{-1})^2}. \end{aligned}$$

Proof. The first two equations follow from Proposition 9.1, Proposition 9.2 and Proposition 9.3. The third follows because $F_{\alpha_1} = F_{\beta_1}$ q-commutes with all the other root vectors $F_{\beta_2}, \dots, F_{\beta_n}$ (see also the discussion before Definition 4.19).

For the last equation we use Proposition 9.4:

$$\begin{aligned}\varphi_{F_\Sigma,\mathbf{b}}(E_{\alpha_1}) =&\varphi_{F_{\beta_n},b_n}\circ\cdots\circ\varphi_{F_{\beta_1},b_1}(E_{\alpha_1})\\ =&\varphi_{F_{\beta_n},b_n}\circ\cdots\circ\varphi_{F_{\beta_2},b_2}\left(E_{\alpha_1}-F_{\beta_1}^{-1}\frac{(b_1-b_1^{-1})(qb_1^{-1}K_{\alpha_1}-q^{-1}b_1K_{\alpha_1}^{-1})}{(q-q^{-1})^2}\right)\\ =&\varphi_{F_{\beta_n},b_n}\circ\cdots\circ\varphi_{F_{\beta_3},b_3}(E_{\alpha_1}-q^2b_2\frac{b_2-b_2^{-1}}{q-q^{-1}}F_{\beta_2}^{-1}F_{\alpha_2}K_{\alpha_1}\\ &-b_2^{-1}F_{\beta_1}^{-1}\frac{(b_1-b_1^{-1})(qb_1^{-1}b_2^{-1}K_{\alpha_1}-q^{-1}b_1b_2K_{\alpha_1}^{-1})}{(q-q^{-1})^2})\\ &\vdots\\ =&E_{\alpha_1}-q^2\sum_{j=2}^{n}b_jb_{j+1}^{-1}\cdots b_n^{-1}\frac{b_j-b_j^{-1}}{q-q^{-1}}F_{\beta_j}^{-1}T_{s_2}\cdots T_{s_{j-1}}(F_{\alpha_j})K_{\alpha_1}\\ &-b_2^{-1}\cdots b_n^{-1}F_{\beta_1}^{-1}\frac{(b_1-b_1^{-1})(qb_1^{-1}\cdots b_n^{-1}K_{\alpha_1}-q^{-1}b_1\cdots b_nK_{\alpha_1}^{-1})}{(q-q^{-1})^2}\square\end{aligned}$$

Proposition 9.6 *Let λ be a weight such that $\lambda(K_{\alpha_i})\in\pm q^{\mathbb{N}}$ for $i=2,\dots,n$ and $\lambda(K_{\alpha_1})\notin\pm q^{\mathbb{N}}$. Let $\mathbf{b}=(b_1,\dots,b_n)\in(\mathbb{C}^*)^n$. Let $i\in\{2,\dots,n\}$. Then E_{α_i} acts injectively on the U_q-module $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ if and only if $b_i\notin\pm q^{\mathbb{Z}}$ and F_{α_i} acts injectively on $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ if and only if $b_{i-1}\notin\pm q^{\mathbb{Z}}$.*

Proof. By Proposition 4.10 and Corollary 4.20 a root vector acts injectively on the U_q-module

$$\varphi_{F_\Sigma,(b_1,\dots,b_n)}.L(\lambda)_{F_\Sigma}$$

if and only if it acts injectively on

$$\varphi_{F_\Sigma,(\varepsilon_1q^{i_1}b_1,\dots,\varepsilon_nq^{i_n}b_n)}.L(\lambda)_{F_\Sigma}$$

for any $i_1,\dots,i_n\in\mathbb{Z}$ and $\varepsilon_1,\dots,\varepsilon_n\in\{\pm1\}$.

Assume there exists a $0\neq v\in\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ such that $E_{\alpha_i}v=0$. We have $v=F_{\beta_1}^{a_1}\cdots F_{\beta_n}^{a_n}\otimes v'$ for some $a_1,\dots,a_n\in\mathbb{Z}_{\leq0}$ and some $v'\in L(\lambda)$. So $E_{\alpha_i}v=0$ implies

$$0=\varphi_{F_\Sigma,\mathbf{b}}(E_{\alpha_i})F_{\beta_1}^{a_1}\cdots F_{\beta_n}^{a_n}\otimes v'=F_{\beta_1}^{a_1}\cdots F_{\beta_n}^{a_n}\otimes\varphi_{F_\Sigma,\mathbf{c}}(E_{\alpha_i})v'$$

where $\mathbf{c}=(q^{a_1}b_1,\dots,q^{a_n}b_n)$. So there exists a $v'\in L(\lambda)$ such that $\varphi_{F_\Sigma,\mathbf{c}}(E_{\alpha_i})v'=0$. That is

$$\left(E_{\alpha_i}+q^{-1}c_i\frac{c_i-c_i^{-1}}{q-q^{-1}}F_{\beta_i}^{-1}F_{\beta_{i-1}}K_{\alpha_i}^{-1}\right)v'=0$$

or equivalently

$$F_{\beta_i}E_{\alpha_i}v'=q^{-1}c_i\frac{c_i^{-1}-c_i}{q-q^{-1}}F_{\beta_{i-1}}K_{\alpha_i}^{-1}v'.$$

Since $L(\lambda)$ is a highest weight module we have some $r\in\mathbb{N}$ such that $E_{\alpha_i}^rv'\neq0$ and $E_{\alpha_i}^{r+1}v'=0$. Fix this r. We get

$$E_{\alpha_i}^{(r)}F_{\beta_i}E_{\alpha_i}v'=E_{\alpha_i}^{(r)}q^{-1}c_i\frac{c_i^{-1}-c_i}{q-q^{-1}}F_{\beta_{i-1}}K_{\alpha_i}^{-1}v'$$

and calculating the right hand side and left hand side we get

$$q^{r-1}[r]F_{\beta_{i-1}}K_{\alpha_i}^{-1}E_{\alpha_i}^{(r)}v' = q^{-1+2r}c_i\frac{c_i^{-1}-c_i}{q-q^{-1}}F_{\beta_{i-1}}K_{\alpha_i}^{-1}E_{\alpha_i}^{(r)}v'.$$

So we must have

$$q^{r-1}[r] = q^{-1+2r}c_i\frac{c_i^{-1}-c_i}{q-q^{-1}}$$

or equivalently $c_i = \pm q^{-r}$. Since $c_i \in q^{\mathbb{Z}}b_i$ we have proved the first claim.

The other claim is shown similarly (see e.g. the calculations done in the proof of Proposition 10.5. The calculations will be the same in this case). $\square$

Proposition 9.7 *Let M be a weight U_q-module of finite Jordan-Hölder length with finite dimensional weight spaces. Let $\alpha \in \Pi$. If E_α and F_α both act injectively on M then E_α and F_α act injectively on every composition factor of M.*

Proof. Let V be a simple U_q-submodule of M. Let μ be a weight of V. Then V_μ is a simple $(U_q)_0$-module by Theorem 2.7 and $E_\alpha F_\alpha$ and $F_\alpha E_\alpha$ act injectively on V_μ by assumption. Since $\dim M_\mu < \infty$ this implies that $F_\alpha E_\alpha$ and $E_\alpha F_\alpha$ act injectively on the $(U_q)_0$ module $(M/V)_\mu \cong M_\mu/V_\mu$. Since M/V is the sum of its weight spaces this implies that $E_\alpha F_\alpha$ and $F_\alpha E_\alpha$ act injectively on M/V. This in turn implies that E_α and F_α act injectively on M/V. Doing induction on the Jordan-Hölder length of M finishes the proof. $\square$

The above proposition is true for a general simple Lie algebra $\mathfrak{g}$ and we will use it in the next section as well.

Theorem 9.8 *Let λ be a weight such that $\lambda(K_{\alpha_i}) \in \pm q^{\mathbb{N}}$ for $i = 2, \ldots, n$ and $\lambda(K_{\alpha_1}) \notin \pm q^{\mathbb{N}}$. Let $\mathbf{b} = (b_1, \ldots, b_n) \in (\mathbb{C}^*)^n$. Then $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ is simple and torsion free if and only if $b_i \notin \pm q^{\mathbb{Z}}$, $i = 1, \ldots, n$ and $\lambda(K_{\alpha_1})^{-1}b_1 \cdots b_n \notin \pm q^{\mathbb{Z}}$.*

Proof. By Proposition 5.10 $L(\lambda)$ is a subquotient of

$${}^{\overline{s_1}}\left(\varphi_{F_\Sigma,(\lambda(K_{\alpha_1}),1,\ldots,1)}.L(\lambda)_{F_\Sigma}\right).$$

So by Lemma 5.3 we get (using that $L(\lambda) = {}^{s_1}\left({}^{\overline{s_1}}L(\lambda)\right)$) for any $\mathbf{c} = (c_1, \ldots, c_n) \in (\mathbb{C}^*)^n$

$$\left(\varphi_{F_\Sigma,\mathbf{c}}.L(\lambda)_{F_\Sigma}\right)^{ss} \cong {}^{\overline{s_1}}\left(\varphi_{F_\Sigma,(\lambda(K_{\alpha_1})c_1^{-1}\cdots c_n^{-1},c_2,\ldots,c_n)}.L(\lambda)_{F_\Sigma}\right)^{ss}.$$

We have $\lambda(K_{\alpha_2}) = \varepsilon q^r$ for some $r \in \mathbb{N}$ and some $\varepsilon \in \{\pm 1\}$. We see in the proof of Lemma 5.11 that $L(\lambda)$ is a subqoutient of

$${}^{\overline{s_2}}\left(\varphi_{F_\Sigma,(\varepsilon,\varepsilon,1,\ldots,1)}.L(\lambda)_{F_\Sigma}\right).$$

We get by Lemma 5.3 (using that $L(\lambda) = {}^{s_2}\left({}^{\overline{s_2}}L(\lambda)\right)$) for any $\mathbf{c} = (c_1, \ldots, c_n) \in (\mathbb{C}^*)^n$

$$\left(\varphi_{F_\Sigma,\mathbf{c}}.L(\lambda)_{F_\Sigma}\right)^{ss} \cong {}^{\overline{s_2}}\left(\varphi_{F_\Sigma,(\varepsilon c_2,\varepsilon c_1,c_3,\ldots,c_n)}.L(\lambda)_{F_\Sigma}\right)^{ss}.$$

Combining the above we get

$$\begin{aligned}(\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma})^{ss} &\cong {}^{\overline{s_2}}\left(\varphi_{F_\Sigma,(\varepsilon b_2,\varepsilon b_1,b_3,\dots,b_n)}.L(\lambda)_{F_\Sigma}\right)^{ss} \\ &\cong {}^{\overline{s_2}}\left({}^{\overline{s_1}}\left(\varphi_{F_\Sigma,(\lambda(K_{\alpha_1})b_1^{-1}\cdots b_n^{-1},\varepsilon b_1,b_3,\dots,b_n)}.L(\lambda)_{F_\Sigma}\right)\right)^{ss} \\ &\cong {}^{\overline{s_1 s_2}}\left(\varphi_{F_\Sigma,(\lambda(K_{\alpha_1})b_1^{-1}\cdots b_n^{-1},\varepsilon b_1,b_3,\dots,b_n)}.L(\lambda)_{F_\Sigma}\right)^{ss}.\end{aligned}$$

Since $T_{s_1}^{-1}T_{s_2}^{-1}(E_{\alpha_1}) = E_{\alpha_2}$ and $T_{s_1}^{-1}T_{s_2}^{-1}(F_{\alpha_1}) = F_{\alpha_2}$ we get by Proposition 9.6 that E_{α_1} acts injectively on ${}^{\overline{s_1 s_2}}\left(\varphi_{F_\Sigma,(\lambda(K_{\alpha_1})b_1^{-1}\cdots b_n^{-1},\varepsilon b_1,b_3,\dots,b_n)}.L(\lambda)_{F_\Sigma}\right)$ if and only if $b_1 \notin \pm q^{\mathbb{Z}}$ and F_{α_1} acts injectively on ${}^{\overline{s_1 s_2}}\left(\varphi_{F_\Sigma,(\lambda(K_{\alpha_1})b_1^{-1}\cdots b_n^{-1},\varepsilon b_1,b_3,\dots,b_n)}.L(\lambda)_{F_\Sigma}\right)$ if and only if $\lambda(K_{\alpha_1})^{-1}b_1\cdots b_n \notin \pm q^{\mathbb{Z}}$.

Assume $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ is torsion free. Then all root vectors act injectively on $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$. We claim $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ is simple: Let $V \subset \varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ be a simple module. Then V is admissible of the same degree d as $L(\lambda)$ by Proposition 5.6 and because all root vectors act injectively $\dim V_{q^\mu\lambda} = d$ for all $\mu \in Q$. So $V = \varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$. Thus $(\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma})^{ss} = \varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$. Then by the above

$$\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma} \cong {}^{\overline{s_1 s_2}}\left(\varphi_{F_\Sigma,(\lambda(K_{\alpha_1})b_1^{-1}\cdots b_n^{-1},\varepsilon b_1,b_3,\dots,b_n)}.L(\lambda)_{F_\Sigma}\right).$$

This shows that when $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ is torsion free we must have $\lambda(K_{\alpha_1})^{-1}b_1\cdots b_n \notin \pm q^{\mathbb{Z}}$. By Proposition 9.6 $b_i \notin \pm q^{\mathbb{Z}}$, $i = 1,\dots,n$.

Assume on the other hand that $b_i \notin \pm q^{\mathbb{Z}}$ for $i \in \{1,\dots,n\}$ and $\lambda(K_{\alpha_1})^{-1}b_1\cdots b_n \notin \pm q^{\mathbb{Z}}$. By Proposition 9.6 we get that the simple root vectors $E_{\alpha_2},\dots,E_{\alpha_n}$ and $F_{\alpha_1},\dots,F_{\alpha_n}$ all act injectively on $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$. We need to show that E_{α_1} acts injectively on the module. By the above

$$(\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma})^{ss} \cong {}^{\overline{s_1 s_2}}\left(\varphi_{F_\Sigma,(\lambda(K_{\alpha_1})b_1^{-1}\cdots b_n^{-1},\varepsilon b_1,b_3,\dots,b_n)}.L(\lambda)_{F_\Sigma}\right)^{ss}$$

and the root vectors $E_{\alpha_1}, F_{\alpha_1}$ act injectively on

$${}^{\overline{s_1 s_2}}\left(\varphi_{F_\Sigma,(\lambda(K_{\alpha_1})b_1^{-1}\cdots b_n^{-1},\varepsilon b_1,b_3,\dots,b_n)}.L(\lambda)_{F_\Sigma}\right).$$

Then by Proposition 9.7 E_{α_1} act injectively on all composition factors of $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$.

Let V be a simple U_q-submodule of $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$. By the above all simple root vectors act injectively on V and then like above this implies $V = \varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ i.e. $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ is simple and torsion free. ☐

By the comments after Corollary 8.15 the above Theorem completes the classification of simple torsion free modules in type A.

10 Classification of simple torsion free modules. Type C.

In this section we assume $\mathfrak{g}$ is of type C_n (i.e. $\mathfrak{g} = \mathfrak{sp}_{2n}$) with $n \geq 2$. Let $\Pi = \{\alpha_1,\dots,\alpha_n\}$ denote the simple roots such that $(\alpha_i|\alpha_{i+1}) = -1$, $i = 2,\dots,n-1$, $\langle \alpha_2, \alpha_1^\vee \rangle = -1$ and $\langle \alpha_1, \alpha_2^\vee \rangle = -2$ i.e. α_1 is long and $\alpha_2,\dots,\alpha_n$ are short.

Set $\beta_j = s_1 \cdots s_{j-1}(\alpha_j) = \alpha_1 + \cdots + \alpha_j$, then $\Sigma = \{\beta_1, \dots, \beta_n\}$ is a set of commuting roots with corresponding root vectors $F_{\beta_j} = T_{s_1} \cdots T_{s_{j-1}}(F_{\alpha_j})$. We will show some commutation formulas and use these to calculate $\varphi_{F_\Sigma,\mathbf{b}}$ on most of the simple root vectors.

Choose a reduced expression of w_0 starting with $s_1 \cdots s_n s_1 \cdots s_{n-1}$ and define root vectors $F_{\gamma_1}, \dots, F_{\gamma_N}$ from this expression. Note that $F_{\beta_i} = F_{\gamma_i}$ for $i = 1, \dots, n$. Note for use in the proposition below that for $j \in \{1, \dots, n-1\}$,

$$\gamma_{n+j} = s_1 \cdots s_n s_1 \cdots s_{j-1}(\alpha_j) = \alpha_1 + 2\alpha_2 + \alpha_3 + \cdots \alpha_{j+1}$$

and

$$\begin{aligned} F_{\gamma_{n+j}} =& T_{s_1} \cdots T_{s_n} T_{s_1} \cdots T_{s_{j-1}}(F_{\alpha_j}) \\ =& T_{s_1} \cdots T_{s_{j+1}} T_{s_1} \cdots T_{s_{j-1}}(F_{\alpha_j}). \end{aligned}$$

In particular $F_{\alpha_1+2\alpha_2} = T_{s_1} T_{s_2}(F_{\alpha_1})$.

Proposition 10.1 *Let $i \in \{2, \dots, n\}$ and $j \in \{1, \dots, n\}$*

$$[F_{\alpha_i}, F_{\beta_j}]_q = \begin{cases} [2]F_{\alpha_1+2\alpha_2}, & \text{if } j = i = 2 \\ F_{\alpha_1+2\alpha_2+\alpha_3+\cdots+\alpha_j}, & \text{if } i = 2 \text{ and } j > 2 \\ F_{\beta_i}, & \text{if } j = i-1 \\ 0, & \text{otherwise} \end{cases}$$

and

$$[E_{\alpha_i}, F_{\beta_j}] = \begin{cases} [2]F_{\beta_1} K_{\alpha_2}^{-1}, & \text{if } j = 2 = i \\ F_{\beta_{i-1}} K_{\alpha_i}^{-1}, & \text{if } j = i > 2 \\ 0, & \text{otherwise.} \end{cases}$$

Proof. We will show the proposition for the F's first and then for the E's. Assume first that $j < i-1$. Then clearly $[F_{\alpha_i}, F_{\beta_j}]_q = [F_{\alpha_i}, F_{\beta_j}] = 0$ since α_i is not connected to any of the simple roots $\alpha_1, \dots, \alpha_j$ appearing in β_j.

Then assume $j \geq i > 2$. We must have $\alpha_i = \gamma_k$ for some $k > n$ since $\{\gamma_1, \dots, \gamma_N\} = \Phi^+$. By Theorem 3.2 $[F_{\alpha_i}, F_{\beta_j}]_q$ is a linear combination of monomials of the form $F_{\gamma_{j+1}}^{a_{j+1}} \cdots F_{\gamma_{k-1}}^{a_{k-1}}$. For a monomial of this form to appear with nonzero coefficient we must have

$$\sum_{h=j+1}^{k-1} a_h \gamma_h = \alpha_i + \beta_j = \alpha_1 + \cdots + \alpha_{i-1} + 2\alpha_i + \alpha_{i+1} + \dots \alpha_j.$$

For this to be possible one of the positive roots γ_s, $j < s < k$ must be equal to $\alpha_1 + \alpha_2 + \cdots + \alpha_m$ for some $m \leq j$ but $\alpha_1 + \alpha_2 + \cdots + \alpha_m = \gamma_m$ by construction and $m \leq j < s$ so $m \neq s$. We conclude that this is not possible.

Assume $j = i-1$. We have

$$\begin{aligned} [F_{\alpha_i}, F_{\beta_{i-1}}]_q =& [T_{s_1} \cdots T_{s_{i-2}}(F_{\alpha_i}), T_{s_1} \cdots T_{s_{i-2}}(F_{\alpha_{i-1}})]_q \\ =& T_{s_1} \cdots T_{s_{i-2}}\left([F_{\alpha_i}, F_{\alpha_{i-1}}]_q\right) \\ =& T_{s_1} \cdots T_{s_{i-2}} T_{s_{i-1}}(F_{\alpha_i}) \\ =& F_{\beta_i}. \end{aligned}$$

Assume $j = 2 = i$. Then

$$\begin{aligned}
[F_{\alpha_2}, F_{\beta_2}]_q =& F_{\alpha_2} F_{\beta_2} - F_{\beta_2} F_{\alpha_2} \\
=& F_{\alpha_2}(F_{\alpha_2} F_{\alpha_1} - q^2 F_{\alpha_1} F_{\alpha_2}) - (F_{\alpha_2} F_{\alpha_1} - q^2 F_{\alpha_1} F_{\alpha_2}) F_{\alpha_2} \\
=& (q^2 F_{\alpha_1} F_{\alpha_2}^2 - q[2] F_{\alpha_2} F_{\alpha_1} F_{\alpha_2} + F_{\alpha_2}^2 F_{\alpha_1}) \\
=& [2] T_{s_2}^{-1}(F_{\alpha_1}) \\
=& [2] T_{s_2}^{-1} T_{s_2} T_{s_1} T_{s_2}(F_{\alpha_1}) \\
=& [2] T_{s_1} T_{s_2}(F_{\alpha_1}) \\
=& [2] F_{\alpha_1 + 2\alpha_2}.
\end{aligned}$$

Assume $i = 2$ and $j = 3$. Then

$$\begin{aligned}
F_{\alpha_1 + 2\alpha_2 + \alpha_3} =& F_{\gamma_{n+2}} \\
=& T_{s_1} T_{s_2} T_{s_3} T_{s_1}(F_{\alpha_2}) \\
=& T_{s_1} T_{s_2} T_{s_1} T_{s_3}(F_{\alpha_2}) \\
=& T_{s_1} T_{s_2} T_{s_1}(F_{\alpha_2} F_{\alpha_3} - q F_{\alpha_3} F_{\alpha_2}) \\
=& F_{\alpha_2} F_{\beta_3} - q F_{\beta_3} F_{\alpha_2}.
\end{aligned}$$

Finally assume $i = 2$ and $j > 3$. We have

$$\begin{aligned}
F_{\alpha_1 + 2\alpha_2 + \alpha_3 + \cdots + \alpha_j} =& F_{\gamma_{n+j-1}} \\
=& T_{s_1} \cdots T_{s_{j-2}} T_{s_{j-1}} T_{s_j} T_{s_1} \cdots T_{s_{j-3}} T_{s_{j-2}}(F_{\alpha_{j-1}}) \\
=& T_{s_1} \cdots T_{s_{j-2}} T_{s_1} \cdots T_{s_{j-3}} T_{s_{j-1}} T_{s_{j-2}} T_{s_j}(F_{\alpha_{j-1}}) \\
=& T_{s_1} \cdots T_{s_{j-2}} T_{s_1} \cdots T_{s_{j-3}} T_{s_{j-1}} T_{s_{j-2}}(F_{\alpha_{j-1}} F_{\alpha_j} - q F_{\alpha_j} F_{\alpha_{j-1}}) \\
=& T_{s_1} \cdots T_{s_{j-2}} T_{s_1} \cdots T_{s_{j-3}}(F_{\alpha_{j-2}} T_{s_{j-1}}(F_{\alpha_j}) - q T_{s_{j-1}}(F_{\alpha_j}) F_{\alpha_{j-2}}) \\
=& F_{\alpha_2} F_{\beta_j} - q F_{\beta_j} F_{\alpha_2} \\
=& [F_{\alpha_2}, F_{\beta_j}]_q
\end{aligned}$$

using the facts that $T_{s_{j-1}} T_{s_{j-2}}(F_{\alpha_{j-1}}) = F_{\alpha_{j-2}}$ and $T_{s_1} \cdots T_{s_{j-2}} T_{s_1} \cdots T_{s_{j-3}}(F_{\alpha_{j-2}}) = F_{\alpha_2}$ by Proposition 8.20 in [Jan96] (The proposition is about the E root vectors but the proposition is true for the F's as well).

For the E's: Assume first $j < i$: Since F_{β_j} is a polynomial in $F_{\alpha_1}, \dots, F_{\alpha_j}$, E_{α_i} commutes with F_{β_j} when $j < i$.

Assume then $j = i$: We have by the above

$$F_{\beta_i} = [F_{\alpha_i}, F_{\beta_{i-1}}]_q$$

so

$$\begin{aligned}
[E_{\alpha_i}, F_{\beta_i}] =& [E_{\alpha_i}, (F_{\alpha_i} F_{\beta_{i-1}} - q^{-(\beta_{i-1}|\alpha_i)} F_{\beta_{i-1}} F_{\alpha_i})] \\
=& [E_{\alpha_i}, F_{\alpha_i}] F_{\beta_{i-1}} - q_{\alpha_{i-1}} F_{\beta_{i-1}} [E_{\alpha_i}, F_{\alpha_i}] \\
=& \frac{K_{\alpha_i} - K_{\alpha_i}^{-1}}{q - q^{-1}} F_{\beta_{i-1}} - q_{\alpha_{i-1}} F_{\beta_{i-1}} \frac{K_{\alpha_i} - K_{\alpha_i}^{-1}}{q - q^{-1}} \\
=& F_{\beta_{i-1}} \frac{q_{\alpha_{i-1}} K_{\alpha_i} - q_{\alpha_{i-1}}^{-1} K_{\alpha_i}^{-1} - q_{\alpha_{i-1}} K_{\alpha_i} + q_{\alpha_{i-1}} K_{\alpha_i}^{-1}}{q - q^{-1}} \\
=& \frac{q_{\alpha_{i-1}} - q_{\alpha_{i-1}}^{-1}}{q - q^{-1}} F_{\beta_{i-1}} K_{\alpha_i}^{-1} \\
=& \begin{cases} [2] F_{\beta_{i-1}} K_{\alpha_i}^{-1}, & \text{if } i = 2 \\ F_{\beta_{i-1}} K_{\alpha_i}^{-1}, & \text{otherwise.} \end{cases}
\end{aligned}$$

Finally assume $j > i$: Observe first that we have

$$T_{s_{i+1}} \cdots T_{s_{j-1}} F_{\alpha_j} = \sum_{s=1}^{m} u_s F_{\alpha_{i+1}} u_s'$$

for some $m \in \mathbb{N}$ and some u_s, u_s' that are polynomials in $F_{\alpha_{i+2}}, \dots F_{\alpha_j}$. Note that $T_{s_i}(u_s) = u_s$ and $T_{s_i}(u_s') = u_s'$ for all s since α_i is not connected to any of the simple roots $\alpha_{i+2}, \dots \alpha_j$. So

$$\begin{aligned}
T_{s_i} T_{s_{i+1}} \cdots T_{s_{j-1}} F_{\alpha_j} =& T_{s_i} \left(\sum_{s=1}^{m} u_s F_{\alpha_{i+1}} u_s' \right) \\
=& \sum_{s=1}^{m} u_s T_{s_i}(F_{\alpha_{i+1}}) u_s' \\
=& \sum_{s=1}^{m} u_s (F_{\alpha_{i+1}} F_{\alpha_i} - q F_{\alpha_i} F_{\alpha_{i+1}}) u_s' \\
=& \sum_{s=1}^{m} u_s F_{\alpha_{i+1}} u_s' F_{\alpha_i} - q F_{\alpha_i} \sum_{s=1}^{m} u_s F_{\alpha_{i+1}} u_s' \\
=& T_{s_{i+1}} \cdots T_{s_{j-1}} (F_{\alpha_j}) F_{\alpha_i} - q F_{\alpha_i} T_{s_{i+1}} \cdots T_{s_{j-1}} (F_{\alpha_j}).
\end{aligned}$$

Thus we see that

$$\begin{aligned}
F_{\beta_j} =& T_{s_1} \dots T_{s_i} \cdots T_{s_{j-1}} (F_{\alpha_j}) \\
=& T_{s_{i+1}} \cdots T_{s_{j-1}} (F_{\alpha_j}) T_{s_1} \cdots T_{s_{i-1}} (F_{\alpha_i}) - q F_{\alpha_i} T_{s_{i+1}} \cdots T_{s_{j-1}} (F_{\alpha_j}) \\
=& T_{s_{i+1}} \cdots T_{s_{j-1}} (F_{\alpha_j}) F_{\beta_i} - q F_{\beta_i} T_{s_{i+1}} \cdots T_{s_{j-1}} (F_{\alpha_j})
\end{aligned}$$

and therefore

$$\begin{aligned}
[E_{\alpha_i}, F_{\beta_j}] =& T_{s_{i+1}} \cdots T_{s_{j-1}} (F_{\alpha_j}) [E_{\alpha_i}, F_{\beta_i}] - q [E_{\alpha_i}, F_{\beta_i}] T_{s_{i+1}} \cdots T_{s_{j-1}} (F_{\alpha_j}) \\
=& [r_i] (T_{s_{i+1}} \cdots T_{s_{j-1}} (F_{\alpha_j}) F_{\beta_{i-1}} K_{\alpha_i}^{-1} - q F_{\beta_{i-1}} K_{\alpha_i}^{-1} T_{s_{i+1}} \cdots T_{s_{j-1}} (F_{\alpha_j})) \\
=& [r_i] (F_{\beta_{i-1}} T_{s_{i+1}} \cdots T_{s_{j-1}} (F_{\alpha_j}) K_{\alpha_i}^{-1} - F_{\beta_{i-1}} T_{s_{i+1}} \cdots T_{s_{j-1}} (F_{\alpha_j}) K_{\alpha_i}^{-1}) \\
=& 0
\end{aligned}$$

where

$$r_i = \begin{cases} 2 & \text{if } i = 2 \\ 1 & \text{otherwise.} \end{cases} \qquad \square$$

Proposition 10.2 *Let $i \in \{2, \dots, n\}$. Let $a \in \mathbb{Z}_{>0}$. Then*

$$[F_{\alpha_i}, F_{\beta_{i-1}}^a]_q = [a]_{\beta_{i-1}} F_{\beta_{i-1}}^{a-1} F_{\beta_i}$$

and for $b \in \mathbb{C}^*$

$$\varphi_{F_{\beta_{i-1}},b}(F_{\alpha_i}) = \begin{cases} b^2 F_{\alpha_2} + \frac{b^2-b^{-2}}{q^2-q^{-2}} F_{\beta_1}^{-1} F_{\beta_2}, & \text{if } i = 2 \\ b F_{\alpha_i} + \frac{b-b^{-1}}{q-q^{-1}} F_{\beta_{i-1}}^{-1} F_{\beta_i}, & \text{otherwise.} \end{cases}$$

Proof. The first claim is proved by induction over a. $a = 1$ is shown in Proposition 9.1. The induction step:

$$\begin{aligned} F_{\alpha_i} F_{\beta_{i-1}}^{a+1} &= \left(q_{\beta_{i-1}}^a F_{\beta_{i-1}}^a F_{\alpha_i} + [a]_{\beta_{i-1}} F_{\beta_{i-1}}^{a-1} F_{\beta_i} \right) F_{\beta_{i-1}} \\ &= q_{\beta_{i-1}}^{a+1} F_{\beta_{i-1}}^{a+1} F_{\alpha_i} + q_{\beta_{i-1}}^a F_{\beta_{i-1}}^a F_{\beta_i} + q_{\beta_{i-1}}^{-1} [a]_{\beta_{i-1}} F_{\beta_{i-1}}^a F_{\beta_i} \\ &= q_{\beta_{i-1}}^{a+1} F_{\beta_{i-1}}^{a+1} F_{\alpha_i} + [a+1]_{\beta_{i-1}} F_{\beta_{i-1}}^a F_{\beta_i}. \end{aligned}$$

So we have proved the first claim. We get then for $a \in \mathbb{Z}_{>0}$:

$$\varphi_{F_{\beta_{i-1}},q^a}(F_{\alpha_i}) = F_{\beta_{i-1}}^{-a} F_{\alpha_i} F_{\beta_{i-1}}^a = q_{\beta_{i-1}}^a F_{\alpha_i} + \frac{q_{\beta_{i-1}}^a - q_{\beta_{i-1}}^{-a}}{q_{\beta_{i-1}} - q_{\beta_{i-1}}^{-1}} F_{\beta_{i-1}}^{-1} F_{\beta_i}.$$

Using the fact that $\varphi_{F_{\beta_{i-1}},b}(F_{\alpha_i})$ is Laurent polynomial in b we get the second claim of the proposition. □

Proposition 10.3 *Let $i \in \{2, \dots, n\}$. Let $a \in \mathbb{Z}_{>0}$. Then*

$$[E_{\alpha_i}, F_{\beta_i}^a] = \begin{cases} q^{a-1}[2][a] F_{\beta_2}^{a-1} F_{\beta_1} K_{\alpha_2}^{-1}, & \text{if } i = 2 \\ q^{a-1}[a] F_{\beta_i}^{a-1} F_{\beta_{i-1}} K_{\alpha_i}^{-1}, & \text{otherwise.} \end{cases}$$

and for $b \in \mathbb{C}^*$

$$\varphi_{F_{\beta_i},b}(E_{\alpha_i}) = \begin{cases} E_{\alpha_2} + q^{-1}[2] b \frac{b-b^{-1}}{q-q^{-1}} F_{\beta_2}^{-1} F_{\beta_1} K_{\alpha_2}^{-1}, & \text{if } i = 2 \\ E_{\alpha_i} + q^{-1} b \frac{b-b^{-1}}{q-q^{-1}} F_{\beta_i}^{-1} F_{\beta_{i-1}} K_{\alpha_i}^{-1}, & \text{otherwise.} \end{cases}$$

Proof. The first claim is proved by induction over a. $a = 1$ is shown in Proposition 9.1. The induction step: For $i > 2$:

$$\begin{aligned} E_{\alpha_i} F_{\beta_i}^{a+1} &= \left(F_{\beta_i}^a E_{\alpha_i} + q^{a-1}[a] F_{\beta_i}^{a-1} F_{\beta_{i-1}} K_{\alpha_i}^{-1} \right) F_{\beta_i} \\ &= F_{\beta_i}^{a+1} E_{\alpha_i} + F_{\beta_i}^a F_{\beta_{i-1}} K_{\alpha_i}^{-1} + q^{a+1}[a] F_{\beta_i}^a F_{\beta_{i-1}} K_{\alpha_i}^{-1} \\ &= F_{\beta_i}^{a+1} E_{\alpha_i} + q^a (q^{-a} + q[a]) F_{\beta_i}^a F_{\beta_{i-1}} K_{\alpha_i}^{-1} \\ &= F_{\beta_i}^{a+1} E_{\alpha_i} + q^a [a+1]_{\alpha_{i-1}} F_{\beta_i}^a F_{\beta_{i-1}} K_{\alpha_i}^{-1}. \end{aligned}$$

For $i = 2$:

$$\begin{aligned}
E_{\alpha_2} F_{\beta_2}^{a+1} &= \left(F_{\beta_2}^{a} E_{\alpha_2} + q^{a-1}[2][a] F_{\beta_2}^{a-1} F_{\beta_1} K_{\alpha_2}^{-1} \right) F_{\beta_2} \\
&= F_{\beta_2}^{a+1} E_{\alpha_2} + [2] F_{\beta_2}^{a} F_{\beta_1} K_{\alpha_2}^{-1} + q^{a+1}[2][a] F_{\beta_2}^{a} F_{\beta_1} K_{\alpha_2}^{-1} \\
&= F_{\beta_2}^{a+1} E_{\alpha_2} + q^{a}[2](q^{-a} + q[a]) F_{\beta_2}^{a} F_{\beta_1} K_{\alpha_2}^{-1} \\
&= F_{\beta_2}^{a+1} E_{\alpha_2} + q^{a}[2][a+1] F_{\beta_2}^{a} F_{\beta_1} K_{\alpha_2}^{-1}.
\end{aligned}$$

This proves the first claim. We get then for $a \in \mathbb{Z}_{>0}$

$$\varphi_{F_{\beta_i}, q^a}(E_{\alpha_i}) = F_{\beta_i}^{-a} E_{\alpha_i} F_{\beta_i}^{a} = \begin{cases} E_{\alpha_2} + q^{-2} q^{2a} \frac{q^{2a} - q^{-2a}}{q - q^{-1}} F_{\beta_2}^{-1} F_{\beta_1} K_{\alpha_2}^{-1}, & \text{if } i = 2 \\ E_{\alpha_i} + q^{-1} q^{a} \frac{q^{a} - q^{-a}}{q - q^{-1}} F_{\beta_i}^{-1} F_{\beta_{i-1}} K_{\alpha_i}^{-1}, & \text{otherwise.} \end{cases}$$

Using the fact that $\varphi_{F_{\beta_i}, b}(E_{\alpha_i})$ is Laurent polynomial in b we get the second claim of the proposition. □

We combine the above propositions in the following proposition

Proposition 10.4 *Let $i \in \{3, \dots, n\}$. For $\mathbf{b} = (b_1, \dots, b_n) \in (\mathbb{C}^*)^n$*

$$\begin{aligned}
\varphi_{F_\Sigma, \mathbf{b}}(F_{\alpha_i}) &= \varphi_{F_{\beta_{i-1}}, b_{i-1}}(F_{\alpha_i}) \\
&= b_{i-1} F_{\alpha_i} + \frac{b_{i-1} - b_{i-1}^{-1}}{q - q^{-1}} F_{\beta_{i-1}}^{-1} F_{\beta_i} \\
\varphi_{F_\Sigma, \mathbf{b}}(E_{\alpha_i}) &= \varphi_{F_{\beta_i}, b_i}(E_{\alpha_i}) = E_{\alpha_i} + q^{-1} b_i \frac{b_i - b_i^{-1}}{q - q^{-1}} F_{\beta_i}^{-1} F_{\beta_{i-1}} K_{\alpha_i}^{-1}.
\end{aligned}$$

Furthermore

$$\varphi_{F_\Sigma, \mathbf{b}}(E_{\alpha_2}) = E_{\alpha_2} + q^{-1}[2] b_2 \frac{b_2 - b_2^{-1}}{q - q^{-1}} F_{\beta_2}^{-1} F_{\beta_1} K_{\alpha_2}^{-1}$$

and

$$\varphi_{F_\Sigma, \mathbf{b}}(F_{\alpha_1}) = b_2 \cdots b_n F_{\alpha_1}.$$

With similar proof as the proof of Proposition 9.6 we can show

Proposition 10.5 *Let λ be a weight such that $\lambda(K_\beta) \in \pm q^{\mathbb{N}}$ for all short $\beta \in \Phi^+$ and $\lambda(K_\gamma) \in \pm q^{1+2\mathbb{Z}}$ for all long $\gamma \in \Phi^+$. Let $\mathbf{b} = (b_1, \dots, b_n) \in (\mathbb{C}^*)^n$. E_{α_2} acts injectively on the U_q-module $\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$ if and only if $b_2 \notin \pm q^{\mathbb{Z}}$. Let $i \in \{3, \dots, n\}$. Then E_{α_i} acts injectively on the module $\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$ if and only if $b_i \notin \pm q^{\mathbb{Z}}$ and F_{α_i} acts injectively on $\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$ if and only if $b_{i-1} \notin \pm q^{\mathbb{Z}}$.*

Proof. By Proposition 4.10 and Corollary 4.20 a root vector acts injectively on the U_q-module

$$\varphi_{F_\Sigma, (b_1, \dots, b_n)}.L(\lambda)_{F_\Sigma}$$

if and only if it acts injectively on

$$\varphi_{F_\Sigma, (\varepsilon_1 q^{i_1} b_1, \dots, \varepsilon_n q^{i_n} b_n)}.L(\lambda)_{F_\Sigma}$$

for any $i_1, \dots, i_n \in \mathbb{Z}$ and $\varepsilon_1, \dots, \varepsilon_n \in \{\pm 1\}$.

Assume there exists a $0 \neq v \in \varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$ such that $F_{\alpha_i} v = 0$. We have $v = F_{\beta_1}^{a_1} \cdots F_{\beta_n}^{a_n} \otimes v'$ for some $a_1, \dots, a_n \in \mathbb{Z}_{\leq 0}$ and some $v' \in L(\lambda)$. $F_{\alpha_i} v = 0$ implies

$$0 = \varphi_{F_\Sigma, \mathbf{b}}(F_{\alpha_i}) F_{\beta_1}^{a_1} \cdots F_{\beta_n}^{a_n} \otimes v' = F_{\beta_1}^{a_1} \cdots F_{\beta_n}^{a_n} \otimes \varphi_{F_\Sigma, \mathbf{c}}(F_{\alpha_i}) v'$$

where $\mathbf{c} = (q^{a_1} b_1, \dots, q^{a_n} b_n)$. So there exists a $v' \in L(\lambda)$ such that $\varphi_{F_\Sigma, \mathbf{c}}(F_{\alpha_i}) v' = 0$. That is

$$\left(c_{i-1} F_{\alpha_i} + \frac{c_{i-1} - c_{i-1}^{-1}}{q - q^{-1}} F_{\beta_{i-1}}^{-1} F_{\beta_i} \right) v' = 0$$

or equivalently

$$\left(F_{\beta_{i-1}} F_{\alpha_i} + c_{i-1}^{-1} \frac{c_{i-1} - c_{i-1}^{-1}}{q - q^{-1}} F_{\beta_i} \right) v' = 0.$$

Let $r \in \mathbb{N}$ be such that $F_{\alpha_i}^{(r)} v' \neq 0$ and $F_{\alpha_i}^{(r+1)} v' = 0$ (possible since $\lambda(K_{\alpha_i}) \in \pm q^{\mathbb{N}}$ so $-\alpha_i \in F_{L(\lambda)}$). So the above being equal to zero implies

$$\begin{aligned} 0 =& F_{\alpha_i}^{(r)} \left(F_{\beta_{i-1}} F_{\alpha_i} + c_{i-1}^{-1} \frac{c_{i-1} - c_{i-1}^{-1}}{q - q^{-1}} F_{\beta_i} \right) v' \\ =& \left([r] F_{\beta_i} F_{\alpha_i}^{(r)} + q^{-r} \frac{1 - c_{i-1}^{-2}}{q - q^{-1}} F_{\beta_i} F_{\alpha_i}^{(r)} \right) v' \\ =& \left([r] + q^{-r} \frac{1 - c_{i-1}^{-2}}{q - q^{-1}} \right) F_{\beta_i} F_{\alpha_i}^{(r)} v'. \end{aligned}$$

Since $F_{\beta_i} F_{\alpha_i}^{(r)} v' \neq 0$ this is equivalent to

$$0 = q^r - q^{-r} + q^{-r} - q^{-r} c_{i-1}^{-2} = q^r - q^{-r} c_{i-1}^{-2}$$

or equivalently $c_{i-1} = \pm q^{-r}$.

The other claims are shown similarly. $\square$

Proposition 10.6 *Let λ be a weight such that $\lambda(K_\beta) \in \pm q^{\mathbb{N}}$ for all short $\beta \in \Phi^+$ and $\lambda(K_\gamma) \in \pm q^{1+2\mathbb{Z}}$ for all long $\gamma \in \Phi^+$. Let $\mathbf{b} = (b_1, \dots, b_n) \in (\mathbb{C}^*)^n$. Then $F_{\alpha_1 + 2\alpha_2}$ acts injectively on the U_q-module $\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$.*

Proof. We can show similarly to the above calculations in this section that

$$\varphi_{F_\Sigma, \mathbf{b}}(F_{\alpha_1 + 2\alpha_2}) = b_2^2 F_{\alpha_1 + 2\alpha_2} + (1 - q^2) b_2^2 b_1^{-2} \frac{b_1^2 - b_1^{-2}}{q^2 - q^{-2}} F_{\beta_1}^{-1} F_{\beta_2}^{(2)}.$$

By Proposition 4.10 and Corollary 4.20 a root vector acts injectively on the U_q-module

$$\varphi_{F_\Sigma, (b_1, \dots, b_n)}.L(\lambda)_{F_\Sigma}$$

if and only if it acts injectively on

$$\varphi_{F_\Sigma, (\varepsilon_1 q^{i_1} b_1, \dots, \varepsilon_n q^{i_n} b_n)}.L(\lambda)_{F_\Sigma}$$

for any $i_1, \dots, i_n \in \mathbb{Z}$ and $\varepsilon_1, \dots, \varepsilon_n \in \{\pm 1\}$.

Assume there exists a $0 \neq v \in \varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ such that $F_{\alpha_1+2\alpha_2} v = 0$. We have $v = F_{\beta_1}^{a_1} \cdots F_{\beta_n}^{a_n} \otimes v'$ for some $a_1, \dots, a_n \in \mathbb{Z}$ and some $v' \in L(\lambda)$. So $F_{\alpha_1+2\alpha_2} v = 0$ implies

$$0 = \varphi_{F_\Sigma,\mathbf{b}}(F_{\alpha_1+2\alpha_2}) F_{\beta_1}^{a_1} \cdots F_{\beta_n}^{a_n} \otimes v' = F_{\beta_1}^{a_1} \cdots F_{\beta_n}^{a_n} \otimes \varphi_{F_\Sigma,\mathbf{c}}(F_{\alpha_1+2\alpha_2}) v'$$

where $\mathbf{c} = (q^{a_1} b_1, \dots, q^{a_n} b_n)$. So there exists a $v' \in L(\lambda)$ and $a_1, \dots, a_n \in \mathbb{Z}$ such that for $\mathbf{c} = (q^{a_1} b_1, \dots, q^{a_n} b_n)$, $\varphi_{F_\Sigma,\mathbf{c}}(F_{\alpha_1+2\alpha_2}) v' = 0$. That is

$$\left(c_2^2 F_{\alpha_1+2\alpha_2} + (1-q^2) c_2^2 c_1^{-2} \frac{c_1^2 - c_1^{-2}}{q^2 - q^{-2}} F_{\beta_1}^{-1} F_{\beta_2}^{(2)} \right) v' = 0$$

or equivalently

$$F_{\beta_1} F_{\alpha_1+2\alpha_2} v' + (1-q^2) c_1^{-2} \frac{c_1^2 - c_1^{-2}}{q^2 - q^{-2}} F_{\beta_2}^{(2)} v' = 0.$$

So to prove our claim it is enough to prove that

$$\left(F_{\beta_1} F_{\alpha_1+2\alpha_2} + (1-q^2) c_1^{-2} \frac{c_1^2 - c_1^{-2}}{q^2 - q^{-2}} F_{\beta_2}^{(2)} \right) v' \neq 0$$

for any $v' \in L(\lambda)$ and any $c_1 \in \mathbb{C}^*$.

So let $v' \in L(\lambda)$ and let $c_1 \in \mathbb{C}^*$. Let $r \in \mathbb{N}$ be such that $E_{\alpha_2}^{(r)} v' \neq 0$ and $E_{\alpha_2}^{(r+1)} v' = 0$ (possible since $L(\lambda)$ is a highest weight module). It is straightforward to show that for $a \in \mathbb{N}$:

$$[E_{\alpha_2}^{(a)}, F_{\alpha_1+2\alpha_2}] = q^{-a+1} [2] F_{\beta_2} E_{\alpha_2}^{(a-1)} K_{\alpha_2}^{-1} + q^{4-2a} F_{\beta_1} E_{\alpha_2}^{(a-2)} K_{\alpha_2}^{-2}$$

and

$$[E_{\alpha_2}^{(a)}, F_{\beta_2}^{(2)}] = q^{2-a} [2] F_{\beta_2} F_{\beta_1} E_{\alpha_2}^{(a-1)} K_{\alpha_2}^{-1} + q^{3-2a} [2] F_{\beta_1}^2 E_{\alpha_2}^{(a-2)} K_{\alpha_2}^{-2}.$$

Using this we get

$$\begin{aligned} & E_{\alpha_2}^{(r+2)} \left(F_{\beta_1} F_{\alpha_1+2\alpha_2} + (1-q^2) c_1^{-2} \frac{c_1^2 - c_1^{-2}}{q^2 - q^{-2}} F_{\beta_2}^{(2)} \right) v' \\ =& \left(q^{-2r} + q^{-1-2r} [2] (1-q^2) c_1^{-2} \frac{c_1^2 - c_1^{-2}}{q^2 - q^{-2}} \right) F_{\beta_1}^2 E_{\alpha_2}^{(r)} K_{\alpha_2}^{-2} v' \\ =& q^{-2r} c_1^{-4} F_{\beta_1}^2 E_{\alpha_2}^{(r)} K_{\alpha_2}^{-2} v' \\ \neq& 0 \end{aligned}$$

since F_{β_1} acts injectively on $L(\lambda)$. Thus

$$\left(F_{\beta_1} F_{\alpha_1+2\alpha_2} + (1-q^2) c_1^{-2} \frac{c_1^2 - c_1^{-2}}{q^2 - q^{-2}} F_{\beta_2}^{(2)} \right) v' \neq 0. \qquad \square$$

Theorem 10.7 *Let λ be a weight such that $\lambda(K_\beta) \in \pm q^{\mathbb{N}}$ for all short $\beta \in \Phi$ and $\lambda(K_\gamma) \in \pm q^{1+2\mathbb{Z}}$ for all long $\gamma \in \Phi$. Let $\mathbf{b} = (b_1, \dots, b_n) \in (\mathbb{C}^*)^n$. Then the U_q-module $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ is simple and torsion free if and only if $b_i \notin \pm q^{\mathbb{Z}}$, $i = 2, \dots, n$ and $b_1^2 b_2 \cdots b_n \notin \pm q^{\mathbb{Z}}$.*

Proof. Let $i \in \{2, \dots, n\}$. By Proposition 10.5, E_{α_i} acts injectively on $\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$ if and only if $b_i \notin \pm q^{\mathbb{Z}}$. If $\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$ is torsion free then every root vector acts injectively. So $\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$ being torsion free implies $b_i \notin \pm q^{\mathbb{Z}}$.

Let $\Sigma' = \{\beta_1', \dots, \beta_n'\}$ denote the set of commuting roots with $\beta_1' = \alpha_1 + \alpha_2$, $\beta_2' = \alpha_1 + 2\alpha_2$, $\beta_j' = \alpha_1 + 2\alpha_2 + \alpha_3 + \dots + \alpha_j$, $j = 3, \dots, n$. Let $F'_{\beta_1'} := T_{s_1}(F_{\alpha_2}) = F_{\beta_2}$, $F'_{\beta_2'} := T_{s_1} T_{s_2}(F_{\alpha_1}) = F_{\alpha_1 + 2\alpha_1}$, $F'_{\beta_j'} := T_{s_1} \cdots T_{s_n} T_{s_1} \cdots T_{s_{j-2}}(F_{\alpha_{j-1}}) = T_{s_2}(F_{\beta_j}) = F_{\alpha_1 + 2\alpha_2 + \alpha_3 + \dots + \alpha_j}$, $j = 3, \dots, n$ (in this case we actually have $F'_{\beta_j'} = F_{\beta_j'}$) and $F_{\Sigma'}$ the Ore subset generated by $F'_{\beta_1'}, \dots, F'_{\beta_n'}$. Similarly to the above calculations in this section we can show that for $\mathbf{c} \in (\mathbb{C}^*)^n$

$$\varphi_{F_{\Sigma'}, \mathbf{c}}(F_{\alpha_2}) = c_n^{-1} \cdots c_3^{-1} c_2^{-2} \left(F_{\alpha_2} + q[2] c_1^{-1} \frac{c_1 - c_1^{-1}}{q - q^{-1}} (F'_{\beta_1'})^{-1} F'_{\beta_2'} \right).$$

Let $v \in L(\lambda)$ and let $r \in \mathbb{N}$ be such that $F_{\alpha_2}^{(r)} v \neq 0$ and $F_{\alpha_2}^{(r+1)} v = 0$ (possible since $\lambda(K_{\alpha_2}) \in \pm q^{\mathbb{N}}$). Then we see like in the proof of Proposition 10.5 that $\varphi_{F_{\Sigma'}, \mathbf{c}}(F_{\alpha_2}) v = 0$ if and only if $c_1 = \pm q^{-r}$ thus $\varphi_{F_{\Sigma'}, \mathbf{c}}.L(\lambda)_{F_{\Sigma'}}$ is not torsion free whenever $c_1 \in \pm q^{\mathbb{Z}}$ by Proposition 4.10 and Corollary 4.20.

Set $f(\mathbf{b}) = (b_1^2 b_2 \cdots b_n, b_1^{-1} b_3^{-1} \cdots b_n, b_3, \dots, b_n)$. Then by Lemma 5.3

$$\left(\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}\right)^{ss} \cong \left(\varphi_{F_{\Sigma'}, f(\mathbf{b})}.L(\lambda)_{F_{\Sigma'}}\right)^{ss}.$$

If $\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$ is torsion free then it is simple so

$$\begin{aligned} \varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma} &\cong \left(\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}\right)^{ss} \\ &\cong \left(\varphi_{F_{\Sigma'}, f(\mathbf{b})}.L(\lambda)_{F_{\Sigma'}}\right)^{ss} \\ &\cong \varphi_{F_{\Sigma'}, f(\mathbf{b})}.L(\lambda)_{F_{\Sigma'}}. \end{aligned}$$

We see that $\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$ being torsion free implies $b_1^2 b_2 \cdots b_n \notin \pm q^{\mathbb{Z}}$.

Now assume $b_i \notin \pm q^{\mathbb{Z}}$, $i = 2, \dots, n$ and $b_1^2 b_2 \cdots b_n \notin \pm q^{\mathbb{Z}}$. By Proposition 10.5 and Proposition 9.7 E_{α_i} and F_{α_i}, $i = 3, \dots, n$ act injectively on all composition factors of $\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$.

Let L_1 be a simple submodule of $\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$ and let L_2 be a simple submodule of $\varphi_{F_{\Sigma'}, f(\mathbf{b})}.L(\lambda)_{F_{\Sigma'}}$. By Proposition 10.6, $F_{\alpha_1 + 2\alpha_2}$ acts injectively on $\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}$. Now clearly $\{-\alpha_1 - \alpha_2, -\alpha_1 - 2\alpha_2, \alpha_3, \dots, \alpha_n\} \subset T_{L_1} \cap T_{L_2}$ so $C(L_1) \cap C(L_2)$ generates Q. This implies that $C(L_1) - C(L_2) = Q$. Since $\left(\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}\right)^{ss} \cong \left(\varphi_{F_{\Sigma'}, f(\mathbf{b})}.L(\lambda)_{F_{\Sigma'}}\right)^{ss}$ we have $\operatorname{wt} L_k \subset q^Q (\mathbf{b}^{-1})^\Sigma \lambda$, $k = 1, 2$. Choose $\mu_1, \mu_2 \in Q$ such that $q^{\mu_1} (\mathbf{b}^{-1})^\Sigma \lambda \in \operatorname{Supp}_{\text{ess}}(L_1)$ and $q^{\mu_2} (\mathbf{b}^{-1})^\Sigma \lambda \in \operatorname{Supp}_{\text{ess}}(L_2)$. Then obviously $q^{C(L_1) + \mu_1} (\mathbf{b}^{-1})^\Sigma \lambda \subset \operatorname{Supp}_{\text{ess}}(L_1)$ and $q^{C(L_2) + \mu_2} (\mathbf{b}^{-1})^\Sigma \lambda \subset \operatorname{Supp}_{\text{ess}}(L_2)$. By the above $q^{C(L_1) + \mu_1} (\mathbf{b}^{-1})^\Sigma \lambda \cap q^{C(L_2) + \mu_2} (\mathbf{b}^{-1})^\Sigma \lambda \neq \emptyset$ so $\operatorname{Supp}_{\text{ess}}(L_1) \cap \operatorname{Supp}_{\text{ess}}(L_2) \neq \emptyset$. Let $\nu \in \operatorname{Supp}_{\text{ess}}(L_1) \cap \operatorname{Supp}_{\text{ess}}(L_2)$. By Proposition 5.6, L_1 and L_2 are admissible of the same degree as $L(\lambda)$. So we have as $(U_q)_0$-modules (using that $(L_1)_\nu$ and $(L_2)_\nu$ are simple $(U_q)_0$-modules by Theorem 2.7)

$$\begin{aligned} (L_1)_\nu = \left(\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}\right)_\nu &\cong \left(\left(\varphi_{F_\Sigma, \mathbf{b}}.L(\lambda)_{F_\Sigma}\right)_\nu\right)^{ss} \\ &\cong \left(\left(\varphi_{F_{\Sigma'}, f(\mathbf{b})}.L(\lambda)_{F_{\Sigma'}}\right)_\nu\right)^{ss} \cong \left(\varphi_{F_{\Sigma'}, f(\mathbf{b})}.L(\lambda)_{F_{\Sigma'}}\right)_\nu = (L_2)_\nu. \end{aligned}$$

By Theorem 2.7 this implies $L_1 \cong L_2$.

Let $\Sigma'' = \{\beta_1'', \dots, \beta_n''\}$ denote the set of commuting roots with $\beta_1'' = \alpha_1 + 2\alpha_2$, $\beta_2'' = \alpha_2$, $\beta_j'' = \alpha_1 + 2\alpha_2 + \alpha_3 + \dots + \alpha_j$, $j = 3, \dots, n$. Let $F''_{\beta_1''} :=$

$T_{s_1}T_{s_2}(F_{\alpha_1})$, $F''_{\beta''_2} := F_{\alpha_2}$, $F''_{\beta''_j} := T_{s_2}T_{s_1}T_{s_2}T_{s_3}\cdots T_{s_{j-1}}(F_{\alpha_j}) = T_{s_1}T_{s_2}(F_{\beta_j})$, $j = 3, \dots, n$ and $F_{\Sigma''}$ the Ore subset generated by $F''_{\beta''_1}, \dots, F''_{\beta''_n}$. Note that $F''_{\beta''_j} = T_{s_1}T_{s_2}(F_{\beta_j})$ for all $j \in \{1, \dots, n\}$. The root vectors $F''_{\beta''_1}, \dots, F''_{\beta''_n}$ act injectively on ${}^{\overline{s_2 s_1}}L(\lambda)$. By Theorem 5.12 and Proposition 5.5 $L(\lambda)$ is a submodule of $\left(\varphi_{F_{\Sigma''},\mathbf{d}}.({}^{\overline{s_2 s_1}}L(\lambda))_{F_{\Sigma''}}\right)^{ss}$ for some $\mathbf{d} \in (\mathbb{C}^*)^n$. Then by Lemma 5.3

$$(\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma})^{ss} \cong \left(\varphi_{F_{\Sigma''},g(\mathbf{b})\mathbf{d}}.({}^{\overline{s_2 s_1}}L(\lambda))_{F_{\Sigma''}}\right)^{ss}$$

for some $g(\mathbf{b}) \in (\mathbb{C}^*)^n$.

Observe that for $a_1, \dots, a_n \in \mathbb{N}$:

$$\begin{aligned}
\varphi_{F_{\Sigma''},(q^{a_1},\dots,q^{a_n})}&(-K_{\alpha_1}E_{\alpha_1}) \\
&=\varphi_{F_{\Sigma''},(q^{a_1},\dots,q^{a_n})}(T_{s_1}T_{s_2}(F_{\alpha_1+2\alpha_2})) \\
&=\left(F''_{\beta''_1}\right)^{-a_1}\cdots\left(F''_{\beta''_n}\right)^{-a_n} T_{s_1}T_{s_2}(F_{\alpha_1+2\alpha_2})\left(F''_{\beta''_n}\right)^{a_n}\cdots\left(F''_{\beta''_1}\right)^{a_1} \\
&=T_{s_1}T_{s_2}\left(F_{\beta_1}^{-a_n}\cdots F_{\beta_n}^{-a_n}F_{\alpha_1+2\alpha_2}F_{\beta_n}^{a_n}\cdots F_{\beta_1}^{a_1}\right) \\
&=T_{s_1}T_{s_2}\left(\varphi_{F_\Sigma,(q^{a_1},\dots,q^{a_n})}(F_{\alpha_1+2\alpha_2})\right).
\end{aligned}$$

Since $\varphi_{F_{\Sigma''},\mathbf{c}}(-K_{\alpha_1}E_{\alpha_1})$ and $T_{s_1}T_{s_2}\left(\varphi_{F_\Sigma,\mathbf{c}}(F_{\alpha_1+2\alpha_2})\right)$ are both Laurent polynomial in $\mathbf{c}$ we get by Lemma 4.8 that $\varphi_{F_{\Sigma''},\mathbf{c}}(-K_{\alpha_1}E_{\alpha_1}) = T_{s_1}T_{s_2}\left(\varphi_{F_\Sigma,\mathbf{c}}(F_{\alpha_1+2\alpha_2})\right)$ for any $\mathbf{c} \in (\mathbb{C}^*)^n$. $T_{s_1}T_{s_2}\left(\varphi_{F_\Sigma,\mathbf{c}}(F_{\alpha_1+2\alpha_2})\right)$ acts injectively on ${}^{\overline{s_2 s_1}}L(\lambda)$ for any $\mathbf{c} \in (\mathbb{C}^*)^n$ by Proposition 10.6. This implies that $-K_{\alpha_1}E_{\alpha_1}$ acts injectively on $\varphi_{F_{\Sigma''},g(\mathbf{b})\mathbf{d}}.({}^{\overline{s_2 s_1}}L(\lambda))_{F_{\Sigma''}}$ and this implies that E_{α_1} acts injectively.

Let L_3 be a simple submodule of $\varphi_{F_{\Sigma''},g(\mathbf{b})\mathbf{d}}.({}^{\overline{s_2 s_1}}L(\lambda))_{F_{\Sigma''}}$. We see that $\{-\alpha_2, -\alpha_1-2\alpha_2, \alpha_3, \dots, \alpha_n\} \subset T_{L_3} \cap T_{L_2}$ so $C(L_2) \cap C(L_3)$ generates Q ($\{\alpha_3, \dots, \alpha_n\} \subset T_{L_3}$ because of Proposition 9.7 and the fact that L_3 is a composition factor of $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$). Arguing as above this implies that $L_2 \cong L_3$. We have shown that $L_1 \cong L_2 \cong L_3$. Above we have shown that E_{α_1} acts injectively on L_3, F_{α_2} acts injectively on L_2 and $F_{\alpha_1}, E_{\alpha_2}, F_{\alpha_i}, E_{\alpha_i}$, $i = 3, \dots, n$ act injectively on L_1. In conclusion we have shown that all root vectors act injectively on the simple submodule L_1 of $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ thus $\operatorname{wt} L_1 = \operatorname{Supp}_{\text{ess}}(L_1) = q^Q(\mathbf{b}^{-1})^\Sigma\lambda$ and therefore $L_1 = \varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$. This shows that $\varphi_{F_\Sigma,\mathbf{b}}.L(\lambda)_{F_\Sigma}$ is simple and torsion free with our assumptions on $\mathbf{b}$. ☐

References

[AM15] Henning Haahr Andersen and Volodymyr Mazorchuk, *Category $\mathcal{O}$ for quantum groups*, J. Eur. Math. Soc. (JEMS) **17** (2015), no. 2, 405–431. MR 3317747

[And03] Henning Haahr Andersen, *Twisted Verma modules and their quantized analogues*, Combinatorial and geometric representation theory (Seoul, 2001), Contemp. Math., vol. 325, Amer. Math. Soc., Providence, RI, 2003, pp. 1–10. MR 1988982 (2005b:17025)

[APW91] Henning Haahr Andersen, Patrick Polo, and Ke Xin Wen, *Representations of quantum algebras*, Invent. Math. **104** (1991), no. 1, 1–59. MR 1094046 (92e:17011)

[DP93] C. DeConcini and C. Procesi, *"Quantum groups" in: D-modules, representation theory, and quantum groups*, Lecture Notes in Mathematics, vol. 1565, Springer-Verlag, Berlin, 1993, Lectures given at the Second C.I.M.E. Session held in Venice, June 12–20, 1992. MR 1288993 (95b:17003)

[Fer90] S. L. Fernando, *Lie algebra modules with finite-dimensional weight spaces. I*, Trans. Amer. Math. Soc. **322** (1990), no. 2, 757–781. MR 1013330 (91c:17006)

[Jan96] Jens Carsten Jantzen, *Lectures on quantum groups*, Graduate Studies in Mathematics, vol. 6, American Mathematical Society, Providence, RI, 1996. MR 1359532 (96m:17029)

[Jos95] Anthony Joseph, *Quantum groups and their primitive ideals*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 29, Springer-Verlag, Berlin, 1995. MR 1315966 (96d:17015)

[Lam01] T. Y. Lam, *A first course in noncommutative rings*, second ed., Graduate Texts in Mathematics, vol. 131, Springer-Verlag, New York, 2001. MR 1838439 (2002c:16001)

[LS91] Serge Levendorskiĭ and Yan Soibelman, *Algebras of functions on compact quantum groups, Schubert cells and quantum tori*, Comm. Math. Phys. **139** (1991), no. 1, 141–170. MR 1116413 (92h:58020)

[Lus90] George Lusztig, *Quantum groups at roots of* 1, Geom. Dedicata **35** (1990), no. 1-3, 89–113. MR 1066560 (91j:17018)

[Mat00] Olivier Mathieu, *Classification of irreducible weight modules*, Ann. Inst. Fourier (Grenoble) **50** (2000), no. 2, 537–592. MR 1775361 (2001h:17017)

[Pap94] Paolo Papi, *A characterization of a special ordering in a root system*, Proc. Amer. Math. Soc. **120** (1994), no. 3, 661–665. MR 1169886 (94e:20056)

[Ped15a] Dennis Hasselstrøm Pedersen, *Irreducible quantum group modules with finite dimensional weight spaces. I*, arXiv:1504.07042, 2015.

[Ped15b] ———, *Twisting functors for quantum group modules*, arXiv:1504.07039, 2015.